

# **Ichneumonidae (Hymenoptera)**

## **As Biological Control Agents Of Pests**



## **A Bibliography**

**Hassan Ghahari**

Department of Entomology, Islamic Azad University, Science & Research Campus, P. O. Box 14515/775, Tehran – Iran; h\_ghahhari@yahoo.com

### **Preface**

The Ichneumonidae is one of the most species rich families of all organisms with an estimated 60000 species in the world (Townes, 1969). Even so, many authorities regard this figure as an underestimate! (Gauld, 1991). An estimated 12100 species of Ichneumonidae occur in the Afrotropical region (Africa south of the Sahara and including Madagascar) (Townes & Townes, 1973), of which only 1927 have been described (Yu, 1998). *This means that roughly 16% of the afrotropical ichneumonids are known to science!* These species comprise 338 genera. The family Ichneumonidae is currently split into 37 subfamilies (including, Acaenitinae; Adelognathinae; Agriotypinae; Alomyinae; Anomaloninae; Banchinae; Brachycyrtinae; Campopleginae; Collyrinae; Cremastinae; Cryptinae; Ctenopelmatinae;

Diplazontinae; Eucerotinae; Ichneumoninae; Labeninae; Lycorininae; Mesochorinae; Metopiinae; Microleptinae; Neorhacodinae; Ophioninae; Orthopelmatinae; Orthocentrinae; Oxytorinae; Paxylomatinae; Phrudinae; Phygadeuontinae; Pimplinae; Rhyssinae; Stilbopinae; Tersilochinae; Tryphoninae; Xoridinae) (Yu, 1998).

The Ichneumonidae, along with other groups of parasitic Hymenoptera, are supposedly no more species rich in the tropics than in the Northern Hemisphere temperate regions (Owen & Owen, 1974; Janzen, 1981; Janzen & Pond, 1975), although a number of hymenopteran families, for example the Chalcididae (Hespenheide, 1979) and Encyrtidae (Noyes 1989b) exhibit an increase in species richness with a decrease in latitude. Other hymenopteran taxa such as sawflies (Symphyta), gall-forming Cynipidae and bees (Apoidea) peak in species richness at mid or high latitudes (Michener 1979; Noyes 1989b; Kouki *et al.* 1994). Considerable debate has centred on the apparent species richness anomaly exhibited by a number of hymenopteran parasitoid taxa in the tropics (see Morrison *et al.* 1978; Gauld, 1991; Gauld & Gaston, 1994).

Ichneumonids utilise a diverse array of insects and arachnids as their hosts and play an essential role in the normal functioning of most ecosystems, underlining the need to inventory their diversity. Ichneumonids have been used successfully as biocontrol agents and given the largely undocumented fauna there is a huge potential for their utilisation in managed biocontrol programmes (Gupta, 1991). Comprehensive quantitative biodiversity surveys will enable the identification of hotspots of species richness and endemism; essential base line data that will enable informed future conservation management decisions.

Identifying characteristics for the family Ichneumonidae include:

- Antennae apparently with 16 or more segments.
- Hind trochanters 2-segmented.
- Two recurrent veins.
- Usually larger than braconids; quite variable in color.
- Females often with long ovipositors.

Additional information:

- Species in the family Ichneumonidae are separated from those in Braconidae by having two, rather than one or zero recurrent veins.
- Ichneumons are important internal parasitoids of other insects. Common hosts are larvae and pupae of Coleoptera, Hymenoptera, and Lepidoptera.

## **Bibliography on Ichneumonidae**

### **A**

**Abbas, M.S.T. (1988).** Biological and ecological studies on *Diadegma semiclausum* Hellen (Hym.: Ichneumonidae), a larval parasite of the diamond back moth, *Plutella xylostella* (L.) (Lep.: Plutellidae) in Egypt. *Anzeiger für Schadlingskunde, Pflanzenschutz, Umweltschutz* 61 (1): 1 - 2.

- Abbas, M.S.T., Abuo-Zeid, N.A. and Megahed, M.M. (1993).** On the natural enemies of the potato tuber moth *Phthorimaea operculella* in Egypt. Egyptian Journal of Agricultural Research 71 (4): 943 - 950.
- Abbas, M.S.T. and El Deeb, Y.A.A. (1993).** On the natural enemies of the major pests infesting cotton in Egypt. Egyptian Journal of Agricultural Research, publ. 1996, 71 (1): 131 - 138.
- Abdinbekova, A.A. and Mustafina, K.M. (1988).** Characteristics of the biology and ecology of *Hyposoter didymator* Thund. (Ichneumonidae) - a parasitoid of the cotton moth *Heliothis armigera* Hb. (Noctuidae) in Azerbaidzhan. Izvestiya Akademii Nauk Azerbaidzhanskoi SSR. Seriya Biologicheskikh Nauk 3: 75 - 83.
- Adashkevich, B.P. and Rashidov, M.I. (1986).** Biological control of the cotton bollworm on vegetable crops. Zashchita Rastenii 6: 51 - 52.
- Adiyaman, N. and Aktumsek, A. (1996).** The effects of low temperature on fecundity of pupae and adults of female *Pimpla turionellae* L. (Hymenoptera: Ichneumonidae). Turkish Journal of Zoology 20: SUPPL, 1 - 5.
- Aeschlimann, J.P. (1989).** Revision of the western Palaearctic species of the genus *Hypsicera* Latreille (Hymenoptera: Ichneumonidae). Annales de la Societe Entomologique de France 25 (1): 33 - 39.
- Agoze, M.E., Drezen, J.M., Renault, S. and Periquet, G. (1994).** Analysis of the reproductive potential of diploid males in the wasp *Diadromus pulchellus* (Hymenoptera: Ichneumonidae). Bulletin of Entomological Research 84 (2): 213 - 218.
- Agoze, M.M.E. (1987).** Reproductive activity of two parasitoids: *Diadromus pulchellus* (Wsm.) and *Bruchocida vuilleti* [*Eulpelmus vuilleti*] (Crawford) during a limited time of their host [*Acrolepiopsus assectella* and *Callosobruchus maculatus*, resp.] presence. Bulletin de la Societe Entomologique d'Egypte, publ. 1989, 67: 113 - 121.
- Ahmed, K., Khalique, F. and Afzal, M. (1989).** Effect of agronomic factors on the incidence of *Heliothis armigera* (Hubn.) and its parasite *Campoletis chlorideae* (U.) in chickpea field. Pakistan Journal of Scientific and Industrial Research 32 (10): 694 - 697.
- Ahmed, K.N. and Khan, A.R. (1995).** Biological note on *Campoletis chlorideae* Uchida (Hymenoptera: Ichneumonidae). Bangladesh Journal of Zoology 23 (2): 243 - 244.
- Ajayi, O., Uvah, I.I., Fussell, L.K. and Weder, J. (1989).** Review of research on millet entomology in Nigeria: 1977 - 1987. Proceedings of the Regional Pearl Millet Improvement Workshop, IAR, Samaru, Nigeria, 15-19 August 1988. 21 - 30.
- Akol, A.M., Sithanatham, S., Njagi, P.G.N., Varela, A. and Mueke, J.M. (2002).** Relative safety of sprays of two neem insecticides to *Diadegma mollipla* (Holmgren), a parasitoid of the diamondback moth: effects on adult longevity and foraging behaviour. Crop Protection 21 (9): 853 - 859.
- Akre, R.D. and Myhre, E.A. (1994).** Nesting biology of *Dolichovespula norvegicoides* (Hymenoptera: Vespidae). Entomological News 105 (1): 39 - 46.

- Aktumsek, A. (1996).** Effect of host and sex on fatty acid composition in the parasitoid *Itoplectis maculator* F. (Hymenoptera: Ichneumonidae). Turkish Journal of Zoology 20: SUPPL, 7 - 10.
- Al Ayedh, H.Y., Kondratieff, B.C., Blodgett, S.L. and Peairs, F.B. (1996).** Evaluation of hymenopterous biological control agents of the alfalfa weevil larvae *Hypera postica* (Coleoptera: Curculionidae) in northcentral Colorado. Journal of the Kansas Entomological Society 69 (4): 326 - 336.
- Al Saffar, Z.Y. and Aldrich, J.C. (1997).** Factors influencing the survival of *Pontania proxima* that attack crack willow *Salix fragilis*. Biology and Environment: Proceedings of the Royal Irish Academy, Section B. 97 (3): 219 - 223.
- Ali, M.I., Islam, A. and Kabir, S.M.H. (1995).** Biological control of insect and mite pests in important agricultural crops of Bangladesh: a review. Journal of the Asiatic Society of Bangladesh Science 21 (2): 149 - 208.
- Ali, M.I. and Karim, M.A. (1995).** Host range, abundance and natural enemies of diamondback moth in Bangladesh. Bangladesh Journal of Entomology 5 (1-2): 25 - 32.
- Aliyev, A.A.O. (1999).** Fauna of the subfamily Ichneumoninae (Hymenoptera: Ichneumonidae) of Azerbaijan with new records. Turkish Journal of Zoology 23 (1): 1 - 12.
- Alizadeh, I., Safaralizadeh, M.H. and Shayestef, N. (2000).** Biology of alfalfa weevil (*Hypera postica* Gyll.) and identification of its' parasitoids and dominant species in Maku and Chalderan regions. Proceedengs of 14th Iranian Plant Protection Congress, p. 27.
- Allen, G.R. (1990).** *Uraba lugens* Walker (Lepidoptera: Noctuidae): larval survival and parasitoid biology in the field in South Australia. Journal of the Australian Entomological Society 29 (4): 301 - 312.
- Aluja, M., Guillen, J., Liedo, P., Cabrera, M., Rios, E., De La Rosa, G., Celedonio, H. and Mota, D. (1990).** Fruit infesting tephritids (Diptera: Tephritidae) and associated parasitoids in Chiapas, Mexico. Entomophaga 35 (1): 39 - 48.
- Amend, J. and Basedow, T. (1997).** Combining release/establishment of *Diadegma semiclausum* (Hellen) (Hym.: Ichneumonidae) and *Bacillus thuringiensis* Berl. for control of *Plutella xylostella* (L.) (Lep.: Yponomeutidae) and other lepidopteran pests in the Cordillera Region of Luzon (Philippines). Journal of Applied Entomology 121 (6): 337 - 342.
- Ananthkrishnan, T.N. (1995).** Biological control of social forest and plantation crop insects. Science Publishers, Inc.; New Hampshire, USA; Lebanon, xii + 225 pp.
- Anciso, J.R. and Quick, T.C. (1990).** Parasitoids of the diamondback moth, *Plutella xylostella* (L.) in the lower Rio Grande Valley, Texas. Journal of the Rio Grande Valley Horticultural Society 43: 89 - 90.
- Andrei-Ruiz, M.C. and Villemant, C. (1999).** Biodiversity of the Ichneumonoidea (Hymenoptera) in the green oak forest of Fango (Corsica). Integrated protection of

- Quercus* spp. forests. Proceedings of the meeting held at Rabat-Sale, Morocco, 26 - 29 October, 1998. Bulletin OILB SROP 22 (3): 151 - 156.
- Anento, J.L., Kolarov, J. and Selfa, J. (2002).** First record for Europe (Alicante, Spain) of *Temelucha dorsonigra* (Hedwig, 1957) (Hymenoptera: Ichneumonidae: Cremastinae). Boletín de la Real Sociedad Española de Historia Natural, Sección Biológica 97 (1-4): 71 - 73.
- Anento, J.L. and Selfa, J. (1996).** New data on Spanish Acaenitinae (Hymenoptera: Ichneumonidae). Nouvelle Revue d'Entomologie 13 (4): 333 - 336.
- Ang, B.N. and Kok, L.T. (1995).** Seasonal mortality of *Cassida rubiginosa* (Coleoptera: Chrysomelidae) due to incomplete development and parasitism in southwestern Virginia. Journal of Entomological Science 30 (1): 9 - 17.
- Anis Rahman, U.R. and Ellington, J.J. (1999).** A survey of the insect parasitoids of alfalfa weevil *Hypera postica* (Gyllenhal) in the Mesilla valley, New Mexico, USA. Sarhad Journal of Agriculture 15 (6): 603 - 606.
- Anonymous. (1987).** *Acigona ignefusalis* (Hampson). Pyralidae. Millet and sorghum borer. Institut de Recherches Agronomiques Tropicales et des Cultures Vivrières; Montpellier; France, 2 pp.
- Anonymous. (1987).** Biological control of diamondback moth in farmer's fields. Progress Report, 1985. 31 - 33.
- Anonymous. (1987).** Biological control of diamondback moth. Progress Report, 1985. 30 - 31.
- Anonymous. (1988).** The apple moth. Zashchita Rastenii Moskva 1: 49.
- Anonymous. (1991).** Demonstration of IPM of diamondback moth on farmers' fields in the lowlands. Progress Report Asian Vegetable Research and Development Center 21 - 25.
- Anonymous. (1991).** Effect of insecticides on the parasitism of diamondback moth by two larval parasite. Progress Report Asian Vegetable Research and Development Center 26 - 29.
- Anonymous. (1991).** Study of parasitism of diamondback moth by larval parasites in various seasons in Taiwan. Progress Report Asian Vegetable Research and Development Center 25 - 26.
- Anonymous. (1994).** Opinion 1772. Metopiini Raffray, 1904 (Insecta: Coleoptera): spelling emended to Metopiasini, and Metopiini Townsend, 1908 (Insecta: Diptera): spelling amended to Metopiaini, so removing the homonymy with Metopiinae Foerster, [1869] (Insecta: Hymenoptera). Bulletin of Zoological Nomenclature 51 (2): 174 - 175.
- Anonymous. (1999).** Beetles in retained strips and stream reserves. Research Note Forestry Tasmania 9: 1 pp.
- Araujo, L.H.A., Braga-Sobrinho, R., Porfirio-de Almeida, R. and De Mesquita, C.K. (1993).** Biological control of the boll weevil. Pesquisa Agropecuaria Brasileira 28 (2): 257 - 261.

- Arretz, P., Lamborot, L., Guerrero, M.A. and Araya, J.E. (1994).** Parasitism of eggs and larva of *Rachiplusia nu* (Guenee) (Lepidoptera: Noctuidae) on cultivated beans in the Metropolitan Region, Chile. *Boletin de Sanidad Vegetal, Plagas* 20 (4): 909 - 917.
- Asaro, C. and Allen, D.C. (1999).** Biology of pine false webworm (Hymenoptera: Pamphiliidae) during an outbreak. *Canadian Entomologist* 131 (6): 729 - 742.
- Asgari, S., Reineke, A., Beck, M. and Schmidt, O. (2002).** Isolation and characterization of a nepriylisin-like protein from *Venturia canescens* virus-like particles. *Insect Molecular Biology* 11 (5): 477 - 485.
- Aslam, M. (1999).** Identification of some parasites of the introduced pine sawfly, *Diprion similis* (Hartig) (Hymenoptera: Diprionidae) by examination of the mature larval remains. *Sarhad Journal of Agriculture* 15 (1): 45 - 46.
- Asquith, A. and Messing, R.H. (1992).** Attraction of Hawaiian ground litter invertebrates to protein hydrolysate bait. *Environmental Entomology* 21 (5): 1022 - 1028.
- Atanasov, A.Z. (1982).** On the volume and nomenclature of some genera of subfamily Anomaloninae (Hymenoptera, Ichneumonidae). (in Russian) *Acta Zoologica Bulgarica* 19: 75-77.
- Atanasov, A.Z. (1988).** Contribution to the fauna of ichneumonids (Hymenoptera: Ichneumonidae) in Bulgaria. *Acta Zoologica Bulgarica* 36: 82 - 84.
- Athanassov, A.Z., Charmillot, P.J., Jeanneret, P. and Renard, D. (1997).** The larval and pupal parasitoids of the codling moth *Cydia pomonella* L. *Revue Suisse de Viticulture, d'Arboriculture et d'Horticulture* 29 (2): 99 - 106.
- Athanassov, A.Z., Jeanneret, P., Charmillot, P.J. and Renard, D. (1998).** Parasitoids of codling moth and other leafrollers (Lepidoptera: Tortricidae) in apple orchards and forests in south-west Switzerland. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 71 (1-2): 153 - 162.
- Athe, T.V. (1990).** The biology of *Diadegma argenteopilosa* Cameron (Hymenoptera: Ichneumonidae) an internal larval parasitoid of *Spodoptera litura* (Fab.). *Entomologist* 109 (1): 2 - 7.
- Aubert, J.F. (1970).** Nouveau supplément aux Ichneumonides non pétiolées avec description d'un genre nouveau. *Bulletin de la Société Entomologique de Mulhouse* Juillet-Aout: 49-56.
- Aubert, J.-F. (1971).** Ichneumonides Agrothereutines d'un genre nouveau et position systématique de colorator Aub. *Bulletin de la Societe Entomologique de Mulhouse* Septembre-Octobre: 73-76.
- Aubert, J.F. (1974).** Recherche des types de Kriechbaumer au Museum de Munich (Ichneum.Cyclopneusticae, Cryptinae et Ophioninae). *Bulletin Mensuel de la Société Linnéenne de Lyon* 43: 262-272.
- Aubert, J.F. (1974).** Ichneumonides pétiolées inédites avec un genre nouveau. . *Bulletin de la Societe Entomologique de Mulhouse* Octobre-Novembre-Décembre: 53-60.

- Aubert, J.F. (1978).** Les Ichneumonides ouest-palearctiques et leurs hotes. 2. Banchinae et Suppl. Aux Pimplinae. Échauffour: É.D.I.F.A.T. - O.P.I.D.A.
- Aubert, J.F. (1979).** Ichneumonides pétiolées inédites avec quatre genres nouveaux. *Bulletin de la Société Entomologique de Mulhouse* Janvier-Mars: 1-8.
- Aubert, J.F. (1980).** Notes sur diverses Ichneumonides mal connues ou inédites. *Bulletin de la Société Entomologique de Mulhouse* Janvier-Mars: 1-6.
- Aubert, J.F. (1984).** Douze ichneumonides non pétiolées mal connues ou inédites. *Bulletin de la Société Entomologique de Mulhouse* Avril-Juin: 17-23.
- Aubert, J.F. (1985).** Ichneumonides Scolobatinae des collections suédoises (suite) et du Musée de Leningrad. *Bulletin de la Société Entomologique de Mulhouse* Octobre-Novembre-Décembre: 49-58.
- Aubert, J.F. (1986).** Quatorze Ichneumonides pétiolées inédites ou synonymes. *Bulletin de la Société Entomologique de Mulhouse* Avril-Juin: 17-25.
- Aubert, J.F. (1989).** Ichneumonides non pétiolées inédites et quatrième suppl. aux Scolobatinae (Ctenopelmatinae): les *Homaspis* Foerst. *Bulletin de la Société Entomologique de Mulhouse* Janvier-Mars: 1-11.
- Aubert, J.-F. (1993).** Nouvelle précisions sur la systématique de quelques Ichneumonides (Hymenoptera, Ichneumonidae). *Nouvelle Revue d'Entomologie* 10: 211-221.
- Aubert, J.-F. (1993).** Ichneumonide Scolobatinae d'un genre nouveau, *Pionpherta* genus novum (Hymenoptera, Ichneumonidae). *Nouvelle Revue d'Entomologie* 10: 358.
- Aubert, J.F. (1996).** Ichneumonides *Netelia* Gray mal étudiées et 6e supplément aux Scolobatinae (Ctenopelmatinae). *Bulletin de la Société entomologique de Mulhouse* Juillet-septembre 1996: 37-42.
- Aubert, J.F. (1998).** Septième supplément aux Ichneumonides Scolobatinae. *Bulletin de la Société entomologique de Mulhouse* Janvier-mars 1998: 1-2.
- Aubert, J.F., Jonaitis, V.P. and Sawoniewicz, J. (1993).** New corrections on the systematics of some ichneumonids (Hymenoptera: Ichneumonidae) (1). *Nouvelle Revue d'Entomologie* 10 (3): 211 - 221.
- Auger, J., Lecomte, C., Paris, J. and Thibout, E. (1989).** Identification of leek-moth and diamondback-moth frass volatiles that stimulate parasitoid, *Diadromus pulchellus*. *Journal of Chemical Ecology* 15 (4): 1391 - 1398.
- Austin, A.D., White, T.C.R., Maelzer, D.A. and Taylor, D.G. (1993).** Biology of *Etiella behrii* Zeller (Lepidoptera: Pyralidae): a pest of seed lucerne in South Australia. *Transactions of the Royal Society of South Australia Incorporated* 117 (1-2): 67 - 76.
- Austin, A.D. and Dangerfield, P.C. (1997).** A new species of *Jarra* Marsh and Austin (Hymenoptera: Braconidae) with comments on other parasitoids associated with the eucalypt longicorn *Phoracantha semipunctata* (F.) (Coleoptera: Cerambycidae). *Australian Journal of Entomology* 36 (4): 327 - 331.
- Austin, A.D. and Dillon, N. (1997).** Extraction and PCR of DNA from parasitoid wasps that have been chemically dried. *Australian Journal of Entomology* 36 (3): 241 - 244.

- Avci, U. and Ozbek, H. (1990).** Lepidopterous cabbage pests and their parasitoids in Erzurum. Proceedings of the Second Turkish National Congress of Biological Control 319 - 330.
- Avila, F.W. and Goff, M.L. (1998).** Arthropod succession patterns onto burnt carrion in two contrasting habitats in the Hawaiian Islands. *Journal of Forensic Sciences* 43 (3): 581 - 586.
- Azerefegne, F. (1999).** The sweet potato butterfly *Acraea acerata* in Ethiopia. Ecology and economic importance. *Acta Universitatis Agriculturae Sueciae Agraria* 195: 62 pp.
- Azhbenov, V.K. (1986).** The grey grain moth. *Zashchita Rastenii Kiev* 7: 26 - 28.
- Azidah, A.A., Fitton, M.G. and Quicke, D.L.J. (2000).** Identification of the *Diadegma* species (Hymenoptera: Ichneumonidae: Campopleginae) attacking the diamondback moth, *Plutella xylostella* (Lepidoptera: Plutellidae). *Bulletin of Entomological Research* 90 (5): 375 - 389.

## B

- Baarlen, P.V., Sunderland, K.D. and Topping, C.J. (1994).** Eggsac parasitism of money spiders (Araneae: Linyphiidae) in cereals, with a simple method for estimating percentage parasitism of *Erigone* spp. eggsacs by Hymenoptera. *Journal of Applied Entomology* 118 (3): 217 - 223.
- Babendreier, D. (2000).** Life history of *Aptesis nigrocincta* (Hymenoptera: Ichneumonidae) a cocoon parasitoid of the apple sawfly, *Hoplocampa testudinea* (Hymenoptera: Tenthredinidae). *Bulletin of Entomological Research* 90 (4): 291 - 297.
- Babendreier, D., Polesny, F., Muller, W. and Olszak, R.W. (1996).** Studies on two ichneumonid parasitoids as potential biological control agents of the European apple sawfly *Hoplocampa testudinea* Klug (Hymenoptera: Tenthredinidae). International conference on integrated fruit production, at Cedzyna, Poland, 28 August - 2 September 1995. *Bulletin OILB SROP* 19 (4): 236 - 240.
- Babendreier, D. and Hoffmeister, T.S. (2002).** Superparasitism in the solitary ectoparasitoid *Aptesis nigrocincta*: the influence of egg load and host encounter rate. *Entomologia Experimentalis et Applicata* 105 (2-3): 63 - 69.
- Babendreier, D. and Polesny, F. (1999).** Observations on the biology and phenology of *Aptesis nigrocincta* (Hym.: Ichneumonidae) parasitizing cocoons of the apple sawfly, *Hoplocampa testudinea* (Hym.: Tenthredinidae). IOBC-WPRS Working Group "Integrated Plant Protection in Orchards". Workshop on arthropod pest problems in pome fruit production at Einsiedeln, Switzerland, November 30 - December 3, 1997. *Bulletin OILB SROP* 22 (7): 57 - 61.
- Badilla, F.F., Solis, A.I. and Alfaro, D. (1991).** Biological control of the sugarcane stem borer *Diatraea* spp. (Lepidoptera: Pyralidae) in Costa Rica. *Manejo Integrado de Plagas* 20 - 21: 39 - 44.



- Baehrecke, E.H., Vinson, S.B. and Williams, H.J. (1990).** Foraging behavior of *Campoletis sonorensis* in response to *Heliothis virescens* and cotton plants. *Entomologia Experimentalis et Applicata* 55 (1): 47 - 57.
- Baehrecke, E.H., Williams, H.J. and Vinson, S.B. (1989).** Electroantennogram responses of *Campoletis sonorensis* (Hymenoptera: Ichneumonidae) to chemicals in cotton (*Gossypium hirsutum* L.). *Journal of Chemical Ecology* 15 (1): 37 - 45.
- Bagheri, M.R., Hatami, B. and Nematollahi, M.R. (2004).** The first record of *Diadegma semiclausum* Hallen, endoparasitoid of larvae of *Plutella xylostella* on wallflower in greenhouse of Isfahan. *Proceedings of 16<sup>th</sup> Iranian Plant Protection Congress*, p. 162.
- Bagnoli, B., Lucchi, A. and Lozzia, C. (2001).** Bionomics of *Cryptoblabes gnidiella* (Milliere) (Pyralidae: Phycitinae) in Tuscan vineyards. *Proceedings of the IOBC-WPRS Working Group "Integrated Control in Viticulture" at Ponte de Lima, Portugal, 2-7 March, 2001*. *Bulletin OILB SROP* 24 (7): 79 - 83.
- Bahana, J.W. (1990).** Bioecological studies on *Dentichasmias busseolae* Heinrich and its potential for biological control of *Chilo partellus* Swinhoe. *Insect Science and its Application* 11 (4-5): 765 - 772.
- Bahena, F., Budia, F., Adan, A., Del Estal, P. and Vinuela, E. (1999).** Scanning electron microscopy of *Hyposoter didymator* (Hymenoptera: Ichneumonidae) in host *Mythimna umbrigerana* (Lepidoptera: Noctuidae) larvae. *Annals of the Entomological Society of America* 92 (1): 144 - 152.
- Bailey, K.E.J. and Alexander, K.N.A. (1993).** Pre-hibernation parasitoid-induced mortality in larvae of *Ladoga camilla* (L.) (Lepidoptera: Nymphalidae). *British Journal of Entomology and Natural History* 6 (2): 36.
- Baker, B.R. (1992).** Emergence and pairing of *Rhyssa persuasoria* L. (Hymenoptera: Ichneumonidae). *British Journal of Entomology and Natural History* 5 (4): 184 - 185.
- Baldacchino, F. and Moleas, T. (1998).** Infestations of *Plutella xylostella* on broccoli in Apulia and the possibility of integrated control. *Informatore Fitopatologico* 48 (4): 71 - 75.
- Baldini, U.A. (2001).** *Tremex fuscicornis*: triple threat. *Chile Forestal* 283: 28 - 30.
- Ballal, C.R., Rao, N.S. and Joshi, S. (2000).** Technique for segregating male and female *Campoletis chlorideae* Uchida at the cocoon stage. *Pest Management in Horticultural Ecosystems* 6 (2): 106 - 109.
- Banchhor, G. (2000).** Seasonal activity of *Helicoverpa armigera* (Hubner) and larval parasitoid, *Campoletis chlorideae* Uchida on chickpea. *Insect Environment* 6 (3): 101 - 102.
- Banita, E., Popov, C., Luca, E., Cojocar, D., Paunescu, G. and Vilau, F. (1992).** Elements of integrated control of wheat stem sawflies (*Cephus pygmaeus* Latr. and *Trachelus tabidus* L.). *Probleme de Protectia Plantelor* 20 (3-4): 169 - 185.
- Barbattini, R. (1986).** *Ostrinia nubilalis* Hb. (Lepidoptera: Pyralidae) on maize in Friuli. III. Study on entomophagous insects. *Frustula Entomologica*, publ. 1989, 9: 63 - 75.

- Barbulescu, A. and Popov, C. (2001).** Development of an integrated control system of pests and diseases of winter wheat and barley crops. *Analele Institutului de Cercetari pentru Cereale si Plante Tehnice*, Fundulea 68: 373 - 384.
- Barlow, N.D., Beggs, J.R. and Moller, H. (1998).** Spread of the wasp parasitoid *Sphecofaga vesparum vesparum* following its release in New Zealand. *New Zealand Journal of Ecology* 22 (2): 205 - 208.
- Barlow, N.D., Moller, H. and Beggs, J.R. (1996).** A model for the effect of *Sphecofaga vesparum vesparum* as a biological control agent of the common wasp in New Zealand. *Journal of Applied Ecology* 33 (1): 31 - 44.
- Barron, J. (1994).** A new species of the *Perilissus anatinus* species group and notes on the relationships of the species groups of *Perilissus* (Hymenoptera: Ichneumonidae: Ctenopelmatinae). *Canadian Entomologist* 126 (2): 219 - 224.
- Barron, J.R. (1989).** Status of the parasite *Agrypon flaveolatum* (Gravenhorst) (Hymenoptera: Ichneumonidae), introduced to control the winter moth in Nova Scotia and British Columbia. *Canadian Entomologist* 121 (1): 11 - 26.
- Barron, J.R. (1990).** The Nearctic species of *Homaspis* (Hymenoptera: Ichneumonidae: Ctenopelmatinae). *Canadian Entomologist* 122 (3-4): 191 - 216.
- Barron, J.R. (1992).** The Nearctic species of *Perilissus* (Hymenoptera: Ichneumonidae: Ctenopelmatinae). *Canadian Entomologist* 124 (2): 211 - 272.
- Barron, J.R. (1995).** The Nearctic species of *Lathrolestes* (Hymenoptera: Ichneumonidae: Ctenopelmatinae). *Contributions of the American Entomological Institute* 28 (3): 135 pp.
- Bartlett, R., Pickering, J., Gauld, I. and Windsor, D. (1999).** Estimating global biodiversity: tropical beetles and wasps send different signals. *Ecological Entomology* 24 (1): 118 - 121.
- Bartninkaite, I. and Jonaitis, V. (1996).** Influence of the physiological state of the Ermine moth *Yponomeuta evonymella* L. to its relations with parasites and microorganisms. *Lietuvos entomologu darbai Lietuvos entomologu draugijos 30 meciui*, 150 - 157.
- Basedow, T. (1999).** Alterations of arthropod fauna in anthropogenic ecosystems: insecticide use, beneficials and developments in agriculture in Panama, Luzon/Philippines and in two regions of Germany. *Pesticidi* 14 (2): 153 - 164.
- Basibuyuk, H.H. and Quicke, D.L.J. (1999).** Gross morphology of multiporous plate sensilla in the Hymenoptera (Insecta). Symposium issue: Phylogeny of the Hymenoptera. *Zoologica Scripta* 28 (1-2): 51 - 67.
- Basset, Y. (1991).** The taxonomic composition of the arthropod fauna associated with an Australian rainforest tree. *Australian Journal of Zoology* 39 (2): 171 - 190.
- Battisti, A. (1994).** Effects of entomopathogenic nematodes on the spruce web-spinning sawfly *Cephalcia arvensis* Panzer and its parasitoids in the field. *Biocontrol Science and Technology* 4 (1): 95 - 102.

- Battisti, A., Bernardi, M. and Ghirardo, C. (2000).** Predation by the hoopoe (*Upupa epops*) on pupae of *Thaumetopoea pityocampa* and the likely influence on other natural enemies. *BioControl* 45 (3): 311 - 323.
- Baur, M.E., Yeargan, K.V. and Cornelius, P.L. (1996).** Allocation of searching time within the soybean canopy by parasitoids attacking the green cloverworm. *Journal of Insect Behavior* 9 (2): 223 - 235.
- Baur, M.E. and Yeargan, K.V. (1994).** Developmental stages and kairomones from the primary parasitoid *Cotesia marginiventris* (Hymenoptera: Braconidae) affect the response of the hyperparasitoid *Mesochorus discitergus* (Hymenoptera: Ichneumonidae) to parasitized caterpillars. *Annals of the Entomological Society of America* 87 (6): 954 - 961.
- Baur, M.E. and Yeargan, K.V. (1994).** Behavioural interactions between the hyperparasitoid *Mesochorus discitergus* (Hymenoptera: Ichneumonidae) and four species of noctuid caterpillars: evasive tactics and capture efficiency. *Journal of Entomological Science* 29 (3): 420 - 427.
- Baur, M.E. and Yeargan, K.V. (1996).** Movement and response to semiochemicals by parasitoids of *Plathypena scabra* (Lepidoptera: Noctuidae). *Journal of the Kansas Entomological Society* 69 (2): 122 - 132.
- Bayram, S. and Ozkan, C. (1998).** Studies on the morphological changes depending on age and host on the internal reproductive system of [parasitoid] *Venturia canescens* Grav. (Hymenoptera: Ichneumonidae). *Turkiye Entomoloji Dergisi* 22 (2): 149 - 159.
- Beck, M., Reineke, A., Lorenz, H., Theopold, U. and Schmidt, O. (2001).** Two distinct reproductive strategies are correlated with an ovarian phenotype in co-existing parthenogenetic strains of a parasitic wasp. *Journal of Insect Physiology* 47 (10): 1189 - 1195.
- Beck, M., Siekmann, G., Theopold, U., Schmidt, O. and Li, D.M. (1999).** A maternal gene mutation correlates with an ovary phenotype in a parthenogenetic wasp population. *Insect Biochemistry and Molecular Biology* 29 (5): 453 - 460.
- Beck, M., Theopold, U. and Schmidt, O. (2000).** Evidence for serine protease inhibitor activity in the ovarian calyx fluid of the endoparasitoid *Venturia canescens*. *Journal of Insect Physiology* 46 (9): 1275 - 1283.
- Beck, M., Theopold, U., Schmidt, O., Austin, A.D. and Downton, M. (2000).** Two genetically distinct *Venturia canescens* strains display different reproductive strategies. Fourth International Hymenoptera Conference, held in Canberra, Australia, in January 1999. *Hymenoptera: evolution, biodiversity and biological control* 38 - 45.
- Beck, N.G. and Cameron, P.J. (1990).** Comparison of lepidopteran pest populations and their parasitoids in three vegetable brassicas. *Proceedings of the Forty Third New Zealand Weed and Pest Control Conference* 21 - 25.
- Beggs, J.R., Austin, A.D. and Downton, M. (2000).** Impact and control of introduced *Vespula* wasps in New Zealand. Fourth International Hymenoptera Conference, held in Canberra,

- Australia, in January 1999. Hymenoptera: evolution, biodiversity and biological control 404 - 409.
- Beggs, J.R., Harris, R.J. and Read, P.E.C. (1996).** Invasion success of the wasp parasitoid *Sphecophaga vesparum vesparum* (Curtis) in New Zealand. *New Zealand Journal of Zoology* 23 (1): 1 - 9.
- Beggs, JR., Harris, R.J. and Read, P.E.C. (1996).** Invasion success of the wasp parasitoid *Sphecophaga vesparum vesparum* (Curtis) in New Zealand. *New Zealand Journal of Zoology* 23 (1): 1 - 9.
- Beggs, J.R., Rees, J.S. and Harris, R.J. (2002).** No evidence for establishment of the wasp parasitoid, *Sphecophaga vesparum burra* (Cresson) (Hymenoptera: Ichneumonidae) at two sites in New Zealand. *New Zealand Journal of Zoology* 29 (3): 205 - 211.
- Beggs, J.R. and Harris, R.J. (2000).** Can the wasp parasitoid *Sphecophaga vesparum* significantly reduce the density of *Vespula* wasps? *New Zealand Journal of Zoology* 27 (1): 73 - 74.
- Begon, M., Sait, S.M., Thompson, D.J., Gange, A.C. and Brown, V.K. (1997).** Two's company, three's a crowd: host-pathogen - parasitoid dynamics. Multitrophic interactions in terrestrial systems: the 36th symposium of the British Ecological Society, Royal Holloway College, University of London, 1995. 307 - 332.
- Bekkaoui, A. and Thibout, E. (1991).** Chemical information in the host-parasitoid couple *Acrolepiopsis assectella* (Lep.) - *Diadromus pulchellus* Hym. *Redia* 74 (3): Appendix, 299 - 300.
- Bekkaoui, A. and Thibout, E. (1992).** Study of cuticular non-volatile compounds of *Acrolepiopsis assectella* (Lepidoptera: Yponomeutoidea) involved in host recognition by the parasitoids *Diadromus pulchellus* and *D. collaris* (Hymenoptera: Ichneumonidae). *Entomophaga* 37 (4): 627 - 639.
- Bekkaoui, A. and Thibout, E. (1993).** Role of the cocoon of *Acrolepiopsis assectella* (Lep.: Hyponomeutoidea) in host recognition by the parasitoid *Diadromus pulchellus* (Hym.: Ichneumonidae). *Entomophaga* 38 (1): 101 - 113.
- Beliveau, C., Laforge, M., Cusson, M. and Bellemare, G. (2000).** Expression of a *Tranosema rostrale* polydnavirus gene in the spruce budworm, *Choristoneura fumiferana*. *Journal of General Virology* 81 (7): 1871 - 1880.
- Benedet, F., Bigot, Y., Renault, S., Pouzat, J. and Thibout, E. (1999).** Identification and perception of stimulus used by the endoparasitoid *Diadromus pulchellus* (Hymenoptera: Ichneumonidae) during host acceptance. *Actes de la IV. Conference Internationale Francophone d'Entomologie, Saint-Malo, France, 5-9 juillet 1998. Annales de la Societe Entomologique de France.* 35: Supp., 364 - 370.
- Benedet, F., Bigot, Y., Renault, S., Pouzat, J. and Thibout, E. (1999).** Polypeptides of *Acrolepiopsis assectella* cocoon (Lepidoptera: Yponomeutoidea): an external host-acceptance kairomone for the parasitoid *Diadromus pulchellus* (Hymenoptera: Ichneumonidae). *Journal of Insect Physiology* 45 (4): 375 - 384.

- Benedet, F., Leroy, T., Gauthier, N., Thibaudeau, C., Thibout, E. and Renault, S. (2002).** Gustatory sensilla sensitive to protein kairomones trigger host acceptance by an endoparasitoid. *Proceedings of the Royal Society of London. Series B, Biological Sciences* 269: 1503, 1879 - 1886.
- Benfatto, D. (1984 - 1985).** Observations on the chemical control of citrus pests: use of pheromone traps to monitor *Archips rosanus* L. and *Prays citri* Mill. *Annali dell'Istituto Sperimentale per l'Agrumicoltura*, publ. 1988, 17 - 18: 213 - 238.
- Benson, J., Pasquale, A., Van Driesche, R. and Elkinton, J. (2003).** Assessment of risk posed by introduced braconid wasps to *Pieris virginiensis*, a native woodland butterfly in New England. *Biological Control* 26 (1): 83 - 93.
- Berberet, R.C. and Bisges, A.D. (1998).** Potential for competition among natural enemies of larvae of *Hypera postica* (Coleoptera: Curculionidae) in the southern plains. *Environmental Entomology* 27 (3): 743 - 751.
- Berg, R., Schuchmann-Feddersen, I. and Schmidt, O. (1988).** Bacterial infection induces a moth (*Ephestia kuhniella*) protein which has antigenic similarity to virus-like particle proteins of a parasitoid wasp (*Venturia canescens*). *Journal of Insect Physiology* 34 (6): 473 - 480.
- Bergmann, D.J. and Oseto, C.Y. (1990).** Life tables of the banded sunflower moth (Lepidoptera: Tortricidae) in the Northern Great Plains. *Environmental Entomology* 19 (5): 1418 - 1421.
- Bernstein, C., Heizmann, A. and Desouhant, E. (2002).** Intraspecific competition between healthy and parasitised hosts in a host-parasitoid system: consequences for life-history traits. *Ecological Entomology* 27 (4): 415 - 423.
- Bernstein, C. and Driessen, G. (1996).** Patch-making and optimal search patterns in the parasitoid *Venturia canescens*. *Journal of Animal Ecology* 65 (2): 211 - 219.
- Berry, J.A. (1990).** Two parasitoid complexes: *Hierodoris atychioides* (Butler) (Lepidoptera: Oecophoridae) and *Icerya purchasi* Maskell (Homoptera: Margarodidae). *New Zealand Entomologist* 13: 60 - 62.
- Berry, J.A. (1991).** Hymenoptera primary types in the New Zealand Arthropod Collection. *New Zealand Journal of Zoology* 18 (3): 323 - 341.
- Berry, J.A., Harris, R.J., Read, P.E.C. and Donovan, B.J. (1997).** Morphological and colour differences between subspecies of *Sphecophaga vesparum* (Hymenoptera: Ichneumonidae). *New Zealand Journal of Zoology* 24 (1): 35 - 46.
- Bertrandi, F. and Schedl, W. (1994).** Biology of the juniper-sawfly, *Monoctenus juniperi* (L.) (Hymenoptera: Diprionidae) and its complex of parasitoids and predators. *Zoologische Jahrbucher, Abteilung fur Systematik, Okologie und Geographie der Tiere* 121 (4): 567 - 592.
- Beukeboom, L.W. and Pijnacker, L.P. (2000).** Automictic parthenogenesis in the parasitoid *Venturia canescens* (Hymenoptera: Ichneumonidae) revisited. *Genome* 43 (6): 939 - 944.

- Beyarslan, A. and Kolarov, J. (1994).** Investigations on Ichneumonidae (Hymenoptera) fauna of Turkey. II. Cryptinae. Turkish Journal of Zoology 18 (4): 227 - 231.
- Bezemer, T.M. and Mills, N.J. (2001).** Host density responses of *Mastrus ridibundus*, a parasitoid of the codling moth, *Cydia pomonella*. Biological Control 22 (2): 169 - 175.
- Bharati, L.R. and Kushwaha, K.S. (1988).** Parasitoids of leaffolder (LF) pupae from Haryana, India. International Rice Research Newsletter 13 (5): 31.
- Bhattacharya, A., Mishra, Y.D., Naqvi, A.H. and Sen, A.K. (1995).** Attraction of some insects associated with lac towards various coloured lights. Journal of Insect Science 8 (2): 205 - 206.
- Bian, X.Y. and Sheng, M.L. (1995).** A new record of Ichneumonidae (Hymenoptera) [*Acaenitus dubitator*] from Xinjiang, China. Entomotaxonomia 17 (4): 276.
- Biddinger, D.J., Felland, C.M. and Hull, L.A. (1994).** Parasitism of tufted apple bud moth (Lepidoptera: Tortricidae) in conventional insecticide and pheromone-treated Pennsylvania apple orchards. Environmental Entomology 23 (6): 1568 - 1579.
- Bidlack, D.S., Grant, J.F. and Pless, C.D. (1991).** Seasonal incidence of larvae of the tobacco budworm and corn earworm, and their parasitoids, on tobacco in eastern Tennessee. Journal of Agricultural Entomology 8 (3): 155 - 162.
- Biever, K.D., Chauvin, R.L., Reed, G.L. and Wilson, R.C. (1992).** Seasonal occurrence and abundance of lepidopterous pests and associated parasitoids on collards in the northwestern United States. Journal of Entomological Science 27 (1): 5 - 18.
- Bigot, Y., Drezen, J.M., Sizaret, P.Y., Rabouille, A., Hamelin, M.H. and Periquet, G. (1995).** The genome segments of DpRV, a commensal reovirus of the wasp *Diadromus pulchellus* (Hymenoptera). Virology New York 210 (1): 109 - 119.
- Bigot, Y., Hamelin, M.H. and Periquet, G. (1991).** Molecular analysis of the genomic organization of the Hymenoptera *Diadromus pulchellus* and *Eupelmus vuilleti*. Journal of Evolutionary Biology 4 (4): 541 - 556.
- Bigot, Y., Rabouille, A., Doury, G., Sizaret, P.Y., Delbost, F., Hamelin, M.H. and Periquet, G. (1997).** Biological and molecular features of the relationships between *Diadromus pulchellus* ascovirus, a parasitoid hymenopteran wasp (*Diadromus pulchellus*) and its lepidopteran host, *Acrolepiopsis assectella*. Journal of General Virology 78 (5): 1149 - 1163.
- Bigot, Y., Rabouille, A., Sizaret, P.Y., Hamelin, M.H. and Periquet, G. (1997).** Particle and genomic characteristics of a new member of the Ascoviridae: *Diadromus pulchellus* ascovirus. Journal of General Virology 78 (5): 1139 - 1147.
- Bigot, Y., Drezen, J.M., Sizaret, P.Y., Rabouille, A., Hamelin, M.H. and Periquet, G. (1995).** The genome segments of DpRV, a commensal reovirus of the wasp *Diadromus pulchellus* (Hymenoptera). Virology New York 210 (1): 109 - 119.
- Bilapate, G.G. (1989).** Investigations on *Heliothis armigera* (Hubner) in Marathwada XXIV. Key mortality factors on cotton and pigeonpea. Journal of Maharashtra Agricultural Universities 14 (3): 258 - 262.

- Bilapate, G.G., Mokate, R.B., Lovekar, R.C. and Bagade, D.N. (1988).** Population ecology of *Heliothis armigera* (Hubner) and its parasites on pulses. Journal of Maharashtra Agricultural Universities 13 (3): 299 - 302.
- Billqvist, A. and Ekblom, B. (2001).** The influence of host plant species on parasitism of pollen beetles (*Meligethes* spp.) by *Phradis morionellu*. Entomologia Experimentalis et Applicata 98 (1): 41 - 47.
- Bin, F., Wackers, F., Romani, R. and Isidoro, N. (1999).** Tyloids in *Pimpla turionellae* (L.) are release structures of male antennal glands involved in courtship behaviour (Hymenoptera: Ichneumonidae). International Journal of Insect Morphology and Embryology 28 (1-2): 61 - 68.
- Biswas, G.C., Kabir, SMH. and Das, G.P. (2001).** Insect pests of sesame, *Sesamum indicum* Linn. in Bangladesh, their succession and natural enemies. Indian Journal of Entomology 63 (2):117 - 124.
- Bland, K.P. and Liston, A.D. (1999).** *Pristiphora borea* (Konow, 1904) (Hymenoptera: Tenthredinidae) reared from larvae on *Betula nana* in Scotland. Entomologist's Gazette 50 (2): 103 - 104.
- Blass, S. and Ruthmann, A. (1989).** Fine structure of the accessory glands of the female genital tract of the ichneumonid *Pimpla turionellae* (Insecta: Hymenoptera). Zoomorphology 108 (6): 367 - 377.
- Blazejewska, A. and Blazejewski, F. (1987).** Reduction of *Meligethes aeneus* F. larvae (Col.: Nitidulidae) by *Isurgus heterocerus* Thoms. (Hymn.: Ichneumonidae) on winter rape and white mustard. Roczniki Nauk Rolniczych, Seria E, Ochrona Roslin. publ. 1988, 17 (2): 169 - 179.
- Blissard, G.W., Theilmann, D.A. and Summers, M.D. (1989).** Segment W of *Campoletis sonorensis* virus: expression, gene products, and organization. Virology New York 69 (1):78 - 89.
- 165. Bloem, K.A. and Duffey, S.S. (1990).** Effect of protein type and quantity on growth and developmental of larval *Heliothis zea* and *Spodoptera exigua* and the endoparasitoid *Hyposoter exiguae*. Entomologia Experimentalis et Applicata 54 (2): 141 - 148.
- Bloem, K.A. and Duffey, S.S. (1990).** Interactive effect of protein and rutin on larval *Heliothis zea* and the endoparasitoid *Hyposoter exiguae*. Entomologia Experimentalis et Applicata 54 (2): 149 - 160.
- Bobadilla, G.D., Vargas, C.H. and Jimenez, R.M. (1993).** Natural enemies of the main Lepidoptera pests of tamarugo *Prosopis tamarugo* Philippi. IDESIA 12: 5 - 11.
- Boeve, J.L. (1996).** Chemoecology of larvae of the European apple sawfly. Ninth International Symposium on Insect - Plant Relationships held on 24-30 June 1995 in Gwatt, Switzerland. Entomologia Experimentalis et Applicata 80 (1): 286 - 288.
- Boeve, J.L., Lengwiler, U., Tollsten, L., Dorn, S. and Turlings, T.C.J. (1996).** Volatiles emitted by apple fruitlets infested by larvae of the European apple sawfly. Phytochemistry 42 (2): 373 - 381.

- Boeve, J.L., Witzgall, P. and El Sayed, A. (1999).** Chemical ecology of the European apple sawfly, *Hoplocampa testudinea*. Proceedings of an IOBC Symposium, Scents in orchards - plant and insect semiochemicals from orchard environments, held in Dachau, Germany, during 21-24 September, 1998. Bulletin OILB SROP 22 (9): 15 - 19.
- Bokor, P. and Cagan, L. (1999).** Occurrence and bionomics of *Eriborus terebrans* (Gravenhorst) (Hymenoptera: Ichneumonidae), a parasitoid of the European corn borer, *Ostrinia nubilalis* Hbn. (Lepidoptera: Pyralidae), in central Europe. Plant Protection Science 35 (1): 17 - 22.
- Bonhof, M.J., Overholt, W.A., Polaszek, A. and Van Huis, A. (1997).** Natural enemies of cereal stemborers in East Africa: a review. Insect Science and its Application 17 (1): 19 - 35.
- Bonsall, M.B. and Hassell, M.P. (1998).** Population dynamics of apparent competition in a host-parasitoid assemblage. Journal of Animal Ecology 67 (6): 918 - 929.
- Borah, B.K. and Dutta, S.K. (2001).** Natural enemies of legume pod borer, *Maruca testulalis* Geyer in Jorhat, Assam. Insect Environment 7 (3): 113 - 114.
- Bordera, S. (1998).** Fauna and distribution of Iberian *Gelina* (Hymenoptera: Ichneumonidae). Fragmenta Entomologica 30 (2): 279 - 293.
- Bordera, S., Selfa, J. and Jimenez, R. (1988).** *Netelia* Gray (Hym.: Ichneumonidae) from the province of Valencia. Boletín de la Asociación Española de Entomología 12: 143 - 151.
- Bordera, S. and Selfa, J. (1992).** Contribution to the knowledge of the *Hemitelina* (Hym.: Ichneumonidae) of the Iberian Peninsula, I: Genus *Aclastus* Foerster, 1868. Boletín de la Asociación Española de Entomología 16: 171 - 176.
- Bordera, S. and Selfa, J. (1992).** Data on the Iberian fauna of the genus *Bathythrix* Foerster 1868 (Hym.: Ichneumonidae). Boletín de la Asociación Española de Entomología 16: 193 - 197.
- Bordera, S. and Selfa, J. (1993).** Contribution to the knowledge of the *Hemitelina* (Hymenoptera: Ichneumonidae) from the Iberian Peninsula, II. Boletín de la Asociación Española de Entomología 17 (2): 255 - 261.
- Bordera, S. and Selfa, J. (1993).** Data on the *Acrolytina* fauna of the Iberian Peninsula (Hymenoptera: Ichneumonida. Nouvelle Revue d'Entomologie 10 (3): 281 - 288.
- Borges, J.D. (1990).** Parasitism of eggs and pupae of *Dirphia araucariae*. Pesquisa Agropecuaria Brasileira 25 (2): 243 - 246.
- Bostanian, N.J., Goulet, H., Masner, L., Avilla, J. and Polesny, F. (2001).** Towards insecticide-free apple orchards in Quebec. Proceedings of the IOBC-WPRS Fifth International Conference on Integrated Fruit Protection, Lleida, Spain, 22-26 October, 2000. Bulletin OILB SROP 24 (5): 389 - 391.
- Boston, M.R. and Nash, R. (1989).** Twenty-one species of Hymenoptera (Ichneumonidae: Ichneumoninae) new to Ireland, with a list of ichneumonine species in the Ulster Museum, Belfast. Irish Naturalists' Journal 23 (4): 134 - 137.



- Bowers, J.P., Richman, D.B. and Ellington, J.J. (1998).** Recovery of *Tetrastichus incertus* (Ratzeburg) from *Hypera postica* (Gyllenhal) in the Mesilla Valley, Dona Ana County, New Mexico. *Southwestern Entomologist* 23 (1): 89 - 90.
- Braga, S.M.P., Dias, M.M. and Pentead-Dias, A.M. (2001).** Bionomic aspects of *Eois tegularia* (Guenee) and *Eois glauculata* (Walker) (Lepidoptera: Geometridae: Larentiinae) and their parasitoids. *Revista Brasileira de Zoologia* 18 (3): 837 - 840.
- Bragg, D.E. (1994).** Notes on parasitoids of *Platyptilia carduidactyla* (Riley) (Lepidoptera: Pterophoridae) in transition zone southeastern Washington. *Pan Pacific Entomologist* 70 (4): 322 - 323.
- Branco, M.C. and Medeiros, M.A. (2001).** Impact of insecticides on Diamondback Moth parasitoids on cabbage in the Federal District of Brazil. *Pesquisa Agropecuaria Brasileira* 36 (1): 7 - 13.
- Brar, K.S. and Shenhmar, M. (1996).** Evaluation of parasitoids for the management of *Chilo infuscatellus* (Snellen) and *Scirpophaga excerptalis* (Fabricius) on sugarcane in Punjab. *Indian Sugar* 46 (2): 121 - 123.
- Braun, D.M., Goyer, R.A. and Lenhard, G.J. (1990).** Biology and mortality agents of the fruit tree leafroller (Lepidoptera: Tortricidae), on baldcypress in Louisiana. *Journal of Entomological Science* 25 (1): 176 - 184.
- Brenner, G.J., Oboyski, P.T. and Banko, P.C. (2002).** Parasitism of *Cydia* spp. (Lepidoptera: Tortricidae) on *Sophora chrysophylla* (Fabaceae) along an elevation gradient of dry subalpine forest on Mauna Kea, Hawaii. *Pan Pacific Entomologist* 78 (2): 101 - 109.
- Bressan-Nascimento, S. and Thibout, E. (2000).** Effects of sulfur-containing precursors of Allium secondary metabolites on behavior and physiology of *Diadromus pulchellus*, a specialist parasitoid of *Acrolepiopsis assectella*. *Journal of Chemical Ecology* 26 (10): 2459 - 2472.
- Breuer, M. and Devkota, B. (1990).** Control of *Thaumetopoea pityocampa* (Den. & Schiff.) by extracts of *Melia azedarach* L. (Meliaceae). *Journal of Applied Entomology* 110 (2): 128 - 135.
- Brewer, M.J., Donahue, J.D. and Morrison, E.J. (1997).** *Bathyplectes parasitoids* (Hymenoptera: Ichneumonidae) of alfalfa weevil, *Hypera postica*, (Coleoptera: Curculionidae) in Wyoming. *Journal of the Kansas Entomological Society* 70 (3): 197 - 202.
- Bribosia, E., Champagne, R. and Bylemans, D. (1999).** The useful organisms in our orchards: the parasitoids of larvae. *Fruiteelt nieuws* 12: 22, 24 - 25.
- Briceno, G.R.A. and Sharkey, M.J. (2000).** *Bassus macadamiae* sp. n. (Hymenoptera: Braconidae: Agathidinae), parasitoid of *Ecdytolopha torticornis* and *E. aurantium* (Lepidoptera: Tortricidae) in Macadamia nut crops in Central and South America. *Journal of Hymenoptera Research* 9 (1): 99 - 103.

- Brock, J.P. and Shaw, M.R. (1997).** *Perithous albicinctus* (Gravenhorst), a large pimpline ichneumon-wasp new to Britain (Hymenoptera: Ichneumonidae). *Entomologist's Gazette* 48 (1): 49 - 50.
- Broglia-Micheletti, S.M.F. and Berti-Filho, E. (2000).** Parasitoids of *Cerconota anonella* (Sepp., 1830) (Lep.: Oecophoridae) in soursop (*Annona muricata* L.). *Scientia Agricola* 57 (3): 565 - 566.
- Brushwein, JR. and Childers, C.C. (1990).** Parasitoids associated with the immature stages of *Selenisa sueroides* (Lepidoptera: Noctuidae). *Florida Entomologist* 73 (2): 337 - 339.
- Buchi, R. (1991).** Monitoring of parasitoids in oilseed rape pests during 1989 in Switzerland. *Bulletin SROP* 14 (6): 54 - 60.
- Buchi, R. (2002).** Mortality of pollen beetle (*Meligethes* spp.) larvae due to predators and parasitoids in rape fields and the effect of conservation strips. *Agriculture, Ecosystems and Environment* 90 (3): 255 - 263.
- Buck, A. (1997).** Comparison of *Bathyplectes curculionis* and *Bathyplectes anurus* as biological control agents for the alfalfa weevil. *Journal of Natural Resources and Life Sciences Education* 26 (1): 75 - 77.
- Bueno, V.H.P. (1988).** Occurrence of nine species of Ichneumonidae (Hymenoptera) in *Brachiaria decumbens* pastures in Tres Coracoes, Minas Gerais. *Anais da Sociedade Entomologica do Brasil*. 17: supl., 175 - 178.
- Bueno, V.H.P. and Fraga, A.I.A. (1988).** Parasitoids associated with *Eueides isabella dianasa* (Cramer, 1782) (Lepidoptera: Heliconiini) in passion fruit and aspects of the biology of *Tetrastichus* sp. (Hymenoptera, Eulophidae). *Turrialba* 38 (2): 83 - 86.
- Bugg, R.L., Ellis, R.T. and Carlson, R.W. (1989).** Ichneumonidae (Hymenoptera) using extrafloral nectar of faba bean (*Vicia faba* L., Fabaceae) in Massachusetts. *Biological Agriculture and Horticulture* 6 (2): 107 - 114.
- Bugg, R.L. and Wilson, L.T. (1989).** *Ammi visnaga* (L.) Lamarck (Apiaceae): associated beneficial insects and implications for biological control, with emphasis on the bell-pepper agroecosystem. *Biological Agriculture and Horticulture* 6 (3): 241 - 268.
- Buleza, V.V. (1990).** Search and host choice by *Exetastes cinctipes* Retz. (Hymenoptera: Ichneumonidae) - parasitoid of the genus *Mamestra* (Lepidoptera: Noctuidae). *Doklady, Biological Sciences* 314 (1-6): 563 - 565.
- Buleza, V.V. (2002).** Influence of former experience on modification of the parasitoid *Exetastes atrator* (Hymenoptera: Ichneumonidae) behavior in search for host. *Zoologicheskii Zhurnal* 81 (3): 329 - 335.
- Buron De, I. and Beckage, N.E. (1992).** Characterization of a polydnavirus (PDV) and virus-like filamentous particle (VLFP) in the braconid wasp *Cotesia congregata* (Hymenoptera: Braconidae). *Journal of Invertebrate Pathology* 59 (3): 315 - 327.
- Butcher, R.D.J., Hubbard, S.F. and Whitfield, W.G.F. (2000).** Microsatellite frequency and size variation in the parthenogenetic parasitic wasp *Venturia canescens* (Gravenhorst) (Hymenoptera: Ichneumonidae). *Insect Molecular Biology* 9 (4): 375 - 384.

- Butcher, R.D.J., Whitfield, W.G.F. and Hubbard, S.F. (2000).** Complementary sex determination in the genus *Diadegma* (Hymenoptera: Ichneumonidae). *Journal of Evolutionary Biology* 13 (4): 593 - 606.
- Butcher, R.D.J., Whitfield, W.G.F. and Hubbard, S.F. (2000).** Single-locus complementary sex determination in *Diadegma chrysostictos* (Gmelin) (Hymenoptera: Ichneumonidae). *Journal of Heredity* 91 (2): 104 - 111.
- Buyukguzel, K. (2001).** Positive effects of some gyrase inhibitors on survival and development of *Pimpla turionellae* (Hymenoptera: Ichneumonidae) larvae reared on an artificial diet. *Journal of Economic Entomology* 94 (1): 21 - 26.
- Buyukguzel, K. and Yazgan, S. (1996).** Effects of some antibiotics on survival and development of the endoparasitoid *Pimpla turionellae* L. (Hymenoptera: Ichneumonidae). *Turkish Journal of Zoology* 20 (1): 1 - 7.

## C

- Caballero, P., Vargas-Osuna, E., Aldebis, H.K. and Santiago-Alvarez, C. (1990).** Parasitoids associated to natural populations of *Spodoptera littoralis* Boisduval and *S. exigua* Hb. (Lepidoptera: Noctuidae). *Boletin de Sanidad Vegetal, Plagas* 16 (1): 91 - 96.
- Caballero, P., Vargas-Osuna, E. and Santiago-Alvarez, C. (1989).** Presence of cutworms on various crops in Andalucia and Extremadura and their associated parasitoids and pathogens. *Boletin de Sanidad Vegetal, Plagas* 15 (1): 3 - 7.
- Caballero, P., Vargas-Osuna, E. and Santiago-Alvarez, C. (1990).** Development of *Apanteles telengai* (Hym.: Braconidae) and *Campoletis annulata* (Hym.: Ichneumonidae) in granulosis virus (GV) infected *Agrotis segetum* (Lep.: Noctuidae) larvae. *Journal of Applied Entomology* 110 (4): 358 - 364.
- Caballero, P., Vargas - Osuna, E. and Santiago - Alvarez, C. (1991).** Parasitization of granulosis-virus infected and noninfected *Agrotis segetum* larvae and the virus transmission by three hymenopteran parasitoids. *Entomologia Experimentalis et Applicata* 58 (1): 55 - 60.
- Cagan, L. and Bokor, P. (1998).** *Sinophorus turionus* Ratz, the parasitoid of the European corn borer, *Ostrinia nubilalis* Hbn. in Slovakia, Czech Republic and Southwestern Poland. *Acta Phytopathologica et Entomologica Hungarica* 33 (3-4): 435 - 445.
- Calvo, R. and Suarez, E. (1991).** Induction of galls in the stem of *Hyptis urticoides* Kunth (Labiatae), by *Glyphipterix* sp. (Lepidoptera: Glyphipterigidae) in Heredia, Costa Rica. *Brenesia* 35: 85 - 95.
- Camillo, E. (1999).** Bionomics of *Trypoxylon (Trypargilum) antropovi* and *Trypoxylon (Trypargilum) aestivale* (Hymenoptera, Sphecidae). *Iheringia, Serie Zoologia* 87: 29 - 36.
- Camillo, E., Garofalo, C.A. and Serrano, J.C. (1994).** Observations on the biology of *Trypoxylon (Trypargilum) rogenhoferi* Kohl (Hymenoptera: Sphecidae). *Anais da Sociedade Entomologica do Brasil* 23 (2): 299 - 310.

- Campadelli, G. (1993).** *Chrysobothris solieri* Lap. Informatore Fitopatologico 43 (12): 25 - 27.
- Campadelli, G. (1996).** Natural enemies of *Zeuzera pyrina* (Lepidoptera Cossidae). Informatore Fitopatologico 46 (9): 13 - 16.
- Campadelli, G. (1998).** Some parasitoids of *Xiphydria longicollis* in the San Vitale pine forest (Ravenna). Informatore Fitopatologico 48 (11): 3 - 6.
- Campadelli, G. and Dindo, M.L. (1994).** *Monochamus galloprovincialis* (Oliv.) (Coleoptera Cerambycidae) in S. Vitale pine forest. Informatore Fitopatologico 44 (1): 31 - 34.
- Campadelli, G. and Scaramozzino, P.L. (1994).** Hymenopterous parasitoids of xylophagous insects in Romagna. Bollettino dell'Istituto di Entomologia 'Guido Grandi' della Universita degli Studi di Bologna 48: 115 - 121.
- Campadelli, G. (1996).** Some parasitoids of *Zeuzera pyrina* L. (Lep.: Cossidae) in Emilia - Romagna. Bollettino dell'Istituto di Entomologia 'Guido Grandi' della Universita degli Studi di Bologna 50: 127 - 131.
- Campos, F., Donskov, N., Arnason, J., Philogene, B.J.R., Atkinson, J., Morand, P. and Werstiuk, N.H. (1990).** Biological effects and toxicokinetics of DIMBOA in *Diadegma terebrans* (Hymenoptera: Ichneumonidae), an endoparasitoid of *Ostrinia nubilalis* (Lepidoptera: Pyralidae). Journal of Economic Entomology 83 (2): 356 - 360.
- Campos De, W.G. and Cure, J.R. (1992).** Parasitism of a natural population of *Myonia piraloides* Walker, 1854 (Lepidoptera: Dioptidae), in *Eucalyptus cloeziana* reforestation. Anais da Sociedade Entomologica do Brasil 21 (1): 241 - 249.
- Canales-Canales, A. and Valdivieso-Jara, L. (1999).** Handbook for biological control in olives. Servicio Nacional de Sanidad Agraria; Jesus Maria; Peru 37 pp.
- Carballo, M., Hernandez, M., Rutilio-Quezada, J. and Quezada, J.R. (1989).** Effect of insecticides and weeds on *Plutella xylostella* (L.) and its parasitoid *Diadegma insulare* (Cress) in cabbage crop. Manejo Integrado de Plagas 11: 1 - 20.
- Carlson, R.W. (1979).** Family Ichneumonidae. In: K.V. Krombein, P.D. Hurd, D.R. Smith, & B.D. Burks (eds.), *Catalog of Hymenoptera in America north of Mexico*. Volume 1. Washington, D.C.: Smithsonian Institution. 1198 pp.
- Carlton, C.E. and Kring, T.J. (1994).** *Melitara prodenialis* Walker on prickly pear in Arkansas. Southwestern Entomologist 19 (1): 23 - 31.
- Carmean, D., Miller, J.C. and Scaccia, B. (1989).** Overwintering of *Phryganidia californica* in the Oregon Cascades and notes on its parasitoids (Lepidoptera: Dioptidae). Pan Pacific Entomologist 65 (1): 74 - 76.
- Carpenter, J.E. (1995).** *Ichneumon promissorius* (Erichson) (Hymenoptera: Ichneumonidae): factors affecting fecundity, oviposition, and longevity. Journal of Entomological Science 30 (2): 279 - 286.
- Carpenter, J.E., Ferkovich, S.M. and Greany, P.D. (2001).** Fecundity and longevity of *Diapetimorpha introita* (Cresson) (Hymenoptera: Ichneumonidae) reared on artificial diets:

- effects of a lipid extract from host pupae and culture media conditioned with an insect cell line. Armyworm Symposium 2000. Florida Entomologist 84 (1): 43 - 49.
- Carpenter, J.E., Pair, S.D. and Fitt, G.P. (1994).** *Ichneumon promissorius* (Hymenoptera: Ichneumonidae): development on North American hosts. Journal of Economic Entomology 87 (4): 929 - 932.
- Carpenter, J.E. and Greany, P.D. (1998).** Comparative development and performance of artificially reared versus host-reared *Diapetimorpha introita* (Cresson) (Hymenoptera: Ichneumonidae) wasps. Biological Control 11 (3): 203 - 208.
- Carver, M., Hart, P.J. and Wellings, P.W. (1993).** Aphids (Hemiptera: Aphididae) and associated biota from The Kingdom of Tonga, with respect to biological control. Pan Pacific Entomologist 69 (3): 250 - 260.
- Casanas-Arango, A., Trujillo, E.E., De Hernandez, A.M. and Taniguchi, G. (1990).** Field biology of *Cyanotricha necyria* Felder (Lep.: Diopitidae), a pest of *Passiflora* spp., in southern Colombia's and Ecuador's Andean region. Journal of Applied Entomology 109 (1): 93 - 97.
- Casiraghi, M., Andrietti, F., Bonasorom, F. and Martinoli, A. (2001).** A note on host detection by *Buathra tarsoleuca* (Schrank) (Hymenoptera: Ichneumonidae), a parasite of *Ammophila sabulosa* (L.) and *Podalonia affinis* (Kirby) (Hymenoptera: Sphecidae). Journal of Insect Behavior 14 (3): 299 - 312.
- Cave, R.D. (1992).** Inventory of parasitic organisms of the striped grass looper, *Mocis latipes* (Lepidoptera: Noctuidae), in Honduras. Florida Entomologist 75 (4): 592 - 598.
- Cha, J.Y., Kim, J.G. and Lee, J.W. (2000).** New records of the genera *Hypsicera* Latreille and *Acerataspis* Uchida (Hymenoptera: Ichneumonidae: Metopiinae) from Korea Insecta Koreana 17 (4): 277 - 285.
- Cha, J.Y., Lee, J.W. and Kim, C.W. (1988).** A systematic study of the Ichneumonidae (Hymenoptera) from Korea X. The tribe Tryphonini (Tryphoninae). Entomological Research Bulletin 14: 45 - 57.
- Cha, J.Y., Lee, J.W. and Kwon, Y.J. (1999).** Zoogeographic analysis on the subfamily Tryphoninae (Hymenoptera: Ichneumonidae). Korean Journal of Applied Entomology 38 (1): 1 - 10.
- Cha, J.Y. and Lee, J.W. (1998).** Taxonomic key of the subfamily Tryphoninae (Hymenoptera: Ichneumonidae), with newly recorded species from Korea. Korean Journal of Entomology 28 (3): 203 - 210.
- Cha, J.Y. and Lee, J.W. (2000).** A taxonomic review of the genus *Colpotrochia* Holmgren (Hymenoptera: Ichneumonidae: Metopiinae) in Korea. Insecta Koreana 17 (3): 221 - 228.
- Chai, X.M., He, Z.H., Jiang, P., Wu, Z.D., Pan, C.R., Hu, R.D. and Ruan, Z.M. (2000).** Studies on natural enemies of *Dendrolimus punctatus* in Zhejiang Province. Journal of Zhejiang Forestry Science and Technology 20 (4): 1 - 56, 61.

- Chakrabarti, S., Ghosh, D. and Debnath, N. (1990).** Hymenopteran parasitoids of aphidophagous dipteran in Western Himalaya, India. *Journal of Aphidology*, publ. 1991, 4 (1-2): 1 - 5.
- 259. Chandler, K.J. and Benson, A.J. (1991).** Evaluation of armyworm infestation in north Queensland sugarcane ratoon crops. *Proceedings of the Australian Society of Sugar Cane Technologists* 13: 79 - 82.
- Chandramohan, N., Babu, T. and Nanjan, K. (1992).** *Acleris extensana* (Meyrick) a new tortricid pest of rose in Nilgiris. *South Indian Horticulture* 40 (5): 304.
- Chandramohan, N. and Chelliah, S. (1990).** Natural enemies of rice yellow stemborer, *Scirpophaga incertulas* (W.) and its relationship with weather elements. *Journal of Biological Control* 4 (2): 89 - 92.
- Chao, H.F. (1994).** On a new species of the genus *Cylloceria* Schi Dte (Hymenoptera: Ichneumonidae: Cylloceriinae). *Wuyi Science Journal* 11: 116 - 119.
- Chao, H.F. and Tang, YQ. (1991).** On a collection of *Enicospilus* and *Dicamptus* taken by Dr. H. Townes from Taiwan, with a description of a new species of *Enicospilus*. *Wuyi Science Journal* 8C: 51 - 54.
- Charles, J.G., Dugdale, J.S. and White, V. (1987).** Preliminary studies of *Eutorna phaulocosma* (Lepidoptera: Oecophoridae) in New Zealand. *New Zealand Journal of Zoology* publ. 1988, 14 (4): 519 - 526.
- Charlet, L.D. (2001).** Biology and seasonal abundance of parasitoids of the banded sunflower moth (Lepidoptera: Tortricidae) in sunflower. *Biological Control* 20 (2): 113 - 121.
- Chauvin, G., E.I. Agoze, M., Hamon, C. and Huignard, J. (1988).** Ultrastructure of the spermatozoa of haploid and diploid males of *Diadromus pulchellus* Wesmael (Hymenoptera: Ichneumonidae). Ultrastructure des spermatozoides des males haploides et diploides de *Diadromus pulchellus* Wesmeal (Hymenoptera: Ichneumonidae). *International Journal of Insect Morphology and Embryology* 17 (4-5): 359 - 366.
- Chen, S.L., Li, Y.G., Luo, P.T. and Ou, Z.S. (1992).** Bionomics of *Amata germana* (Felder) and its control. *Entomological Knowledge* 29 (4): 209 - 211.
- Chen, S.L., Li, Y.G. and Ou, Z.S. (1990).** Bionomics of *Eurema hecabe* L. and its control. *Insect Knowledge* 27 (1): 29 - 34.
- Chen, W.S., Tu, C.C. and Chang, F.I. (1988).** Ecological studies on the new insect pests of asparagus. *Asparagus Research Newsletter* 6 (1): 107 - 111, 24.
- Chen, Y.M., Wang, F.L. and Diangm, Z.K. (1999).** Investigation on parasitic enemies of *Ostrinia furnacalis* larvae. *Plant Protection* 25 (4): 45 - 46.
- Cheng, W.Y., Chen, S.M. and Wang, Z.T. (1999).** Survey of larval and pupal parasitoids of cane borers in spring cane. *Report of the Taiwan Sugar Research Institute* 164: 1 - 14.
- Chiang, F.M. and Sun, C.N. (1991).** Detoxifying enzymes and susceptibility to several insecticides of *Apanteles plutellae* (Hymenoptera: Braconidae) and *Diadegema semiclausum* (Hymenoptera: Ichneumonidae), parasitoids of diamondback moth (Lepidoptera: Plutellidae) larvae. *Environmental Entomology* 20 (6): 1687 - 1690.

- Childs, R., Casagrande, R.A., Tewksbury, L. and Van Driesche, R.G. (1997).** Establishment, distribution, and impact in southern New England of *Lathrolestes nigricollis* (Thompson) (Hymenoptera: Ichneumonidae), an introduced parasitoid of the birch leafminer, *Fenusa pusilla* (Lepeletier) (Hymenoptera: Tenthredinidae). *Canadian Entomologist* 129 (4): 601 - 611.
- Chiu, S.C. (1984).** Keys to the genera of Ichneumonidae (Hymenoptera) of Taiwan. *Taiwan Agricultural Research Institute, Special Publication* 17: 1-58.
- Cho, H.C., Kim, T.H. and Paik, J.C. (1992).** Hymenopteran parasitoids of the onion maggot, *Delia antiqua* (Meigen) (Diptera: Anthomyiidae) in Korea. *Korean Journal of Entomology* 22 (3): 209 - 214.
- Civolani, S. and Pasqualini, E. (1999).** Short-term toxicity of different formulations of insecticides to some groups of predatory arthropods. *Informatore Agrario* 55: 20, 87 - 90.
- Clark, T.L., Foster, J.E., Witkowski, J.F., Siegfried, B.D. and Spencer, T.A. (1997).** Parasitoids recovered from European corn borer, *Ostrinia nubilalis* Hubner, (Lepidoptera: Pyralidae) larvae in Nebraska. *Journal of the Kansas Entomological Society* 70 (4): 365 - 367.
- Coffelt, M.A. and Schultz, P.B. (1993).** Larval parasitism of orangestriped oakworm (Lepidoptera: Saturniidae) in the urban shade tree environment. *Biological Control* 3 (2):127 - 134.
- Coll, M., Gavish, S. and Dori, I. (2000).** Population biology of the potato tuber moth, *Phthorimaea operculella* (Lepidoptera: Gelechiidae), in two potato cropping systems in Israel. *Bulletin of Entomological Research* 90 (4): 309 - 315.
- Colombera, S., Alma, A., Arzone, A. and Lozzia, C. (2001).** Comparison between the parasitoids of *Lobesia botrana* and *Eupoecilia ambiguella* in conventional and integrated vineyards. Proceedings of the IOBC-WPRS Working Group "Integrated Control in Viticulture" at Ponte de Lima, Portugal, 2-7 March, 2001. *Bulletin OILB SROP* 24 (7): 91 - 96.
- Conlong, D.E. and Goebel, R. (2002).** Biological control of *Chilo sacchariphagus* (Lepidoptera: Crambidae) in Mocanbique: the first steps. Proceedings of the 76th Annual Congress of the South African Sugar Technologists' Association, Mount Edgecombe, South Africa, 30 July - 2 August 2002. Proceedings of the Annual Congress South African Sugar Technologists' Association. 76: 310 - 320; published as a CD.
- Conlong, D.E. and Kasl, B. (2001).** Stimulo - deterrent diversion of *Eldana saccharina* (Lep.: Pyralidae) and *Xanthopimpla stemmator* (Hymenoptera: Ichneumonidae) - preliminary results. Proceedings of the 75th Annual Congress of the South African Sugar Technologists' Association, Durban, South Africa, 31 July-3 August 2001. Proceedings of the Annual Congress South African Sugar Technologists' Association 75: 180 - 182.
- Conlong, D.E. and Mugalula, A. (2001).** *Eldana saccharina* (Lep.: Pyralidae) and its parasitoids at Kinyara Sugar Works, Uganda. Proceedings of the 75th Annual Congress of the South African Sugar Technologists' Association, Durban, South Africa, 31 July - 3

- August 2001. Proceedings of the Annual Congress South African Sugar Technologists' Association 75: 183 - 185.
- Constantineanu, M.I. (1969).** Description d'un nouveau genre de la R.S. Roumanie de la sous-famille des Ichneumoninae (Hym., Ichn.) et d'une espèce nouvelle. *Lucrarile Statiunii de Cercetari Marine "Prof. Ioan Borcea" Agigea* 3: 255-258.
- Constantineanu, M.I. (1973).** Un genre et cinq espèces d'ichneumonides nouveaux pour la science. *Revue Roumaine de Biologie* 18: 23-31.
- Constantineanu, M.I. and Pisica, C. (1970).** L'étude de la tribu des Pimplini (Hym. Ichneum.) de la République Socialiste România (Insectes auxiliaires à la sylviculture et à l'agriculture). *Analele Stiintifice ale Universitatii "Al. I. Cuza" din Iasi, Mongraphii* 2: 1-106.
- Constantineanu, M.I. and Pisica, C. (1977).** Hymenoptera. Familia Ichneumonidae. Subfamiliile Ephialtinae, Lycorininae, Xoridinae si Acaenitinae. *Fauna Republicii Socialiste Romania* 9(7): 1-305.
- Constantineanu, R. and Constantineanu, I. (1988).** Parasitoid ichneumonids (Hymenoptera: Ichneumonidae) recovered from the defoliator moths *Lymantria dispar* L. and *Malacosoma neustria* L. in Romania. *Revue Roumaine de Biologie, Biologie Animale* 33 (2): 81 - 85.
- Constantineanu, R. and Constantineanu, I. (1997).** Contributions to the knowledge of ichneumonids (Hymenoptera: Ichneumonidae) from Macin Mountains (II). *Analele Stiintifice ale Universitatii "Al I Cuza" din Iasi. Serie Noua Sectiunea I Biologie Animala* 41- 43: 59 - 67.
- Cook, S.P., Webb, R.E. and White, G.B. (1995).** Estimates of gypsy moth (Lepidoptera: Lymantriidae) mortality at two sample heights following application of gypsy moth nuclear polyhedrosis virus. *Journal of Entomological Science* 30 (4): 507 - 510.
- Coop, L., Knight, A. and Fisher, G. (1989).** Parasitism of orange tortrix on caneberry, *Rubus* spp. in Western Oregon and Washington. *Journal of the Entomological Society of British Columbia* 86: 63 - 65.
- Coop, L.B. and Croft, B.A. (1990).** Diapause and life history attributes of *Phytodietus vulgaris* (Hymenoptera: Ichneumonidae), a parasitoid of *Argyrotaenia citrana* (Lepidoptera: Tortricidae). *Annals of the Entomological Society of America* 83 (6): 1148 - 1151.
- Copley, K.J. and Grant, J.F. (1998).** Statewide distribution of parasitoids of the alfalfa weevil in Tennessee. *Journal of Agricultural Entomology* 15 (1):43 - 51.
- Cordero, J. and Cave, R.D. (1992).** Natural enemies of *Plutella xylostella* (Lep.: Plutellidae) on crucifers in Honduras. *Entomophaga* 37 (3): 397 - 407.
- Cordero, R.J. and Cave, R.D. (1990).** Parasitism of *Plutella xylostella* L. (Lepidoptera: Plutellidae) by *Diadegma insulare* (Cresson) (Hymenoptera: Ichneumonidae) in cabbages *Brassica oleracea* var. capitata in Honduras. *Manejo Integrado de Plagas* 16: 19 - 22.



- Corley, J.C. (1999).** Host density signal in relation to aggregation in the parasitoid *Venturia canescens*. Netherlands Journal of Zoology 49 (2): 97 - 109.
- Cortez, M.H. and Trujillo, A.J. (1994).** Incidence of the fall armyworm and its natural enemies in three maize agrosystems. Turrialba 44 (1): 1 - 9.
- Cortez-Madrigal, H. (1998).** Impact of weeds on the biocontrol of *Spodoptera frugiperda* (Lep.: Noctuidae) in corn. Revista Chapingo. Serie Ingenieria Agropecuaria 1 (1): 19 - 23.
- Coruh, S., Ozbek, H. and Kolarov, J. (2002).** New and rare taxa of Ichneumonidae (Hymenoptera) from Turkey. Journal of the Entomological Research Society 4 (1):1 - 4.
- Cossentine, J.E. and Jensen, L.B. (1995).** *Orthosia hibisci* Guenee (Lepidoptera: Noctuidae): indigenous parasitoids and the impact of *Earinus limitarus* (Say) (Hymenoptera: Ichneumonidae) on its host feeding activity. Canadian Entomologist 127 (4): 473 - 477.
- Costa, E.C. and Link, D. (1989).** Behaviour of predators and parasitoids on soyabean crop in relation to insecticide application. Revista do Centro de Ciencias Rurais 19 (4): 317 - 323.
- Covassi, M. and Francardi, V. (1994).** On the presence of *Argyresthia illuminatella* Zeller in fir woods of central and south Italy with notes on its biology (Lepidoptera: Argyresthiidae). Redia 77 (1): 123 - 131.
- Cowgill, S.E. (1995).** Influence of the chickpea cropping system on *Helicoverpa armigera* (Lep.: Noctuidae) populations and their rate of parasitism by *Campoletis chlorideae* (Hym.: Ichneumonidae). Entomophaga 40 (3-4): 307 - 315.
- Craig, T.P., Itami, J.K. and Price, P.W. (1990).** The window of vulnerability of a shoot-galling sawfly to attack by a parasitoid. Ecology 71 (4): 1471 - 1482.
- Cronin, J.T. (1989).** Inverse density-dependent parasitism of the bagworm, *Thyridopteryx ephemeraeformis* (Lepidoptera: Psychidae). Environmental Entomology 18 (3): 403 - 407.
- Cronin, J.T. and Gill, D.E. (1989).** The influence of host distribution, sex, and size on the level of parasitism by *Itopectis conquisitor* (Hymenoptera: Ichneumonidae). Ecological Entomology. 14 (2): 163 - 173.
- Crosby, D.F. (1994).** Further parasite associations for some Australian butterflies (Lepidoptera). Australian Entomologist 21 (3): 81 - 88.
- Cross, J., Jay, C., Avilla, J. and Polesny, F. (2001).** Exploiting the parasitoids *Lathrolestes ensator* and *Platygaster demades* for control of apple sawfly and apple leaf midge in IPM in apple orchards. Proceedings of the IOBC-WPRS Fifth International Conference on Integrated Fruit Protection, Lleida, Spain, 22-26 October, 2000. Bulletin OILB SROP 24 (5): 161 - 165.
- Cruz, I., Figueiredo, M.L.C., Goncalves, E.P., Lima, D.A.N. and Diniz, E.E. (1997).** Effect of age of larvae of *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae) on the performance of the parasitoid *Campoletis flavicincta* (Ashmead) (Hymenoptera: Ichneumonidae) and foliar consumption by parasitized and unparasitized larvae. Anais da Sociedade Entomologica do Brasil 26 (2): 229 - 234.

- Cruz, I., Figueiredo, M.L.C., Valicente, F.H. and Oliveira, A.C. (1997).** Application rate trials with a nuclear polyhedrosis virus to control *Spodoptera frugiperda* (Smith) on maize. *Anais da Sociedade Entomologica do Brasil* 26 (1): 145 - 152.
- Cruz, I., Lima, D.A.N., Figueiredo, M.L.C. and Valicente, F.H. (1995).** Aspects of the biology of the parasitoid *Campoletis flavicincta* (Ashmead) with larvae of *Spodoptera frugiperda* (Smith). *Anais da Sociedade Entomologica do Brasil* 24 (2): 201 - 208.
- Csoka, G., Lesko, K. and Ambrus, A. (1989).** Biology and damage of *Dendrolimus pini* L. (Lepidoptera: Lasiocampidae) in Hungary. *Novenyvedelem* 25: (2): 61 - 65.
- Cui, J.J. and Xia, J.Y. (1998).** Effects of early seasonal strain of Bt transgenic cotton on population dynamics of main pests and their natural enemies. *Acta Gossypii Sinica* 10 (5): 255 - 262.
- Cusson, M., Barron, J.R., Goulet, H., Regniere, J. and Doucet, D. (1998).** Biology and status of *Tranosema rostrale* rostrale (Hymenoptera: Ichneumonidae), a parasitoid of the eastern spruce budworm (Lepidoptera: Tortricidae). *Annals of the Entomological Society of America* 91 (1): 87 - 93.
- Cusson, M., Laforge, M., Miller, D., Cloutier, C. and Stoltz, D. (2000).** Functional significance of parasitism - induced suppression of juvenile hormone esterase activity in developmentally delayed *Choristoneura fumiferana* larvae. *General and Comparative Endocrinology* 117 (3): 343 - 354.
- Cusson, M., Laforge, M., Regniere, J., Beliveau, C., Trudel, D., Thireau, J.C., Bellemare, G., Keirstead, N. and Stolz, D. (2002).** Multiparasitism of *Choristoneura fumiferana* by the ichneumonid *Tranosema rostrale* and the tachinid *Actia interrupta*: occurrence in the field and outcome of competition under laboratory conditions. *Entomologia Experimentalis et Applicata* 102 (2): 125 - 133.
- Cusson, M., Lucarotti, C., Stoltz, D., Krell, P. and Doucet, D. (1998).** A polydnavirus from the spruce budworm parasitoid, *Tranosema rostrale* (Ichneumonidae). *Journal of Invertebrate Pathology* 72 (1): 50 - 56.

## D

- Dai, X.F. (1989).** Observations on *Campoletis chlorideae* (Hym.: Ichneumonidae), an important native parasitoid of cotton bollworm [? *Helicoverpa armigera*]. *Chinese Journal of Biological Control* 5 (4): 183.
- Dai, X.F. (1990).** Biology of *Campoletis chlorideae* (Hym.: Ichneumonidae) and its control of cotton bollworm in the field. *Chinese Journal of Biological Control* 6 (4): 153 - 156.
- Dapoto, G. and Giganti, H. (1994).** Bioecology of *Nematus desantisi* (Hymenoptera: Tenthredinidae: Nematinae) in Rio Negro y Neuquen, Argentina. *Bosque* 15 (1): 27 - 32.
- Darvish Mojeini, T. and Sedivy, J. 2001.** New report of parasitoid ichneumonid wasps of cotton bollworm, *Helicoverpa armigera* (Hub.) (Lep.: Noctuidae) in Iran. *Journal of Entomological Society of Iran*, 21(1): 107-108.

- Dasch, C.E. (1974).** Neotropic Mesochorinae (Hymenoptera: Ichneumonidae). *Memoirs of the American Entomological Institute* 22: 1-509.
- Dasch, C.E. (1979).** Ichneumon-flies of America north of Mexico: 8. Subfamily Cremastinae. *Memoirs of the American Entomological Institute* 29: 1-702.
- Dasch, C.E. (1984).** Ichneumon-flies of America north of Mexico: 9. Subfamilies Theriinae and Anomaloninae. *Memoirs of the American Entomological Institute* 36: 1-610.
- Dasch, C.E. (1992).** The ichneumon-flies of America north of Mexico: Part 12. Subfamilies Microleptinae, Helictinae, Cylloceriinae and Oxytorinae (Hymenoptera: Ichneumonidae). *Memoirs of the American Entomological Institute* 52: 1-470.
- Davatchi, A. and Shojai, M. (1969).** Les hymenopteres entomophages de l'Iran (etudes faunistiques). Univ. Tehran, Faculty agric., 107: 1-88.
- Davies, D.H. and Vinson, S.B. (1988).** Interference with function of plasmatocytes of *Heliothis virescens* in vivo by calyx fluid of the parasitoid *Campoletis sonorensis*. *Cell and Tissue Research* 251 (2): 467 - 475.
- Davies, T.H. (1990).** A possible new record of a parasite attacking the bagworm moth, *Liothula omnivorus* (Fereday). *Weta* 13 (1): 12.
- Davis, A.J. (1987).** Effects of experiences and kairomones on searching behaviour of *Diadegma eucerophaga* (Ichneumonidae) for *Plutella xylostella* (Plutellidae) on Brassica plants. *Mededelingen van de Faculteit Landbouwwetenschappen, Rijksuniversiteit Gent* 52 (2a): 403 - 411.
- Dawah, H.A. and Rothfritz, H. (1996).** Generic-level identification of final instar larvae of Eurytomidae and their parasitoids associated with grasses (Poaceae) in N.W. Europe (Hymenoptera: Braconidae, Eulophidae, Eupelmidae, Eurytomidae, Ichneumonidae, Pteromalidae). *Journal of Natural History* 30 (10): 1517 - 1526.
- Day, W.H. (2002).** Biology, host preferences, and abundance of *Mesochorus curvulus* (Hymenoptera: Ichneumonidae), a hyperparasite of *Peristenus* spp. (Hymenoptera: Braconidae) parasitizing plant bugs (Miridae: Hemiptera) in alfalfa-grass forage crops. *Annals of the Entomological Society of America* 95 (2): 218 - 222.
- Dayakar, S. and Ray, S.N. (1999).** Natural parasitization of *Helicoverpa armigera* (Hub.) in pigeonpea ecosystems at Pantnagar. *Insect Environment* 4 (4):136.
- Dbar, R.S. (1984).** Review of the Palearctic species of the genus *Cymodusa* Holmgren (Hymenoptera, Ichneumonidae). (in Russian) *Entomologicheskoye Obozreniye* 63: 802-812.
- Dbar, R.S. (1984).** A new genus and new species of ichneumonids of the tribe Limneriini (Hymenoptera, Ichneumonidae) from the USSR and the Mongolian People's Republic. (in Russian) *Nasekomye. Mongol.* 9: 438-445.
- Dbar, R.S. (1985).** Revision of the Palearctic species of the genus *Cymodusa* Holmgren (Hymenoptera, Ichneumonidae). II. (in Russian) *Entomologicheskoye Obozreniye* 64: 588-600.

- Dbar, R.S. and Saparmamedova, N.K. (1988).** A new species of ichneumonid of the genus *Eriborus* (Hymenoptera: Ichneumonidae) - a parasitoid of the Turkmenian lymantriid *Portesia cargalica* Moore in the central Kopetdag. *Izvestiya Akademii Nauk Turkmenskoi SSR. Seriya Biologicheskikh Nauk* 6: 62 - 64.
- DeGooyer, T.A., Pedigo, L.P. and Giles, K.L. (1995).** Population dynamics of the alfalfa weevil (Coleoptera: Curculionidae) in central and southern Iowa. *Journal of the Kansas Entomological Society* 68 (3): 268 - 278.
- Deka, N.K., Prasad, D. and Chand, P. (1987).** Role of *Campoletis chlorideae* Uchida in *Heliothis* management as affected by sowing dates of chickpea. *Research and Development Reporter* 4 (2): 165 - 172.
- Delrio, G. (1975).** Revision des espèces ouest-palearctiques du genre *Netelia* Gray (Hym. Ichneumonidae). *Studi Sassaesi Sez. III - Annali della Facolta di Agraria dell'Universita di Sassari* 23: 1-126.
- Delvare, G. and Kirk, A. (1999).** Potential of parasitic hymenoptera for the biocontrol of *Plutella xylostella* (L.) (Lep.: Yponomeutidae): a review from past experience and recent exploration. *Proceedings of the Fifth International Conference on Pests in Agriculture, Part 3, Montpellier, France, 7-9 December, 1999.* 655 - 663.
- Deng, L. and Webb, B.A. (1999).** Cloning and expression of a gene encoding a *Campoletis sonorensis* polydnavirus structural protein. *Archives of Insect Biochemistry and Physiology* 40 (1): 30 - 40.
- Dennill, G.B. and Pretorius, W.L. (1995).** The status of diamondback moth, *Plutella xylostella* (Linnaeus) (Lepidoptera: Plutellidae), and its parasitoids on cabbages in South Africa. *African Entomology* 3 (1): 65 - 71.
- Desai, J.C. and Patel, J.R. (2001).** Hyperparasites of syrphids in Middle Gujarat. *Indian Journal of Entomology* 63 (2):195 - 197.
- Devi, N. and Raj, D. (1995).** Biology and parasitization of diamondback moth, *Plutella xylostella* L. infesting cauliflower in mid hill region of Himachal Pradesh (India). *Journal of Entomological Research* 19 (1): 83 - 86.
- Devi, N.S., Singh, O.H., Devjani, P. and Singh, T.K. (2002).** Natural enemies of *Helicoverpa armigera* Hubner on chickpea. *Annals of Plant Protection Sciences* 10 (2): 179 - 183.
- Dhembare, A.J. (1999).** Preference of host *Helicoverpa armigera* by larval parasite *Campoletis chlorideae* Uchida. *Journal of Experimental Zoology, India* 2 (1): 27 - 28.
- Diaconu, A., Pisica, C., Andriescu, I. and Lozan, A. (2000).** The complex of parasitoids of the feeding larvae of *Cydia pomonella* L. (Lep.: Tortricidae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 73 (1-2): 13 - 22.
- Diaconu, A. and Lozan, A. (2000).** Ectoparasitoid braconids (Hym.: Braconidae) of certain species of leafrollers (Lep.: Tortricidae) of fruit trees. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 73 (1-2): 5 - 11.

- Dias, M.M. (1991).** Immature stages of *Procitheronia principalis* (Walker, 1855), compared with those of some species of the genus *Citheronia* Hubner, (1819) (Lepidoptera: Saturniidae). *Revista Brasileira de Entomologia* 35 (2): 269 - 282.
- Diaz, B.F.A. (1993).** Updated list of the ichneumonids (Hymenoptera: Parasitica) of Venezuela. *Manejo Integrado de Plagas* 29: 40 - 42.
- Diaz, F.A. (2000).** The Venezuelan species of *Pimpla* (Hymenoptera: Ichneumonidae). *Journal of Hymenoptera Research* 9 (2): 246 - 253.
- Dib-Hajj, S.D., Webb, B.A. and Summers, M.D. (1993).** Structure and evolutionary implications of a 'cysteine-rich' *Campoletis sonorensis* polydnavirus gene family. *Proceedings of the National Academy of Sciences of the United States of America* 90 (8): 3765 - 3769.
- Didonet, J., Panato, A.P., Dos Santos, G.R. and Peluzio, J.M. (1999).** Insect fauna associated to acerola plant (*Malpighia* spp.) in Gurupi, Tocantins State. *Revista Brasileira de Fruticultura* 21 (2): 212 - 215.
- Digweed, S.C. (1998).** Mortality of birch leafmining sawflies (Hymenoptera: Tenthredinidae): impacts of natural enemies on introduced pests. *Environmental Entomology* 27 (6): 1357 - 1367.
- Dijkerman, H.J. (1988).** Notes on the parasitisation behaviour and larval development of *Trieceles tricarinatus* and *Triclistus yponomeutae* (Hymenoptera: Ichneumonidae), endoparasitoids of the genus *Yponomeuta* (Lepidoptera: Yponomeutidae). *Proceedings of the Koninklijke Nederlands Akademie van Wetenschappen, Series C, Biological and Medical Sciences* 91 (1): 19 - 30.
- Dijkerman, H.J. (1990).** Suitability of eight *Yponomeuta*-species as hosts of *Diadegma armillata*. *Entomologia Experimentalis et Applicata* 54 (2): 173 - 180.
- Dijkerman, H.J. (1991).** Acceptance and suitability of *Yponomeuta*-species as hosts for *Trieceles tricarinatus* and *Triclistus yponomeutae*. *Proceedings of the Koninklijke Nederlands Akademie van Wetenschappen. Series C, Biological and Medical Sciences* 94 (4): 445 - 453.
- Dijkerman, H.J. and Koenders, J.T.H. (1988).** Competition between *Trieceles tricarinatus* and *Triclistus yponomeutae* in multiparasitized hosts. *Entomologia Experimentalis et Applicata* 47 (3): 289 - 295.
- Diller, E. (1970).** Eine neue Gattung der Unterfamilie Diplazontinae (Hym., Ichneumonidae). *Nachrichtenblatt der Bayerischen Entomologen* 19: 8-10.
- Diller, E. (1981).** Bemerkungen zur Systematik der Phaeogenini mit einem vorläufigen Katalog der Gattungen (Hymenoptera, Ichneumonidae). *Entomofauna* 2: 93-111.
- Diller, E. (1982).** Diplazontinae der australischen Region (Hymenoptera, Ichneumonidae, Diplazontinae). *Entomofauna* 3: 287-322.
- Diller, E. (1984).** Studie über Diplazontinae aus Sudamerika (Hymenoptera, Ichneumonidae, Diplazontinae). *Entomofauna* 5: 101-123.

- Diller, E. (1985).** Eine neue Gattung und Art der unterfamilie Ichneumoninae (Hymenoptera, Ichneumonidae). *Entomofauna* 6: 1-7.
- Diller, E. (1994).** Beschreibungen einer neuen Subtribus, neuer Gattungen und Arten der Tribus Phaeogenini (Insecta: Hymenoptera: Ichneumonidae: Phaeogenini). *Annalen des Naturhistorischen Museums in Wien* 96B: 125-136.
- Diller, E. & K. Schönitzer. (1995).** Eine neue Gattung und Art der Phaeogenini aus der neotropischen Region (Insecta: Hymenoptera, Ichneumonidae, Ichneumoninae). *Entomofauna* 16: 453-464.
- Dillon, N., Austin, A.D. and Bartowsky, E. (1996).** Comparison of preservation techniques for DNA extraction from hymenopterous insects. *Insect Molecular Biology* 5 (1): 21 - 24.
- Dimova, M. (1987).** The parasitic entomofauna of *Anarsia lineatella* Zell. (Gelechiidae) and *Grapholitha molesta* Busck. (Lepidoptera: Tortricidae) and its importance in reducing the density of these pests. *Pochvoznanie, Agrokhimiya i Rastitelna Zashchita* 22 (6): 87 - 92.
- Dix, M.E., Hodges, L., Brandle, J.R., Wright, R.J. and Harrell, M.O. (1997).** Effects of shelterbelts on the aerial distribution of insect pests in muskmelon. *Journal of Sustainable Agriculture* 9 (2-3): 5 - 24.
- Doak, P. (2000).** The effects of plant dispersion and prey density on parasitism rates in a naturally patchy habitat. *Oecologia* 122 (4): 556 - 567.
- Doganlar, M. (1987).** Investigations on leafrollers and other Lepidoptera feeding in a similar manner on apple and pear trees, together with their parasitoids, in Erzurum and its neighbourhood. *Doga, Turk Tarim ve Ormancilik Dergisi* 11 (1): 86 - 93.
- Dolgin, M.M. (1990).** Biology of *Pissodes validirostris* in the Komi ASSR. *Lesnoe Khozyaistvo* 8: 59 - 60.
- Dominguez-Jimenez, I., Llanderal-Cazares, C. and Nieto-Hernandez, R. (2000).** *Pristomerus spinator* fabricius (Hymenoptera: Ichneumonidae), a parasite of the potato tuber moth. *Agrociencia* 34 (5): 611 - 617.
- Donovan, B.J. (1989).** Potential enemies of the introduced wasp parasitoid *Sphecofaga vesparum* (Hymenoptera: Ichneumonidae) in New Zealand. *New Zealand Journal of Zoology* 16 (3): 365 - 367.
- Donovan, B.J. (1992).** Problems caused by immigrant German and common wasps in New Zealand, and attempts at biological control. *Bee World* 73 (3): 131 - 148.
- Donovan, B.J. (1996).** Progress with biological control of wasps. *New Zealand Beekeeper* 3 (4): 5, 14 - 15; 12 - 13.
- Donovan, B.J. (2000).** Correction of errors in two recent wasp-related papers from Landcare Research Ltd. *New Zealand Journal of Zoology* 27 (1): 71 - 73.
- Donovan, B.J., Marris, J.W.M., Schroeder, N.C. and Read, P.E.C. (1990).** Inoculation of wasp nests with the parasitoid *Sphecofaga vesparum*. *Proceedings of the Forty Third New Zealand Weed and Pest Control Conference* 195 - 196.
- Donovan, B.J., Moller, H., Plunkett, G.M., Read, P.E.C. and Tilley, J.A.V. (1989).** Release and recovery of the introduced wasp parasitoid *Sphecofaga vesparum vesparum*

- (Curis) (Hymenoptera: Ichneumonidae) in New Zealand. *New Zealand Journal of Zoology* 16 (3): 355 - 364.
- Donovan, B.J., Moller, H. and Fordham, R.A. (1991).** Setting priorities for wasp research. *New Zealand Journal of Zoology* 18 (4): 455 - 458.
- Doucet, D. and Cusson, M. (1996).** Alteration of developmental rate and growth of *Choristoneura fumiferana* parasitized by *Tranosema rostrale*: a role of the calyx fluid. *Entomologia Experimentalis et Applicata* 81 (1): 21 - 30.
- Doucet, D. and Cusson, M. (1996).** Role of calyx fluid in alterations of immunity in *Choristoneura fumiferana* larvae parasitized by *Tranosema rostrale*. *Comparative Biochemistry and Physiology. A, Physiology* 114 (4): 311 - 317.
- Douglas, S.D. and Stockey, R.A. (1996).** Insect fossils in middle Eocene deposits from British Columbia and Washington State: faunal diversity and geological range extensions. *Canadian Journal of Zoology* 74 (6): 1140 - 1157.
- Dover, B.A., Davies, D.H. and Vinson, S.B. (1988).** Dose-dependent influence of *Campoletis sonorensis* polydnavirus on the development and ecdysteroid titers of last-instar *Heliothis virescens* larvae. *Archives of Insect Biochemistry and Physiology* 8 (2): 113 - 126.
- Dover, B.A., Strand, M.R., Davies, D.H. and Vinson, S.B. (1989).** Ultrastructure of host tissues exposed to the calyx fluid of the parasitoid, *Campoletis sonorensis* (Cameron) (Hymenoptera: Ichneumonidae). *International Journal of Insect Morphology and Embryology* 18 (1): 47 - 57.
- Dover, B.A., Tanaka, T. and Vinson, S.B. (1995).** Stadium-specific degeneration of host prothoracic glands by *Campoletis sonorensis* calyx fluid and its association with host ecdysteroid titers. *Journal of Insect Physiology* 41 (11): 947 - 955.
- Dover, B.A. and Vinson, S.B. (1990).** Effect of host ligation and starvation on the development and emergence of the parasitoid *Campoletis sonorensis*. *Entomologia Experimentalis et Applicata* 57 (3): 209 - 213.
- Dover, B.A. and Vinson, S.B. (1990).** Stage-specific effects of *Campoletis sonorensis* parasitism on *Heliothis virescens* development and prothoracic glands. *Physiological Entomology* 15 (4): 405 - 414.
- Dowton, M. and Austin, A.D. (1994).** Molecular phylogeny of the insect order Hymenoptera: aprocritan relationships. *Proceedings of the National Academy of Sciences of the United States of America* 91: 9911 - 9915.
- Driessen, G., Bernstein, C., Kacelnik, A. and Van Alphen, J.J.M. (1995).** A count-down mechanism for host search in the parasitoid *Venturia canescens*. *Journal of Animal Ecology* 64 (1): 117 - 125.
- Driessen, G. and Bernstein, C. (1999).** Patch departure mechanisms and optimal host exploitation in an insect parasitoid. *Journal of Animal Ecology* 68 (3): 445 - 459.

- Duale, A.H. and Nwanze, K.F. (1999).** Incidence and distribution in sorghum of the spotted stem borer *Chilo partellus* and associated natural enemies in farmers' fields in Andhra Pradesh and Maharashtra states. *International Journal of Pest Management* 45 (1): 3 - 7.
- Dubey, O.P., Odak, S.C. and Gargav, V.P. (1993).** Population dynamics of gram pod borer. *JNKVV Research Journal publ.* 1995, 27 (1): 59 - 63.
- Durdvey, S.K. (1987).** Contribution to the study of natural enemies of the codling moth *Laspeyresia pomonella* L. (Lepidoptera: Tortricidae) in the Prikopetdag zone. *Izvestiya Akademii Nauk Turkmenskoi SSR. Seriya Biologicheskikh Nauk* 2: 22 - 26.
- Dyer, L.E. and Landis, D.A. (1996).** Effects of habitats, temperature, and sugar availability on longevity of *Eriborus terebrans* (Hymenoptera: Ichneumonidae). *Environmental Entomology* 25 (5): 1192 - 1201.
- Dyer, L.E. and Landis, D.A. (1997).** Influence of noncrop habitats on the distribution of *Eriborus terebrans* (Hymenoptera: Ichneumonidae) in cornfields. *Environmental Entomology* 26 (4): 924 - 932.

## E

- Eberhard, W.G. (2000).** The natural history and behavior of *Hymenoepimecis argyraphaga* (Hymenoptera: Ichneumonidae) a parasitoid of *Plesiometa argyra* (Araneae: Tetragnathidae). *Journal of Hymenoptera Research* 9 (2): 220 - 240.
- Eberhard, W.G. (2001).** Under the influence: webs and building behavior of *Plesiometa argyra* (Araneae: Tetragnathidae) when parasitized by *Hymenoepimecis argyraphaga* (Hymenoptera: Ichneumonidae). *Journal of Arachnology* 29 (3): 354 - 366.
- Eggleton, P. (1991).** Patterns in male mating strategies of the Rhyssini: a holophyletic group of parasitoid wasps (Hymenoptera: Ichneumonidae). *Animal Behavior* 41 (5): 829 - 837.
- Eghtedar, E. (1996).** Biology of *Lobesia botrana* in Fars province. *Applied Entomology and Phytopathology* 63 (1-2): 5 - 6.
- Eichhorn, O. (1996).** Experimental studies upon the parasitoid complex of the gypsy moth (*Lymantria dispar* L.) (Lep.: Lymantriidae) in lower host populations in eastern Austria. *Journal of Applied Entomology* 120 (4): 205 - 212.
- Eichhorn, O. and Bogenschutz, H. (2000).** Studies on the web-spinning spruce sawflies of the genus *Cephalcia* spp. 1 Panz. (Hym.: Pamphiliidae). IV. Population dynamics of *Cephalcia abietis* L. in Upper Swabia. *Journal of Applied Entomology* 124 (3-4): 121 - 139.
- Eisenbach, J. (1996).** Three-trophic-level interactions in cattail hybrid zones. *Oecologia* 105 (2): 258 - 265.
- El Agoze, M., Poirie, M. and Periquet, G. (1995).** Precedence of the first male sperm in successive matings in the Hymenoptera *Diadromus pulchellus*. *Entomologia Experimentalis et Applicata* 75 (3): 251 - 255.
- El Agoze, M. and Periquet, G. (1993).** Viability of diploid males in the parasitic wasp, *Diadromus pulchellus* (Hym.: Ichneumonidae). *Entomophaga* 38 (2): 199 - 206.



- Elgueta, M., Rojas, F., Martin-Piera, F., Morrone, J.J. and Melic, A. (2000).** Chilean Hymenoptera. Hacia un Proyecto CYTED para el Inventario y Estimacion de la Diversidad Entomologica en Iberoamerica: PrIBES 2000. Trabajos del 1er Taller Iberoamericano de Entomologia Sistemica, Villa de Leyva, Colombia, 28 de junio a 5 de Julio de 1999. 245 - 251.
- Eliopoulos, P.A., Athanasiou, C.G., Buchelos, C.H., Adler, C. Navarro, S., Scholler, M. and Stengard -Hansen, L. (2002).** Occurrence of hymenopterous parasitoids of stored product pests in Greece. Proceedings of the IOBC-WPRS Working Group "Integrated Protection in Stored Products", Lisbon, Portugal, 3-5 September, 2001. Bulletin OILB SROP 25 (3): 127 - 139.
- Eller, F.J., Bartelt, R.J., Kulman, H.M. and Jones, R.L. (1989).** Interaction of prechill and chill durations on diapausing prepupal larvae of the yellowheaded spruce sawfly (Hymenoptera: Tenthredinidae) and associated parasitoids. Annals of the Entomological Society of America 82 (3): 361 - 367.
- Embree, D.G., Raske, A.G. and Wickman, B.E. (1991).** The winter moth *Operophtera brumata* in eastern Canada, 1962-1988. Proceedings of a symposium, Towards integrated pest management of forest defoliators, held at the 18th International Congress of Entomology, Vancouver, Canada in 1988. Forest Ecology and Management 39 (1-4): 47 - 54.
- Emelyanov, I.M. and Goncharenko, G.G. (1991).** Enzyme-linked assay determination of the parasitic pine lappet caterpillars *Dendrolimus pini* L. Doklady, Biological Sciences 316 (1-6): 43 - 46.
- Emre, I. and Yazgan, S. (1990).** Effects of dietary components on reproduction of *Pimpla turionellae* L. (Hymenoptera: Ichneumonidae). Doga, Turk Biyoloji Dergisi 14 (2): 96 - 104.
- England, S. (1995).** Diel emergence patterns of the parasitoid *Bathyplectes curculionis* (Thomson) (Hymenoptera: Ichneumonidae) at two photoperiods. Journal of the Kansas Entomological Society 68 (4): 470 - 473.
- Escribano, A., Williams, T., Goulson, D., Cave, R.D., Chapman, J.W. and Caballero, P. (2000).** Effect of parasitism on a nucleopolyhedrovirus amplified in *Spodoptera frugiperda* larvae parasitized by *Campoletis sonorensis*. Entomologia Experimentalis et Applicata 97 (3): 257 - 264.
- Escribano, A., Williams, T., Goulson, D., Cave, R.D. and Caballero, P. (2000).** Parasitoid - pathogen - pest interactions of *Chelonus insularis*, *Campoletis sonorensis*, and a nucleopolyhedrovirus in *Spodoptera frugiperda* larvae. Biological Control 19 (3): 265 - 273.
- Esmaili, M. and Habibi, J. (1974).** Bioecological investigation on *Hypera postica* in Iran. Proceeding V. Plant Protec. Cong. Iran; Univ. Tabriz, 83-84 pp.
- Evans, D.C. and Zambrano, E. (1991).** Insect damage in maize of highland Ecuador and its significance in small farm pest management. Tropical Pest Management 37 (4): 409 - 414.

- Evans, E.W. and England, S. (1996).** Indirect interactions in biological control of insects: pests and natural enemies in alfalfa. *Ecological Applications* 6 (3): 920 - 930.
- Evans, E.W. and Swallow, J.G. (1993).** Numerical responses of natural enemies to artificial honeydew in Utah alfalfa. *Environmental Entomology* 22 (6): 1392 - 1401.

## F

- Farrar, R.R. Jr., Barbour, J.D. and Kennedy, G.G. (1994).** Field evaluation of insect resistance in a wild tomato and its effects on insect parasitoids. *Entomologia Experimentalis et Applicata* 71 (3): 211 - 226.
- Federici, B.A. (1991).** Viewing polydnviruses as gene vectors of endoparasitic Hymenoptera. *Redia*. 74 (3): Appendix, 387 - 392.
- Federici, B.A., Suduth-Klinger, J.M. and Klinger, J.M.S. (1988).** Evidence for DNA in the calyx particles of *Venturia canescens* (Ichneumonidae). *Colloques de l'INRA* 48: 101.
- Felland, C.M. (1990).** Habitat-specific parasitism of the stalk borer (Lepidoptera: Noctuidae) in northern Ohio. *Environmental Entomology* 19 (1): 162 - 166.
- Feng, J.H., Yan, G.Z., Yao, D.F., Li, G.W. and Zhao, Z.L. (1999).** Studies on insect natural enemy diversity of gypsy moth and their role in natural control of the pest (Lepidoptera: Lymantriidae) in the Beijing area. *Scientia Silvae Sinicae* 35 (2): 50 - 56.
- Ferkovich, S.M., Morales-Ramos, J.A., Rojas, M.G., Oberlander, H., Carpenter, J.E. and Greany, P. (1999).** Rearing of ectoparasitoid *Diapetimorpha introita* on an artificial diet: supplementation with insect cell line derived factors. *BioControl* 44 (1): 29 - 45.
- Ferkovich, S.M., Shapiro, J. and Carpenter, J. (2000).** Growth of a pupal ectoparasitoid, *Diapetimorpha introita*, on an artificial diet: stimulation of growth rate by a lipid extract from host pupae. *BioControl* 45 (4): 401 - 413.
- Fernandez, T.J.L., Garces, G.G., Portuondo, F.E., Valdes, T.P. and Exposito, E.I. (2001).** Insects associated with flowers of weeds of the Botanical Garden of Santiago de Cuba, with emphasis on Hymenoptera. *Revista de Biología Tropical* 49 (3-4): 1013 - 1026.
- Ferrari, R., Burgio, G., Pozzati, M. and Zwakhals, K. (1998).** Records of parasitoids of syrphids (Diptera) in Emilia-Romagna. *Informatore Fitopatologico* 48 (4): 76 - 80.
- Figueiredo, D. and Araujo, J. (1987).** Cabbage (*Brassica oleracea* L.) entomofauna. *Boletim da Sociedade Portuguesa de Entomologia*. 3-12: 82, 1 - 14.
- Figueiredo, E., Mexia, A., Albajes, R. and Sekeroglu, E. (2000).** Parasitoid complex associated with Lepidoptera on horticultural protected crops in the Oeste region of Portugal. IOBC-WPRS Working Group "Integrated Control in Protected Crops, Mediterranean Climate". Proceedings of the meeting, Antalya, Turkey, 24-28 April 2000. *Bulletin OILB SROP* 23 (1): 205 - 208.
- Fincke, O.M., Higgins, L. and Rojas, E. (1990).** Parasitism of *Nephila clavipes* (Araneae: Tetragnathidae) by an ichneumonid (Hymenoptera: Polysphinctini) in Panama. *Journal of Arachnology* 18 (3): 321 - 329.

- Fischer, H.Z. and Patel, P.N. (1993).** Population fluctuation and behaviour of butterflies (Lepidoptera: Lasiocampidae) and their parasitoids (Hymenoptera and Diptera) on avocado (*Persea americana*). *Revista de Biologia Tropical*, publ. 1994, 41 (3A): 515 - 519.
- Fischer, S., Samietz, J., Wackers, F.L. and Dorn, S. (2001).** Interaction of vibrational and visual cues in parasitoid host location. *Journal of Comparative Physiology. A, Sensory, Neural, and Behavioral Physiology* 187 (10): 785 - 791.
- Fitton, M.G. (1984).** Subfamily Metopiinae. In: Gauld, I.D. 1984. *An introduction to the Ichneumonidae of Australia*. London: British Museum (Natural History). 413 pp.
- Fitton, M.G. (1987).** A review of the *Banchus*-group of ichneumon-flies, with a revision of the Australian genus *Philogalleria* (Hymenoptera: Ichneumonidae). 12: 33-45.
- Fitton, M.G. and Gauld, I.D. (1976).** The family-group names of the Ichneumonidae (excluding the Ichneumoninae). *Systematic Entomology* 1: 247-258.
- Fitton, M.G. and Gauld, I.D. (1978).** Further notes on family-group names of Ichneumonidae (Hymenoptera). *Systematic Entomology* 3: 245-247.
- Fitton, M. G., Gauld, I. D. and Shaw, M. R. (1982).** The taxonomy and biology of the British Adelognathinae (Hymenoptera: Ichneumonidae). *Journal of Natural History* 16: 275-283.
- Fitton, M.G., Shaw, M.R. and Gauld, I.D. (1988).** Pimpline ichneumon-flies. Hymenoptera, Ichneumonidae (Pimplinae). *Handbooks for the identification of British Insects* 7 (1): 110 pp
- Fitton, M.G. and Boston, M. (1988).** The British species of *Phthorima* (Hymenoptera: Ichneumonidae). *Entomologist's Gazette* 39 (2): 165 - 170.
- Fitton, M. and Ficken, L. (1990).** British ichneumon-flies of the tribe Oedemopsini (Hymenoptera: Ichneumonidae). *Entomologist* 109 (4): 200 - 214.
- Flanders, K.L., Radcliffe, E.B. and Krueger, C.A. (1994).** Natural enemies of alfalfa weevil, *Hypera postica* (Coleoptera: Curculionidae), in Minnesota. *Great Lakes Entomologist* 27 (1): 7 - 18.
- Flanders, K.L. and Radcliffe, E.B. (2000).** Phenology of the alfalfa weevil (Coleoptera: Curculionidae) and its associated parasitoids in Minnesota. *Journal of Entomological Science* 35 (3): 227 - 238.
- Fleming, J.A.G.W. (1991).** The integration of polydnavirus genomes in parasitoid genomes: implications for biocontrol and genetic analyses of parasitoid wasps. *Biological Control* 1 (2): 127 - 135.
- Fleming, J.A.G.W. (1992).** Polydnaviruses: mutualists and pathogens. *Annual Review of Entomology* 37: 401 - 425.
- Fleming, J.A.G.W. and Summers, M.D. (1991).** Polydnavirus DNA is integrated in the DNA of its parasitoid wasp host. *Proceedings of the National Academy of Sciences of the United States of America* 88: 21, 9770 - 9774.

- Fleming, R.A. (1988).** Difficulties implementing a modelling-based integrated pest management program for alfalfa. *Memoirs of the Entomological Society of Canada* 143: 47 - 59.
- Floate, K., Khan, B. and Gibson, G. (1999).** Hymenopterous parasitoids of filth fly (Diptera: Muscidae) pupae in cattle feedlots. *Canadian Entomologist* 131 (3): 347 - 362.
- Force, D.C. (1989).** Observations on the parasitoids of *Mesepiola specca* Davis (Lepidoptera: Incurvariidae). *Pan Pacific Entomologist* 65 (4): 436 - 439.
- Forster, P.I. (1994).** Diurnal insects associated with the flowers of *Gomphocarpus physocarpus* E. Mey. (Asclepiadaceae), an introduced weed in Australia. *Biotropica* 26 (2): 214 - 217.
- Fox, L.R., Kester, K.M. and Eisenbach, J. (1996).** Direct and indirect responses of parasitoids to plants: sex ratio, plant quality and herbivore diet breadth. Ninth International Symposium on Insect-Plant Relationships held on 24-30 June 1995 in Gwatt, Switzerland. *Entomologia Experimentalis et Applicata* 80 (1): 289 - 292.
- Fox, L.R., Letourneau, D.K., Eisenbach, J. and Van Nouhuys, S. (1990).** Parasitism rates and sex ratios of a parasitoid wasp: effects of herbivore and plant quality. *Oecologia* 83 (3): 414 - 419.
- Fox, L.R. and Eisenbach, J. (1992).** Contrary choices: possible exploitation of enemy-free space by herbivorous insects in cultivated vs. wild crucifers. *Oecologia* 89 (4): 574 - 579.
- Frana, J.E. and O'Neil, R.J. (1993).** Parasitism of late instar larvae of the cattail caterpillar, *Simyra henrici* (Grote) (Lepidoptera: Noctuidae) in Indiana. *Journal of the Kansas Entomological Society* 66 (4): 399 - 404.
- Franca, F.H., De Medeiros, M.A. (1998).** Impact of insecticide mixtures on cabbage production and parasitoids of diamondback moth. *Horticultura Brasileira* 16 (2): 132 - 135.
- Francardi, V. and Covassi, M. (1994).** On the bio-ecology of *Epinotia nigricana* (Herrich-Schaeffer) in fir woods of central Italy (Lepidoptera: Tortricidae). *Redia* 77 (1): 55 - 67.
- Fuester, R.W., Gruber, F., Drea, J.J Jr. and Hoyer, H. (1988).** Parasites of *Lymantria dispar* (Lep.: Lymantriidae) in France and their relation to a biological control program. *Acta Oecologica, Oecologia Applicata* 9 (4): 385 - 402.
- Fuester, R.W., Taylor, P.B. and Eisenberg, A. (1989).** Suitability of prepupae and pupae of the gypsy moth (Lepidoptera: Lymantriidae) as hosts of *Coccygomimus disparis* (Hymenoptera: Ichneumonidae). *Environmental Entomology* 18 (2): 338 - 341.
- Fuester, R.W. and Taylor, P.B. (1991).** Host instar preferences and developmental times of two ichneumonid parasites of the gypsy moth (Lepidoptera: Lymantriidae). *Annals of the Entomological Society of America* 84 (4): 429 - 435.
- Fuester, R.W. and Taylor, P.B. (1993).** Developmental time, oviposition behavior, and incidental mortality by *Coccygomimus disparis* (Hymenoptera: Ichneumonidae), a parasite of the gypsy moth (Lepidoptera: Lymantriidae). *Biological Control* 3 (4): 281 - 287.

- Fuester, R.W. and Taylor, P.B. (1996).** Differential mortality in male and female gypsy moth (Lepidoptera: Lymantriidae) pupae by invertebrate natural enemies and other factors. *Environmental Entomology* 25 (2): 536 - 547.
- Fukuda, H. and Hijii, N. (1996).** Different parasitism patterns of two hymenopterous parasitoids (Ichneumonidae and Ibalidae) depending on the development of *Sirex nitobei* (Hym.: Siricidae). *Journal of Applied Entomology* 120 (5): 301 - 305.
- Furlong, M.J., Verkerk, R.H.J. and Wright, D.J. (1994).** Differential effects of the acylurea insect growth regulator teflubenzuron on the adults of two endolarval parasitoids of *Plutella xylostella*, *Cotesia plutellae* and *Diadegma semiclausum*. *Pesticide Science* 41 (4): 359 - 364.
- Furlong, M.J. and Pell, J.K. (1996).** Interactions between the fungal entomopathogen *Zoophthora radicans* Brefeld (Entomophthorales) and two hymenopteran parasitoids attacking the diamondback moth, *Plutella xylostella* L. *Journal of Invertebrate Pathology* 68 (1): 15 - 21.
- Furlong, M.J. and Pell, J.K. (2000).** Conflicts between a fungal entomopathogen, *Zoophthora radicans*, and two larval parasitoids of the diamondback moth. *Journal of Invertebrate Pathology* 76 (2): 85 - 94.
- Furlong, M.J. and Wright, D.J. (1993).** Effect of the acylurea insect growth regulator teflubenzuron on the endo-larval stages of the hymenopteran parasitoids *Cotesia plutellae* and *Diadegma semiclausum* in a susceptible and an acylurea-resistant strain of *Plutella xylostella*. *Pesticide Science* 39 (4): 305 - 312.

## G

- Gaasch, C.M. (1996).** Flight phenology and species distribution of parasitic wasps in a heterogeneous landscape in Georgia's piedmont, with special reference to the Ichneumoninae and Campopleginae (Hymenoptera: Ichneumonidae). M. S. thesis, Univ. of Georgia, Athens, 200pp.
- Gaasch, C. M. and Pickering, J. (1995).** Flight phenology of parasitic wasps (Hymenoptera: Ichneumonidae) in Georgia's piedmont. (in review).
- Gaasch, C. M., Pickering, J. and Wahl, D. B. (1995).** Distribution and community succession of specialist and generalist parasitic wasps (Hymenoptera: Ichneumonidae) in a heterogeneous landscape in Georgia's piedmont. (in review).
- Gaasch, C.M., Pickering, J. and Moore, C.T. (1998).** Flight phenology of parasitic wasps (Hymenoptera: Ichneumonidae) in Georgia's piedmont. *Environmental Entomology* 27 (3): 606 - 614.
- Gaines, D.N. and Kok, L.T. (1995).** *Cotesia orobena* (Hymenoptera: Braconidae), a gregarious endoparasitoid of *Evergestis rimosalis* (Lepidoptera: Pyralidae), and hyperparasitoids in Virginia Brassica crops. *Biological Control* 5 (4): 573 - 580.
- Gamayunova, S.G. (1988).** Bio-ecological features of the parasites of the green oak-roller *Tortrix viridana*. *Lesovodstvo i Agrolesomelioratsiya* 76: 51 - 53.

- Gantner, M. (2001).** A natural regulation of leafrollers (Lepidoptera: Tortricidae) occurring on hazel shrubs in Lublin Region. *Progress in Plant Protection* 41 (2): 661 - 664.
- Garcia, J.L. (1992).** Hymenopteran parasitoids of *Leptophobia aripa* (Boisduval) (Lepidoptera: Pieridae) in the Cataurito Experimental Station, Zamora County, Aragua State. *Boletin de Entomologia Venezolana* 7 (2): 127 - 131.
- Garcia, R.J.L. (1991).** Effect of diazinon, deltamethrin and *Bacillus thuringiensis* var. *kurstaki* on the control of larvae of Lepidoptera on cabbage. *Boletin de Entomologia Venezolana* 6 (1): 19 - 25.
- Garcia, R.J.L. (1991).** Seasonal variation of the cabbage moth *Plutella xylostella* (L.) (Lepidoptera: Yponomeutidae) and of the parasitoid *Diadegma insulare* Cresson (Hymenoptera: Ichneumonidae), in the Cataurito Experimental Station, Aragua State. *Boletin de Entomologia Venezolana* 6 (1): 27 - 35.
- Garg, D.K. (1989).** *Campoletis chlorideae* Uchida, a larval parasite of *Helicoverpa armigera* (Hub.) in the Kumaon Hills, Uttar Pradesh, India. *International Chickpea Newsletter* 20: 8 - 9.
- Gaston, K.J. and Gauld, I.D. (1993).** How many species of *pimplines* (Hymenoptera: Ichneumonidae) are there in Costa Rica? *Journal of Tropical Ecology* 9 (4): 491 - 499.
- Gates, M.W., Heraty, J.M., Schauff, M.E., Wagner, D.L., Whitfield, J.B. and Wahl, D.B. (2002).** Survey of the parasitic hymenoptera on leafminers in California. *Journal of Hymenoptera Research* 11 (2): 213 - 270.
- Gauld, I.D. (1976).** The classification of the Anomaloninae (Hymenoptera: Ichneumonidae). *Bulletin of the British Museum of Natural History (Entomology)* 33: 1-135.
- Gauld, I.D. (1976).** Description of a new genus of Ichneumonidae (Hymenoptera) of economic importance from Sabah. *Bulletin of Entomological Research* 66: 1-4.
- Gauld, I.D. (1977).** A revision of the Ophioninae (Hymenoptera: Ichneumonidae) of Australia. *Australian Journal of Zoology (Supplementary Series)* 49: 1-112.
- Gauld, I.D. (1977).** *Australoglypta* gen. n. In: Danthanarayana, W., D. Farrugia, & I.D. Gauld. 1977. Studies on the biology and systematic position of a new species of ichneumonid parasitizing the light brown apple moth, *Epiphyas postvittana* (Walker) (Lepidoptera: Tortricidae), in Australia. *Bulletin of Entomological Research* 67: 607-617.
- Gauld, I.D. (1979).** An analysis of the classification of the *Ophion* genus-group (Ichneumonidae). *Systematic Entomology* 5: 59-82.
- Gauld, I.D. (1984).** An introduction to the Ichneumonidae of Australia. London: British Museum (Natural History). 413 pp.
- Gauld, I.D. (1985).** The phylogeny, classification and evolution of parasitic wasps of the subfamily Ophioninae (Ichneumonidae). *Bulletin of the British Museum of Natural History (Entomology)* 51: 1-185.
- Gauld, I.D. (1991).** The Ichneumonidae of Costa Rica, 1. *Memoirs of the American Entomological Institute* 47: 1-589.

- Gauld, I.D. (1995).** Ichneumonidae. *In*: Hanson, P.E. & I.D. Gauld (eds.), The Hymenoptera of Costa Rica. New York: Oxford University Press & The Natural History Museum, London. 893 pp.
- Gauld, I.D. (1997).** The Ichneumonidae of Costa Rica, 2. *Memoirs of the American Entomological Institute* 57: 1-485.
- Gauld, I.D. (2000).** The Ichneumonidae of Costa Rica, 3. *Memoirs of the American Entomological Institute* 63: 1-453.
- Gauld, I.D. and Bradshaw, K. (1997).** The subfamily Anomaloninae. *In*: Gauld, I.D. 1997. The Ichneumonidae of Costa Rica, 2. *Memoirs of the American Entomological Institute* 57: 1-485.
- Gauld, I.D. and Holloway, G.A. (1983).** A new genus of endaseine Ichneumonidae from Australia (Hymenoptera). *Contributions of the American Entomological Institute* 20: 191-197.
- Gauld, I.D. and Holloway, G.A. (1986).** Australian ichneumonids of the tribes Labenini and Poecilocryptini. *Bulletin of the British Museum of Natural History (Entomology)* 53: 107-149.
- Gauld, I.D. and Mitchell, P.A. (1981).** The taxonomy, distribution and host preferences of Indo-Papuan parasitic wasps of the subfamily Ophioninae. Slough: Commonwealth Institute of Entomology. 611 pp.
- Gauld, I.D. and Ward, S. (2000).** Subfamily Brachycyrtinae. *In* Gauld, I.D., The Ichneumonidae of Costa Rica, 3. *Memoirs of the American Entomological Institute* 63:1-453.
- Gauld, I. and Wahl, D. (2000).** The townesioninae: a distinct subfamily of Ichneumonidae (Hymenoptera) or a clade of the Banchinae? *Transactions of the American Entomological Society* 126 (3-4): 279 - 292.
- Gauld, I.D. (1988).** Evolutionary patterns of host utilization by ichneumonid parasitoids (Hymenoptera: Ichneumonidae and Braconidae). *Biological Journal of the Linnean Society* 35 (4): 351 - 377.
- Gauld, I.D., Gomez, J.A.U. and Hanson, P. (1998).** A guide to the Pimplinae of Costa Rica (Hymenoptera: Ichneumonidae). *Revista de Biologia Tropical Supplement* 1, 189 pp.
- Gauld, I.D. and Janzen, D.H. (1994).** The classification, evolution and biology of the Costa Rican species of *Cryptophion* (Hymenoptera: Ichneumonidae). *Zoological Journal of the Linnean Society* 110 (4): 297 - 324.
- Gauld, I.D. and Wahl, D.B. (2000).** The Labeninae (Hymenoptera: Ichneumonidae): a study in phylogenetic reconstruction and evolutionary biology. *Zoological Journal of the Linnean Society* 129 (3): 271 - 347.
- Gauld, I.D. and Wahl, D.B. (2002).** The Eucerotinae: a Gondwanan origin for a cosmopolitan group of Ichneumonidae? *Journal of Natural History* 36: 18, 2229 - 2248.
- Gebremeskel, F.B., Liu, S.S., Wang, X.G. and Shi, Z.H. (2001).** The biology of *Diadromus collaris* (Hymenoptera: Ichneumonidae), a pupal parasitoid of *Plutella xylostella*

- (Lepidoptera: Plutellidae), and its interactions with *Oomyzus sokolowskii* (Hymenoptera: Eulophidae). Bulletin of Entomological Research 91 (6): 461 - 469.
- Gebremeskel, F.B., Liu, S.S. and Shi, Z.H. (2002).** Reproductive compatibility and variation in survival and sex ratio between two geographic populations of *Diadromus collaris*, a pupal parasitoid of the diamondback moth, *Plutella xylostella*. BioControl 47 (6): 625 - 643.
- Gebremeskel, F.B., Shi, Z.H. and Liu, S.S. (2002).** Comparative study on reproductive compatibility of two geographic populations of *Diadromus collaris* (Gravenhors) (Hym.: Ichneumonidae), a pupal parasitoid of *Plutella xylostella* (L.) (Lep.: Plutellidae). Journal of Zhejiang University Agriculture and Life Sciences. 28 (2): 136 - 140.
- Gebremeskel, F.B., Shi, Z.H. and Liu, S.S. (2002).** Comparative study on response to temperature of two geographic populations of *Diadromus collaris* (Gravenhors) (Hym.: Ichneumonidae), a pupal parasitoid of *Plutella xylostella* (L.) (Lep.: Plutellidae). Journal of Zhejiang University Agriculture and Life Sciences 28 (5): 480 - 484.
- Gelman, D.B., Carpenter, J.E. and Greany, P.D. (2000).** Ecdysteroid levels/profiles of the parasitoid wasp, *Diapetimorpha introita*, reared on its host, *Spodoptera frugiperda* and on an artificial diet. Journal of Insect Physiology 46 (4): 457 - 465.
- Gelok, E., McGregor, R., Henderson, D. and Poirier, L. (1998).** Seasonal occurrence and parasitism of *Bucculatrix ainsliella* (Lepidoptera: Lyonetiidae) on *Quercus rubra* in Burnaby, British Columbia. Journal of the Entomological Society of British Columbia 95: 111 - 116.
- Gencer, L. (2003).** The parasitoids of *Yponomeuta malinellus* Zeller (Lepidoptera: Yponomeutidae) in Sivas. Turkish Journal of Zoology 27 (1): 43 - 46.
- Generani, M., Pagliano, G., Scaramozzino, P.L. and Strumia, F. (1998).** New Hymenoptera from Montecristo Island (Tuscany-Italy) (Hymenoptera: Tenthredinidae, Gasteruptiidae, Evaniidae, Ichneumonidae, Chrysididae, Tiphidae, Scoliidae, Formicidae, Sphecidae, Apoidea). Frustula Entomologica 21: 75 - 83.
- Genise, J.F. (1992).** The known species from the Tiphidae (Hym.: Tiphioidea) from Argentina and neighbouring countries. Revista de la Sociedad Entomologica Argentina 50 (1-4): 277 - 329.
- Georgiev, G. (2000).** Studies on larval parasitoids of *Paranthrene tabaniformis* (Rott.) (Lepidoptera: Sesiidae) on urban poplars (*Populus* spp.) in Sofia, Bulgaria. Annals of Forest Science 57 (2): 181 - 186.
- Georgiev, G. (2001).** Notes on the biology and ecology of the parasitoids of the poplar clearwing moth, *Paranthrene tabaniformis* (Rott.) (Lep.: Sesiidae) in Bulgaria. II. *Eriborus terebrans* (Gravenhorst, 1826) (Hym.: Ichneumonidae). Journal of Applied Entomology 125 (6): 289 - 292.
- Georgiev, G. and Delkov, A. (2003).** Bioecological characteristics of *Bassus tumidulus* (Nees) (Hym.: Braconidae), a parasitoid of the poplar twig borer, *Gypsonoma aceriana* (Dup.) (Lep.: Tortricidae) in Bulgaria. Journal of Applied Entomology 127 (2): 99 - 102.



- Georgiev, G. and Kolarov, J. (1999).** New ichneumonidae (Hymenoptera) parasitoids of forest insect pests in Bulgaria. *Anzeiger fur Schadlingskunde* 72 (3): 57 - 61.
- Georgiev, G. and Samuelian, S. (1999).** Species composition, structure and impact of larval parasitoids of poplar twig borer, *Gypsonoma aceriana* (Dup.) (Lepidoptera: Tortricidae), on poplar ornamental trees in Sofia. *Anzeiger fur Schadlingskunde, Pflanzenschutz, Umweltschutz* 72 (1): 1 - 4.
- Georgiev, G.T.S. and Tsankov, G. (1995).** New species of parasitoid insects on larvae of the poplar clearwing (*Paranthrene tabaniformis*) in Bulgaria. *Nauka za Gorata* 32 (2): 51 - 58.
- Gibogini, B., Alma, A. and Arzone, A. (1996).** Bio-ethological research on the Hymenoptera of the biocenosis of *Phyllonorycter robiniellus* (Clemens) (Lepidoptera: Gracillariidae). *Bollettino di Zoologia Agraria e di Bachicoltura* 28 (1): 13 - 22.
- Giles, K.L., Obrycki, J.J., DeGooyer, T.A. and Orr, C.J. (1994).** Seasonal occurrence and impact of natural enemies of *Hypera postica* (Coleoptera: Curculionidae) larvae in Iowa. *Environmental Entomology* 23 (1): 167 - 176.
- Giraldo-Vanegas, H. and Garcia, J.L. (1992).** Determination of the number of instars of *Eiphosoma vitticolle* Cresson (Hymenoptera: Ichneumonidae). *Boletin de Entomologia Venezolana* 7 (2): 133 - 137.
- Giraldo-Vanegas, H. and Garcia, R.J.L. (1994).** Behaviour of adults of *Eiphosoma vitticolle* Cresson (Hymenoptera: Ichneumonidae), parasitoid of *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae). *Boletin de Entomologia Venezolana* 9 (1): 15 - 20.
- Giraldo-Vanegas, H. and Garcia, R.J.L. (1994).** Behaviour, description and development time of the immature stages of *Eiphosoma vitticolle* Cresson (Hymenoptera: Ichneumonidae), parasite of *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae). *Agronomia Tropical Maracay* 44 (4): 645 - 665.
- Giraldo-Vanegas, H. and Garcia, R.J.L. (1995).** Influence of feeding on the capacity for reproduction of *Eiphosoma vitticolle* Cresson (Hymenoptera: Ichneumonidae), parasitoid of *Spodoptera frugiperda* (J.E. Smith). *Agronomia Tropical Maracay* 45 (2): 159 - 170.
- Giunchi, P. and De Giovanni, G. (1987).** Life cycle and symbionts of *Gypsonoma minutana* Hubner (Lepidoptera: Tortricidae) developing on poplar in Emilia-Romagna. *Bollettino dell'Istituto di Entomologia 'Guido Grandi' della Universita degli Studi di Bologna* 41: 195 - 207.
- Glavendekic, M.M. (2000).** Studies on *Altica quercetorum* Foud. (Coleopt.: Chrysomelidae) during an outbreak 1992 - 1995 in Serbia. *Anzeiger fur Schadlingskunde* 73 (5): 127 - 128.
- Glinski, Z. and Jarosz, J. (1996).** Role of polydnavirus in the immunosuppression in insects. *Medycyna Weterynaryjna* 52 (8): 502 - 505.
- Glowacka, B. and Malinowski, H. (1991).** The toxicity of deltamethrin to pine beauty moth *Panolis flammea* Schiff. and its parasitoids *Barichneumon bilunulatus* Grav. and *Rictichneumon pachymerus* Katz. *Pesticide Science* 32 (4): 413 - 417.
- Godfrey, L.D., Godfrey, K.E., Hunt, T.E. and Spomer, S.M. (1991).** Natural enemies of European corn borer *Ostrinia nubilalis* (Hubner) (Lepidoptera: Pyralidae) larvae in

- irrigated and drought-stressed corn. *Journal of the Kansas Entomological Society* 64 (3): 279 - 286.
- Goel, S.C. and Kumar, S. (1994).** Record of a new larval parasitoid associated with *Antigastra catalaunalis* (Dup.) (Lepidoptera: Pyralidae). *Journal of the Bombay Natural History Society* 91 (2): 331.
- Goel, S.C. and Kumar, S. (1994).** Record of a new larval parasitoid associated with *Antigastra catalaunalis* (Dup.) (Lepidoptera: Pyralidae). *Journal of the Bombay Natural History Society* 91 (2): 331.
- Goh, K.S., Berberet, R.C., Young, L.J. and Conway, K.E. (1989).** Mortality of the parasite *Bathyplectes curculionis* (Hymenoptera: Ichneumonidae) during epizootics of *Erynia phytomyi* (Zygomycetes: Entomophthorales) in the alfalfa weevil. *Environmental Entomology* 18 (6): 1131 - 1135.
- Gokhman, V.E. (1988).** Ecological and morphological aspects of the formation and evolution of ichneumonid flies of the subfamily Ichneumoninae (Hymenoptera: Ichneumonidae). *Entomologicheskoe Obozrenie* 67 (4): 821 - 825.
- Gokhman, V.E. (1990).** On the evolution of karyotypes of parasitoids of the subfamily Ichneumoninae (Hymenoptera: Ichneumonidae). *Zoologicheskii Zhurnal* 69 (11): 70 - 80.
- Gokhman, V.E. (1990).** A new genus and species of ichneumon flies of Phaeogenini (Hymenoptera, Ichneumonidae) from Mongolia. *Entomologicheskoye Obozreniye* 69: 671-674.
- Gokhman, V.E. (1992).** On the origin of endoparasitism in the subfamily Ichneumoninae (Hymenoptera: Ichneumonidae). *Zhurnal Obshchei Biologii* 53 (4): 600 - 608.
- Gokhman, V.E. (1995).** Trends of biological evolution in the subfamily Ichneumoninae and related groups (Hymenoptera: Ichneumonidae): an attempt of phylogenetic reconstruction. *Russian Entomological Journal* 4 (1-4): 91 - 103.
- Gokhman, V.E. (1997).** Differential chromosome staining in parasitic wasps of the genus *Dirophanes* (Hymenoptera: Ichneumonidae). *Zoologicheskii Zhurnal* 76 (1): 70 - 73.
- Gokhman, V.E. and Krutov, V.V. (1996).** On the external structure of the male antennae in the subfamily Ichneumoninae (Hymenoptera: Ichneumonidae) and related groups. *Zoologicheskii Zhurnal* 75 (8): 1182 - 1194.
- Gokhman, V.E. and Quicke, D.L.J. (1995).** The last twenty years of parasitic Hymenoptera karyology: an update and phylogenetic implications. *Journal of Hymenoptera Research* 4: 41 - 63.
- Gonzalez, J.R., Caballero, A. and Van Montal, J. (1998).** Environmental gradient of the insect community associated with sugarcane cultivation in Cuba. *Cultivos Tropicales* 19 (3): 69 - 73.
- Goodwin, S. (1979).** Changes in numbers in the parasitoid complex associated with the diamondback moth *Plutella xylostella* (L.) (Lepidoptera) in Victoria. *Australian J. Zool.* 27: 981-989.

- Gordon, D.M., Nisbet, R.M., De Roos, A., Gurney, W.S.C. and Stewart, R.K. (1991).** Discrete generations in host-parasitoid models with contrasting life cycles. *Journal of Animal Ecology* 60 (1): 295 - 308.
- Goyer, R.A., Lenhard, G. and Wei, H.X. (1998).** Abundance of parasitoids of *Archips argyrospila* (Lepidoptera: Tortricidae) in Louisiana. *Environmental Entomology* 27 (3): 726 - 730.
- Graf, V. (1976).** *Pectinorex*, novo gênero de Joppini do sul do Brasil (Hymenoptera - Ichneumonidae). *Dusenía* 9: 71-75.
- Graf, V. (1984).** Ichneumofauna da região sudeste do Brasil. IV. *Neoexetastes*, novo gênero de Banchini (Hymenoptera, Ichneumonidae). *Revista Brasileira de Entomologia* 28: 411-414.
- Graf, V. (1991).** Ichneumonidae (Hymenoptera) do sul do Brasil: um novo gênero de Ctenopelmatinae. *Acta Biol. Par., Curitiba* 20: 157-166.
- Graf, V. (1995).** Note on *Xorides* Latreille (Hymenoptera: Ichneumonidae) in Brazil. *Revista Brasileira de Zoologia* 12 (1): 31 - 35.
- Graf, V. and Marzagao, M.R. (1999).** *Labena fiorii* sp. nov. (Hymenoptera: Ichneumonidae), parasitoid of *Hedypathes betulinus* (Klug), borer of erva-mate, and *Chydarteres striatus* (Fabricius), borer of aroeira, (Coleoptera: Cerambycidae). *Revista Brasileira de Zoologia* 16 (1): 185 - 190.
- Greco, C.F. (1998).** Aphidophagous syrphids (Diptera: Syrphidae): quick identification of pre-imaginal stages and list of natural enemies of the most commonly found species in cereals and pastures in the province of Buenos Aires (Argentina). *Acta Entomologica Chilena* 22: 7 - 11.
- Grijpma, P. and Grijpma, P. (1988).** Parasitoids, parasitism percentages and pathogens of the antler moth, *Cerapteryx graminis* (L.) (Lep.: Noctuidae). Rapport Rijksinstituut voor Onderzoek in de Bos en Landschapsbouw 'De Dorschkamp', Wageningen 516: 51 - 61.
- Grikenis, R.A., Ionaitis, V.P. and Shvitra, D.I. (1988).** Mathematical model of a separate ecosystem of insects. *Trudy Akademii Nauk Litovskoi SSR*, B. 1: 116 - 127.
- Grissell, E.E. (1992).** A revision of *Perissocentrus* Crawford (Hymenoptera: Torymidae). *Journal of Hymenoptera Research* 1 (1): 91 - 102.
- Grissell, E.E. and Zack, R.S. (1996).** Torymidae (Hymenoptera) new to Washington State. *Proceedings of the Entomological Society of Washington* 98 (4): 827 - 828.
- Grochowska, M. (1989).** Observations on *Scambus arundinator* (F.) (Hymenoptera: Ichneumonidae), parasite of the fly *Platycephala planifrons* F. (Diptera: Chloropidae). *Polskie Pismo Entomologiczne* 59 (1): 293 - 297.
- Gross, H.R. and Pair, S.D. (1991).** Seasonal distribution, response to host development stage, and screened-cage performance of *Archytas marmoratus* (Diptera: Tachinidae) and *Ophion flavidus* (Hymenoptera: Ichneumonidae) on *Spodoptera frugiperda* (Lepidoptera: Noctuidae). *Florida Entomologist* 74 (2): 237 - 245.

- Gu, D.J., Waage, J.K. and Wright, D.J. (1994).** Sublethal effects of insecticides on *Plutella xylostella* and its parasitoid *Diadegma eucerocephala*. *Acta Phytophylacica Sinica* 21 (3): 265 - 268.
- Guerra, T.M. and Penteado-Dias, A.M. (2002).** Ichneumonidae (Hymenoptera) abundance in Sao Carlos forest - state of Sao Paulo, Brazil. *Acta Scientiarum* 24 (2): 363 - 368.
- Guerrero, S.M.A., Arretz, V.P., Lamborot, C.L. and Araya, J.E. (1993).** Parasitism of *Heliothis zea* (Boddie) (Lepidoptera: Noctuidae) eggs and larvae on corn in the metropolitan region of Chile. *Acta Entomologica Chilena* 18: 65 - 72.
- Guimaraes, F.R., Vargas-Osuna, E., Maracaja, P.B. and Santiago-Alvarez, C. (1995).** Presence of *Spodoptera exigua* Hb. (Lep.: Noctuidae) and its associated biotic agents in Cordoba province. *Boletin de Sanidad Vegetal, Plagas* 21 (4): 641 - 646.
- Gunasena, G.H., Vinson, S.B., Williams, H.J. and Stipanovic, R.D. (1989).** Development and survival of the endoparasitoid *Campoletis sonorensis* (Hymenoptera: Ichneumonidae) reared from gossypol-exposed *Heliothis virescens* (F.) (Lepidoptera: Noctuidae). *Environmental Entomology* 18 (5): 886 - 891.
- Gunasena, G.H., Vinson, S.B. and Williams, H.J. (1989).** Interrelationships between growth of *Heliothis virescens* (Lepidoptera: Noctuidae) and that of its parasitoid, *Campoletis sonorensis* (Hymenoptera: Ichneumonidae). *Annals of the Entomological Society of America* 82 (2): 187 - 191.
- Gunasena, G.H., Vinson, S.B. and Williams, H.J. (1990).** Effects of nicotine on growth, development, and survival of the tobacco budworm (Lepidoptera: Noctuidae) and the parasitoid *Campoletis sonorensis* (Hymenoptera: Ichneumonidae). *Journal of Economic Entomology* 83 (5): 1777 - 1782.
- Gunasena, G.H. and Harris, M.K. (1988).** Parasites of hickory shuckworm and pecan nut casebearer with five new host-parasite records. *Southwestern Entomologist* 13 (2): 107 - 111.
- Guo, S.J. (1999).** Parasitic natural enemies of the cotton bollworm larva and their relationship to meteorological factors. *Journal of Henan Agricultural Sciences* 10: 17 - 19.
- Guo, Y.J. and Heong, K.L. (1992).** An evaluation of the suppressive effect of larval parasitization on populations of rice leaffolders. *Chinese Journal of Biological Control* 8 (1): 1 - 5.
- Gupta, S. and Gupta, V.K. (1984).** *Lagarosoma*, a new Brazilian genus of the tribe Gabuniini (Hymenoptera: Ichneumonidae). *International Journal of Entomology* 26: 235-240.
- Gupta, S. and Gupta, V.K. (1985).** The type species of *Ahilya* (Hymenoptera: Ichneumonidae). *Oriental Insects* 18: 24.
- Gupta, V.K. (1970).** Ichneumon hunting in India. Delhi: privately published. 80 pp.
- Gupta, V.K. (1980).** A revision of the tribe Poemeniini in the Oriental Region (Hymenoptera: Ichneumonidae). *Oriental Insects* 14: 73-130.

- Gupta, V.K. (1985).** A review of the tribe Neoxoridini of the World (Hymenoptera: Ichneumonidae: Pimplinae). *Oriental Insects* 19: 323-329.
- Gupta, V.K. (1987).** The Ichneumonidae of the Indo-Australian area (Hymenoptera). *Memoirs of the American Entomological Institute* 41: 1-1210.
- Gupta, V.K. (1988).** Relationships of the genera of the tryphonine tribe Oedemopsini and a revision of *Acaenitellus* Morley. In: V.K.Gupta (ed.), *Advances in Parasitic Hymenoptera Research*. Leiden/New York: E.J.Brill. 546 pp.
- Gupta, V.K. (1989).** New genera of porizontine Ichneumonidae from Florida (Hymenoptera). *Florida Entomologist* 72: 284-294.
- Gupta, V.K. and Gupta, M.L. (1971).** A new porizontine genus *Urvashia* from India (Hymenoptera: Ichneumonidae). *Oriental Insects* 4: 483-487.
- Gupta, V.K. and Gupta, M.L. (1971).** A new porizontine genus *Menaka* from India (Hymenoptera: Ichneumonidae). *Oriental Insects* 5: 111-116.
- Gupta, V.K. and Gupta, M.L. (1976).** A new porizontine genus *Kartika* from India and Burma (Hymenoptera: Ichneumonidae). *Oriental Insects* 10: 459-496.
- Gupta, V.K. and Jonathan, J.K. (1970).** The genera of the *Dagathia* complex (Hymenoptera: Ichneumonidae). *Oriental Insects* 3: 389-393.
- Gupta, V.K. and Jonathan, J.K. (1971).** A new mesostenine genus *Tanepomidos* from India (Hymenoptera: Ichneumonidae). *Oriental Insects* 5: 145-148.
- Gupta, V.K. and Tikar, D.T. (1969).** Taxonomic identity of pimpline genera *Flavopimpla* Betrem and *Afrephialtes* Benoit (Hymenoptera: Ichneumonidae). *Oriental Insects* 3: 269-277.
- Gupta, V.K. and Tikar, D.T. (1976).** *Ichneumonologia Orientalis*, or a monographic study of the Ichneumonidae of the Oriental Region, part 1. The tribe Pimplini (Hymenoptera: Ichneumonidae: Pimplinae). *Oriental Insects Monographs* 1: 1-312.
- Gupta, M. and Pawar, A.D. (1992).** Biological wealth in rice ecosystem. *Plant Protection Bulletin Faridabad* 44 (3): 6 - 15.
- Gupta, S. (1988).** New distributional records for Ichneumoninae (Hymenoptera: Ichneumonidae) for the Indo-Australian area. *Oriental Insects* 22: 301 - 357.
- Gupta, S.C., Yazdani, S.S., Hameed, S.F. and Agarwal, M.L. (1994).** Seasonal prevalence of larval and pupal parasitoids of sugarcane top borer, *Scirpophaga excerptalis* in North Bihar (India). *Journal of Entomological Research* 18 (1): 83 - 87.
- Gupta, V. (1987).** A revision of the genus *Microcharops* (Hymenoptera: Ichneumonidae). *Contributions of the American Entomological Institute* 23 (2): 1 - 42.
- Gupta, V. (1999).** A new genus and species of mesostenine Ichneumonidae from Malaysia (parasitic Hymenoptera). *Oriental Insects* 33: 95 - 98.
- Gupta, V. and Gupta, S. (1992).** Supplementary list of ichneumonid references No. 1 (Hymenoptera: Ichneumonidae). *Oriental Insects* 26: 335 - 365.
- Gupta, V. and Gupta, S. (1993).** Supplementary list of ichneumonid references, No. 2 (Hymenoptera: Ichneumonidae). *Oriental Insects* 27: 385 - 402.

- Gupta, V.K. (1991).** A review of the exenterine genus *Eridolius* (Hymenoptera: Ichneumonidae) and descriptions of new species from the Oriental Region. *Oriental Insects* 25: 435 - 446.
- Gupta, V.K. (1991).** The parasitic Hymenoptera and biological control of the African Ichneumonidae. *Insect Science and its Application* 12 (1-3): 9 - 18.
- Gupta, V.K. (1993).** The exenterine ichneumonids (Hymenoptera: Ichneumonidae) of China. *Japanese Journal of Entomology* 61 (3): 425 - 441.
- Gupta, V.K. (1999).** A review of the mesostenine genus *Amauromorpha* (Hymenoptera: Ichneumonidae). *Oriental Insects* 33: 267 - 277.
- Gupta, V.K. (2002).** A new genus of *Glyptini* from India (Hymenoptera: Ichneumonidae). *Oriental Insects* 36: 239 - 243.
- Gupta, V.K. (2002).** *Glyptopimpla* Morley (Hymenoptera: Ichneumonidae: Banchinae) - a valid genus with descriptions of new species from the Orient. *Oriental Insects* 36: 221 - 237.
- Gupta, V.K. (1988).** Relationships of the genera of the tryphonine tribe Oedemopsini and a revision of *Acaenitellus* Morley (Hymenoptera: Ichneumonidae: Tryphoninae). *Advances in parasitic Hymenoptera research: Proceedings of the II. Conference on the Taxonomy and Biology of Parasitic Hymenoptera. recd. 1993*, 243 - 258.
- Gur'-yanova, T.M. (1986).** The role of insect behaviour. *Zashchita Rastenii* 2: 30.
- Gur'-yanova, T.M. (1990).** Role of the spatial structure of host and parasite populations in the population dynamics of the sawfly *Neodiprion sertifer*. *Lesovedenie* 6: 67 - 76.
- Gur'-yanova, T.M. (1991).** Principles of integrated control of defoliators in forest biocoenoses. *Lesnoe Khozyaistvo* 8: 8 - 10.
- Güven, T. and Yel, M. (1990).** The ultrastructure of the venom gland in *Pimpla turionellae* (Hymenoptera: Ichneumonidae). *Doga, Turk Biyoloji Dergisi* 14 (1): 17 - 31.

## H

- Habibi, J. (1977).** The bioecology of alfalfa weevil and its control in Karaj. *Ibid.*, 44: 23-41.
- Hagley, E.A.C., Biggs, A.R., Timbers, G.E. and Coutu-Sundy, J. (1993).** Effect of age of the puparium of the apple maggot, *Rhagoletis pomonella* (Walsh) (Diptera: Tephritidae), on parasitism by *Phygadeuon wiesmanni* Sachtl. (Hymenoptera: Ichneumonidae). *Canadian Entomologist* 125 (4): 721 - 724.
- Hagley, E.A.C. and Barber, D.R. (1991).** Foliage-feeding Lepidoptera and their parasites recovered from unmanaged apple orchards in southern Ontario. *Proceedings of the Entomological Society of Ontario* 122: 1 - 7.
- Hailemichael, Y., Smith, J.W. Jr. and Wiedenmann, R.N. (1994).** Host-finding behavior, host acceptance, and host suitability of the parasite *Xanthopimpla stemmator*. *Entomologia Experimentalis et Applicata* 71 (2): 155 - 166.

- Hailemichael, Y. and Smith, J.W. Jr. (1994).** Development and longevity of *Xanthopimpla stemmator* (Hymenoptera: Ichneumonidae) at constant temperature. *Annals of the Entomological Society of America* 87 (6): 874 - 878.
- Hanks, L.M., Campbell, C., Paine, T.D. and Millar, J.G. (1997).** Host range expansion of *Helcostizus rufiscutum* Cushman (Hymenoptera: Ichneumonidae) to *Phoracantha semipunctata* Fabr. (Coleoptera: Cerambycidae) [on *Eucalyptus globulus*] in California. *Pan Pacific Entomologist* 73 (3): 190 - 191.
- Hanski, I. and Van Nouhuys, S. (2000).** Apparent competition between parasitoids mediated by a shared hyperparasitoid. *Ecology Letters* 3 (2): 82 - 84.
- Hanski, I. and Van Nouhuys, S. (2002).** Colonization rates and distances of a host butterfly and two specific parasitoids in a fragmented landscape. *Journal of Animal Ecology* 71 (4): 639 - 650.
- Happe, M., Kirchoff, K., Wassenegger-Wittlich, G. and Madel, G. (1988).** The biology of the parasitic wasp *Diadegma semiclausum* Hellen (Hym.: Ichneumonidae) and its host *Plutella xylostella* Curtis (Lep.: Yponomeutidae). *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 6 (1-3): 146 - 149.
- Harcourt, D.G. and Ellis, C.R. (1991).** Distribution in southern Ontario of *Bathyplectes anurus* (Hymenoptera: Ichneumonidae), a larval parasitoid of the alfalfa weevil. *Proceedings of the Entomological Society of Ontario* 122: 41-45.
- Harrell, M.O., Jones, J.A. and Brohman, M.A. (1996).** Life histories and parasitoids of *Dioryctria borers* (Lepidoptera: Pyralidae) of pines in Nebraska. *Journal of the Kansas Entomological Society* 69 (4): 279 - 284.
- Harrington, S.A., Hutchinson, P., Dutch, M.E., Lawrence, P.J. and Michael, P.J. (1993).** An efficient method of mass rearing two introduced parasitoids of noctuids (Lepidoptera: Noctuidae). *Journal of the Australian Entomological Society* 32 (1): 79 - 80.
- Harris, R.J. and Read, P.E.C. (1999).** Enhanced biological control of wasps. *Science for Conservation* 115: 39 pp.
- Harris, R.J. and Rose, E.A.F. (1999).** Factors influencing reproductive strategies of the vespid parasitoid *Sphecophaga vesparum vesparum* (Hymenoptera: Ichneumonidae). *New Zealand Journal of Zoology* 26 (2): 89 - 96.
- Harvey, I., Marris, G. and Hubbard, S. (1988).** Adaptive patterns in the avoidance of superparasitism by solitary parasitic wasps. *Colloques de l'INRA* 48: 137 - 142.
- Harvey, J.A. (1996).** *Venturia canescens* parasitizing *Galleria mellonella* and *Anagasta kuehniella*: is the parasitoid a conformer or regulator? *Journal of Insect Physiology* 42 (11-12): 1017 - 1025.
- Harvey, J.A., Harvey, I.F. and Thompson, D.J. (1993).** The effect of superparasitism on development of the solitary parasitoid wasp, *Venturia canescens* (Hymenoptera: Ichneumonidae). *Ecological Entomology* 18 (3): 203 - 208.
- Harvey, J.A., Harvey, I.F. and Thompson, D.J. (1994).** Flexible larval growth allows use of a range of host sizes by a parasitoid wasp. *Ecology* 75 (5): 1420 - 1428.

- Harvey, J.A., Harvey, I.F. and Thompson, D.J. (1995).** The effect of host nutrition on growth and development of the parasitoid wasp *Venturia canescens*. *Entomologia Experimentalis et Applicata* 75 (3): 213 - 220.
- Harvey, J.A., Harvey, I.F. and Thompson, D.J. (2001).** Lifetime reproductive success in the solitary endoparasitoid, *Venturia canescens*. *Journal of Insect Behavior* 14 (5): 573 - 593.
- Harvey, J.A., Thompson, D.J. and Heyes, T.J. (1996).** Reciprocal influences and costs of parasitism on the development of *Corcyra cephalonica* and its endoparasitoid *Venturia canescens*. *Entomologia Experimentalis et Applicata* 81 (1): 39 - 45.
- Harvey, J.A., Van Dam, N.M. and Gols, R. (2003).** Interactions over four trophic levels: foodplant quality affects development of a hyperparasitoid as mediated through a herbivore and its primary parasitoid. *Journal of Animal Ecology* 72 (3): 520 - 531.
- Harvey, J.A. and Thompson, D.J. (1994).** Some factors affecting host suitability for the solitary parasitoid wasp, *Venturia canescens* (Hymenoptera: Ichneumonidae). *Norwegian Journal of Agricultural Sciences, SUPP* 16: 321 - 327.
- Harvey, J.A. and Thompson, D.J. (1995).** Developmental interactions between the solitary endoparasitoid *Venturia canescens* (Hymenoptera: Ichneumonidae), and two of its hosts, *Plodia interpunctella* and *Corcyra cephalonica* (Lepidoptera: Pyralidae). *European Journal of Entomology* 92 (2): 427 - 435.
- Harvey, J.A. and Thompson, D.J. (1995).** Host behaviour and its influence on foraging and acceptance by the solitary parasitoid wasp, *Venturia canescens* (Hym.: Ichneumonidae). *Entomophaga* 40 (2): 193 - 210.
- Harvey, J.A. and Vet, L.E.M. (1997).** *Venturia canescens* parasitizing *Galleria mellonella* and *Anagasta kuehniella*: differing suitability of two hosts with highly variable growth potential. *Entomologia Experimentalis et Applicata* 84 (1): 93 - 100.
- Has, A., Dortbudak, N., Caliskaner, S., Erdogan, P. and Aydemir, M. (1995).** Investigations on the distribution, infestation rates, population densities, damage, biological and ecological factors for the control of potato tuber moth, *Phthorimaea operculella* (Zeller), and its control in the central Anatolia region. *Zirai Mucadele Arastirma Yilligi*. 30: 17 - 18.
- Haseeb, M., Amano, H., Nemoto, H., Vogt, H. and Heimbach, U. (2000).** Effects of pesticides on the mortality and beneficial capacity of *Diadegma semiclausum* (Hymenoptera: Ichneumonidae), a primary parasitoid of the diamondback moth. Working group "Pesticides and beneficial organisms", Versailles, France, 27 - 29 October 1999. *Bulletin OILB SROP* 23 (9): 39 - 48.
- Haseeb, M., Amano, H. and Nemoto, H. (2000).** Pesticidal effects on mortality and parasitism rates of *Diadegma semiclausum*, a parasitoid of the diamondback moth. *BioControl* 45 (2): 165 -178.



- Hastings, F.L., Hain, F.P. and Odell, T.M. (2002).** A survey of parasitoids and other organisms affecting gypsy moth (Lepidoptera: *Lymantria dispar* L.) along the leading edge of its southward movement. *Journal of Entomological Science* 37 (2): 207 - 209.
- Hatakeyama, M., Sawa, M., Oishi, K. and Lee, J.M. (2000).** Artificial reproduction in a hymenopteran insect, *Athalia rosae*, using eggs matured with heterospecific yolk proteins and fertilized with cryopreserved sperm. *Archives of Insect Biochemistry and Physiology* 43 (3): 137 - 144.
- Haugen, D.A. and Underdown, M.G. (1990).** Release of parasitoids for *Sirex noctilio* control by transporting infested logs. *Australian Forestry* 53 (4): 266 - 270.
- Haugen, D.A. and Underdown, M.G. (1990).** *Sirex noctilio* control program in response to the 1987 Green Triangle outbreak. *Australian Forestry* 53 (1): 33 - 40.
- Hauser, E. (1994).** Ecology of the parasitoids of *Taleporia tubulosa* (Hymenoptera: Ichneumonoidea/ Lepidoptera: Psychidae). *Entomologia Generalis* 18 (3-4): 227 - 234.
- Havron, A. and Margalith, Y. (1995).** Parasitization of *Vespa orientalis* nests by *Sphēcophaga vesparum* Curtis in southern Israel (Hymenoptera: Vespidae, Ichneumonidae). *Phytoparasitica*. 23 (1): 19 - 25.
- Hawkeswood, T.J. (1994).** Observations on the biology and host plants of the Australian longicorn beetle, *Ceresium seminigrum* Aurivillius (Coleoptera: Cerambycidae). *Giornale Italiano di Entomologia* 7: 36, 5 - 10.
- Hawkeswood, T.J. (1994).** Observations on the biology and host plants of the Australian longicorn beetle, *Ceresium seminigrum* Aurivillius (Coleoptera: Cerambycidae). *Giornale Italiano di Entomologia* 7: 36, 5 - 10.
- He, B., Peng, S.G., He, G.Y., Wei, Y.L. and Tan, X.Q. (1991).** Bionomics of *Actias selene ningpoana* Felder and its control. *Insect Knowledge* 28 (6): 353 - 354.
- He, J.H. (1991).** Three new species of *Megalomya* Uchida from China (Hymenoptera: Ichneumonidae). *Oriental Insects* 25: 145 - 153.
- He, J.H. (1992).** A supplementary catalogue of Ichneumonidae from China (Hymenoptera: Ichneumonidae). *Oriental Insects* 26: 275 - 334.
- He, J.H., Chen, X.X. and Ma, Y. (2001).** A new species of the genus *Rothneyia* Cameron (Hymenoptera: Ichneumonidae) from Xizang, China. *Entomologia Sinica* 8 (2): 111 - 114.
- He, J.H. and Li, Q.C. (1995).** A new species of *Diaparis* Foerster (Hymenoptera: Ichneumonidae: Tersilochinae) from Gansu, China. *Entomotaxonomia* 17 (4): 303 - 305.
- He, K.L., Wang, Z.Y., Wen, L.P., Bai, S.X., Liu, K.H. and Zhou, D.R. (2002).** Investigation on the parasitoids and pathogens of Asian corn borer overwintering larvae in the corn belt of China. *Chinese Journal of Biological Control* 18 (2): 49 - 53.
- He, Y.H., Chen, S.C., Chen, H.S., Wu, J.W., Yan, J.J., Xu, C.H. and Yu, D.F. (1988).** A survey on the natural enemies of Chinese pine caterpillar (*Dendrolimus tabulaeformis* Tsai et Liu) in the Beijing area. *Chinese Journal of Biological Control* 4 (1): 21 - 24.

- Hedderwick, M.P., El. Agoze, M. and Periquet, G. (1988).** Enzymatic polymorphism, inbreeding and occurrence of diploid males in *Diadromus pulchellus* (Hym.: Ichneumonidae). Colloques de l'INRA 48: 109 - 110.
- Hedstrom, L. (1988).** Swedish insect records - report 4. Entomologisk Tidskrift 109 (3-4): 139 - 149.
- Hedstrom, L. (1990).** Swedish insect records - report 6. Entomologisk Tidskrift 111 (4): 133 - 147.
- Heinlein, G., Scholler, M., Prozell, S., Reichmuth, C., Adler, C., Navarro, S., Scholler, M. and Stengard-Hansen, L. (2002).** Oviposition of *Venturia canescens* (Gravenhorst) (Hymenoptera: Ichneumonidae) parasitizing the Indian meal moth *Plodia interpunctella* (Hubner) (Lepidoptera: Pyralidae). Proceedings of the IOBC-WPRS Working Group "Integrated Protection in Stored Products", Lisbon, Portugal, 3-5 September, 2001. Bulletin OILB SROP 25 (3): 109 - 114.
- Heinrich, G.H. (1965).** Burmesische Ichneumoninae I. Entomologisk Tidskrift 86: 1-130.
- Heinrich, G.H. (1965).** Burmesische Ichneumoninae II. Entomologisk Tidskrift 86: 133-177.
- Heinrich, G.H. (1966).** Burmesische Ichneumoninae III. Entomologisk Tidskrift 87: 184-247.
- Heinrich, G.H. (1967).** Synopsis and reclassification of the Ichneumoninae Stenopneusticae of Africa south of the Sahara (Hymenoptera). Vol. I. Altötting, Germany: Farmington State College Press.
- Heinrich, G.H. (1967).** Synopsis and reclassification of the Ichneumoninae Stenopneusticae of Africa south of the Sahara (Hymenoptera). Vol. II. Altötting, Germany: Farmington State College Press.
- Heinrich, G.H. (1968).** Burmesische Ichneumoninae IV. Entomologisk Tidskrift 89: 77-106.
- Heinrich, G.H. (1968).** Burmesische Ichneumoninae V. Entomologisk Tidskrift 89: 197-228.
- Heinrich, G.H. (1968).** Synopsis and reclassification of the Ichneumoninae Stenopneusticae of Africa south of the Sahara (Hymenoptera). Vol. III. Altötting, Germany: Farmington State College Press.
- Heinrich, G.H. (1968).** Contributions to the knowledge of North American Ichneumoninae. Naturaliste Canadien 95: 703-722.
- Heinrich, G.H. (1969).** Burmesische Ichneumoninae VI. Entomologisk Tidskrift 90: 100-130.
- Heinrich, G.H. (1969).** Synopsis and reclassification of the Ichneumoninae Stenopneusticae of Africa south of the Sahara (Hymenoptera). Vol. IV. Altötting, Germany: Farmington State College Press.
- Heinrich, G.H. (1969).** Synopsis and reclassification of the Ichneumoninae Stenopneusticae of Africa south of the Sahara (Hymenoptera). Vol. V. Altötting, Germany: Farmington State College Press.

- Heinrich, G.H. (1971).** Synopsis of Nearctic Ichneumoninae Stenopneusticae with particular reference to the northeastern region (Hymenoptera). Supplement 2. *Naturaliste Canadien* 98: 959-1026.
- Heinrich, G.H. (1972).** Synopsis of Nearctic Ichneumoninae Stenopneusticae with particular reference to the northeastern region (Hymenoptera). Supplement 3. *Naturaliste Canadien* 98: 173-211.
- Heinrich, G.H. (1972).** Zur Systematik der Ichneumoninae Stenopneusticae IX. Eine Spätlese (Hymenoptera, Ichneumonidae). *Mitteilungen. Münchener. Entomologischen Gesellschaft* 60: 80-101.
- Heinrich, G.H. (1973).** Synopsis of Nearctic Ichneumoninae Stenopneusticae with particular reference to the northeastern region (Hymenoptera). Supplement 4. *Naturaliste Canadien* 99: 461-465.
- Heinrich, G.H. (1974).** Burmesische Ichneumoninae VIII. *Annales Zoologici* 31: 407-457.
- Heinrich, G.H. (1974).** Burmesische Ichneumoninae IX. *Annales Zoologici* 32: 103-198.
- Heinrich, G.H. (1975).** Burmesische Ichneumoninae X. *Annales Zoologici* 32: 441-514.
- Heinrich, G.H. (1977).** Ichneumoninae of Florida and neighboring states (Hymenoptera: Ichneumonidae, subfamily Ichneumoninae). 9: 1-350.
- Heinrich, G.H. (1978).** Eastern Palearctic Ichneumoninae. (in Russian). Leningrad: Academy of Science. 80 pp.
- Heinrich, G.H. (1980).** Burmesische Ichneumoninae XI. Tribus Ichneumoninae (5. teil). Subtribus Craticheumonina Heinrich. *Annales Zoologici* 35: 117-192.
- Heitland, W. and Pschorn-Walcher, H. (1992).** A brief note on unusually high superparasitism in some sawfly parasitoids (Hym.: Ichneumonidae: Euryproctini). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 65 (1-2): 75 - 79.
- Hekal, A.M. (1986).** Percentage of parasitism and sex ratio in different parasites of the diapausing larvae of *Pectinophora gossypiella* Saund. *Annals of Agricultural Science, Moshtohor* 24 (4): 2213 - 2221.
- Heliovaara, K., Vaisanen, R. and Varama, M. (1991).** Larval mortality of pine sawflies (Hymenoptera: Diprionidae) in relation to pollution level: a field experiment. *Entomophaga* 36 (2): 315 - 321.
- Hellers, M., Beck, M., Theopold, U., Kamel, M. and Schmidt, O. (1996).** Multiple alleles encoding a virus-like particle protein in the ichneumonid endoparasitoid *Venturia canescens*. *Insect Molecular Biology* 5 (4): 239 - 249.
- Hemerik, L. and Harvey, J.A. (1999).** Flexible larval development and the timing of destructive feeding by a solitary endoparasitoid: an optimal foraging problem in evolutionary perspective. *Ecological Entomology* 24 (3): 308 - 315.
- Hemming, M.F. (1945).** *Opinion* 159. 2: 275-290.
- Henaut, A. (1990).** Study of the sound produced by *Pimpla instigator* (Hymenoptera: Ichneumonidae) during host selection. *Entomophaga* 35 (1): 127 - 139.

- Hendrickson, R.M. Jr., Gruber, F., Mailloux, G. and Drea, J.J. (1991).** Parasite colonizations against *Crioceris asparagi* (L.) and *C. duodecimpunctata* (L.) (Coleoptera: Chrysomelidae) in North America from 1983 to 1988. *Proceedings of the Entomological Society of Washington* 93 (1): 67 - 69.
- Henneberry, T.J., Vail, P.V., Pearson, A.C. and Sevacherian, V. (1991).** Biological control agents of noctuid larvae (Lepidoptera: Noctuidae) in the Imperial Valley of California. *Southwestern Entomologist* 16 (1): 81 - 89.
- Herard, F. and Prevost, G. (1997).** Suitability of *Yponomeuta malinellus* and *Y. cagnagellus* (Lepidoptera: Yponomeutidae) as hosts of *Diadegma armillata* (Hymenoptera: Ichneumonidae). *Environmental Entomology* 26 (4): 933 - 938.
- Herz, A. and Heitland, W. (1995).** The role of different guilds of parasitoids of *Diprion pini* (Hymenoptera: Diprionidae) at times of low host density. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 10 (1-6): 59 - 62.
- Herz, A. and Heitland, W. (1999).** Larval parasitism of a forest pest, the common pine sawfly *Diprion pini* (L.) (Hym.: Diprionidae), during an endemic density phase. *Journal of Applied Entomology* 123 (3): 129 - 137.
- Heyer, W. (1992).** The parasitization of cereal leaf beetles, *Oulema* spp., in the Halle region. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 8 (1-3): 87 - 89.
- Higashiura, Y., Raske, A.G. and Wickman, B.E. (1991).** Pest control in natural and man-made forests in northern Japan. *Proceedings of a symposium, Towards integrated pest management of forest defoliators, held at the 18th International Congress of Entomology, Vancouver, Canada in 1988. Forest Ecology and Management* 39 (1-4): 55 - 64.
- Hildebrand, M., Dickler, E. and Geider, K. (2000).** Occurrence of *Erwinia amylovora* on insects in a fire blight orchard. *Journal of Phytopathology* 148 (4): 251 - 256.
- Hilgarth, R.S. and Webb, B.A. (2002).** Characterization of *Campoletis sonorensis* ichnovirus segment I genes as members of the repeat element gene family. *Journal of General Virology* 83 (10): 2393 - 2402.
- Hill, T.A. and Foster, R.E. (2000).** Effect of insecticides on the diamondback moth (Lepidoptera: Plutellidae) and its parasitoid *Diadegma insulare* (Hymenoptera: Ichneumonidae). *Journal of Economic Entomology* 93 (3): 763 - 768.
- Hill, T.A. and Foster, R.E. (2003).** Influence of selected insecticides on the population dynamics of diamondback moth (Lepidoptera: Plutellidae) and its parasitoid, *Diadegma insulare* (Hymenoptera: Ichneumonidae), in cabbage. *Journal of Entomological Science* 38 (1): 59 - 71.
- Hilpert, H. (1992).** Zur Systematik der Gattung *Ichneumon* Linnaeus, 1758 in der Westpaläarktis (Hymenoptera, Ichneumonidae, Ichneumoninae). *Entomofauna (Supplement 6)*: 1-389.

- Hilszczanski, J. (1994).** Impact of insecticides, applied for the control of the nun moth (*Lymantria monacha*) on ichneumonids (Ichneumonidae: Hymenoptera). *Sylwan* 138 (5): 43 - 49.
- Hilszczanski, J., Grodzki, W., Knizek, and Forster, B. (1998).** Effects of control treatment on population dynamics of koino- and idiobiont ichneumonids. *Methodology of forest insect and disease survey in Central Europe. Proceedings, First Workshop of the IUFRO WP 7.03.10, Ustron Jaszowiec, Poland, 21-24 April, 1998.* 81 - 83.
- Hintze-Podufal, C. and Thiele, C (1988).** Investigations on the microfauna of rose bedeguar galls. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 6: (4-6): 537 - 541.
- Hinz, R. (1987).** Rearing species of the genus *Ichneumon* Linnaeus and allied genera for biological investigations (Hymenoptera: Ichneumonidae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 60 (1-2): 113 - 120.
- Hinz, R. (1991).** Investigations on the biology of some species of Ichneumonini (Hymenoptera: Ichneumonidae). *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 64 (1-2): 190 - 114.
- Hirashima, Y., Abe, M., Tadauchi, O., Konishi, K. and Maeto, K. (1989).** The hymenopterous parasitoids of the diamondback moth, *Plutella xylostella* (Lepidoptera: Yponomeutidae) in Japan. *Esakia* 28: 63 - 73.
- Hirashima, Y., Nohara, K. and Miura, T. (1990).** Studies on the biological control of the diamondback moth, *Plutella xylostella* (Linnaeus). 1. Insect natural enemies and their utilization. *Science Bulletin of the Faculty of Agriculture, Kyushu University* 44 (3): 65 - 70.
- Hoballah, M.E.F. and Turlings, T.C.J. (2001).** Experimental evidence that plants under caterpillar attack may benefit from attracting parasitoids. *Evolutionary Ecology Research* 3 (5): 553 - 565.
- Hoch, G., Gschwantner, T., Schopf, A. and Villemant, C. (1999).** The gypsy moth and its natural enemies at different population densities in the oak forests of Eastern Austria. Integrated protection of *Quercus* spp. forests. *Proceedings of the meeting held at Rabat Sale, Morocco, 26 - 29 October, 1998. Bulletin OILB SROP* 22 (3):121 - 128.
- Hoch, G., Zubrik, M., Novotny, J. and Schopf, A. (2001).** The natural enemy complex of the gypsy moth, *Lymantria dispar* (Lep.: Lymantriidae) in different phases of its population dynamics in eastern Austria and Slovakia - a comparative study. *Journal of Applied Entomology* 125 (5): 217 - 227.
- Hodges, A.C., Hodges, G.S. and Espelie, K.E. (2003).** Parasitoids and parasites of *Polistes metricus* Say (Hymenoptera: Vespidae) in Northeast Georgia. *Annals of the Entomological Society of America* 96 (1): 61 - 64.
- Hoffmeister, T. (1992).** Factors determining the structure and diversity of parasitoid complexes in tephritid fruit flies. *Oecologia* 89 (2): 288 - 297.

- Hohmann, C.L. and Meneguim, A.M. (1993).** Preliminary observations on the occurrence of avocado borer, *Stenoma catenifer* Wals in the state of Parana. Anais da Sociedade Entomologica do Brasil 22 (2): 417 - 419.
- Hokkanen, H., Husberg, G.B. and Soderblom, M. (1988).** Natural enemy conservation for the integrated control of the rape blossom beetle *Meligethes aeneus* F. Annales Agriculturae Fenniae 27 (4): 281 - 194.
- Hokkanen, H.M.T. (1989).** Biological and agrotechnical control of the rape blossom beetle *Meligethes aeneus* (Coleoptera: Nitidulidae). Acta Entomologica Fennica 53: 25 - 29.
- Holtkamp, R.H. (1993).** Parasitoids of *Comostolopsis germana* Prout (Lepidoptera: Geometridae). Journal of the Australian Entomological Society 32 (4): 387 - 388.
- Hondo, M. (1992).** Mortality of post-overwintered larvae of the Mulberry tiger moth, *Thanatarctia imparilis* (Butler) (Lepidoptera: Arctiidae) caused by parasitoids and pathogens. Applied Entomology and Zoology 27 (4): 595 - 598.
- Hopper, K.R. and Roush, R.T. (1993).** Mate finding, dispersal, number released, and the success of biological control introductions. Ecological Entomology 18 (4): 321 - 331.
- Horstmann, K. (1969).** Typenrevision der europäischen Arten der Gattung *Diadegma* Förster (syn. *Angitia* Holmgren). Beiträge zur Entomologie 19: 413-472.
- Horstmann, K. (1970).** Bemerkungen zur Systematik einiger Gattungen der Campopleginae (Hymenoptera, Ichneumonidae). Nachrichtenblatt der Bayerischen Entomologen 19: 77-84.
- Horstmann, K. (1971).** Revision der europäischen Tersilochinen (Hymenoptera, Ichneumonidae). Teil I. Veröffentlichungen der Zoologischen Staatssammlung München 15: 47-138.
- Horstmann, K. (1973).** Revision der Gattung *Nepiesta* Förster (mit einer Übersicht über die Arten der Gattung *Leptoperillus* Schmiedeknecht) (Hymenoptera, Ichneumonidae). Polskie Pismo Entomologiczne 43: 729-741.
- Horstmann, K. (1974).** Typenrevision von Strobl in der Gattung *Hemiteles* Gravenhorst s.l. beschriebenen Arten und Formen (Hymenoptera, Ichneumonidae). Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen 26: 103-112.
- Horstmann, K. (1976).** Wenig bekannte oder neue europäische Hemitelinen-Gattungen (Hymenoptera, Ichneumonidae, Cryptinae). Nachrichtenblatt der Bayerischen Entomologen 25: 22-31.
- Horstmann, K. (1977).** Bemerkungen zur Systematik einiger Gattungen der Campopleginae. II. (Hymenoptera, Ichneumonidae). Mitteilungen Münchener Entomologischer Gesellschaft 67: 65-83.
- Horstmann, K. (1978).** Revision der Gattungen der Mastrina Townes (Hymenoptera, Ichneumonidae, Hemitelinae). Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen 30: 65-70.
- Horstmann, K. (1981).** Revision der europäischen Tersilochinen II (Hymenoptera: Ichneumonidae). *Spixiana* (suppl.) 4: 1-76.

- Horstmann, K. (1986).** Bemerkungen zur Systematik einiger Gattungen der Campopleginae III (Hymenoptera, Ichneumonidae). Mitteilungen Münchener Entomologischen Gesellschaft 76: 143-164.
- Horstmann, K. (1986).** Revision der von Hartig beschriebenen Ichneumoniden-Arten (Hymenoptera). Spixiana 8: 323-335.
- Horstmann, K. (1990).** Die westpaläarktischen Arten einiger Gattungen der Cryptini (Hymenoptera, Ichneumonidae). Mitteilungen Münchener Entomologischen Gesellschaft 79: 65-89.
- Horstmann, K. (1992).** Revision einiger Gattungen und Arten der Phygadeuontini. Mitteilungen Münchener Entomologischen Gesellschaft 81: 229-254.
- Horstmann, K. (1998).** Revisionen einiger Gattungen und Arten der Phygadeuontini II (Hymenoptera, Ichneumonidae, Cryptinae). Entomofauna 19: 433-460.
- Horstmann, K. (2001).** Revision of the Ichneumonidae (Hymenoptera) described by Johann Christian Fabricius. Beiträge zur Entomologie 51 (1): 7 - 50.
- Horstmann, K. & Shaw, M. R. (1984).** The taxonomy and biology of *Diadegma chrysostictos* (Gmelin) and *Diadegma fabriciana* sp. n. (Hymenoptera: Ichneumonidae). Systematic Entomology 9: 329-337.
- Horstmann, K., Floren, A. and Linsenmair, K.E. (1999).** High species richness of Ichneumonidae (Hymenoptera) from the canopy of a Malaysian rain forest. Ecotropica 5 (1): 1 - 12.
- Horvath, Z. (1987).** Role played by *Oberea erythrocephala* Schrank (Col., Cerambycidae) in restricting the individual number in stands of cypress spurge (*Euphorbia cyparissias* L.). Novevényvedelem 23 (12): 545 - 549.
- Hossain, Z., Gurr, G.M., Wratten, S.D. and Raman, A. (2002).** Habitat manipulation in lucerne *Medicago sativa*: arthropod population dynamics in harvested and 'refuge' crop strips. Journal of Applied Ecology 39 (3): 445 - 454.
- Host Schmidt, O., Asgari, S., Beck, M., Theopold, U., Austin, A.D. and Dowton, M. (2000).** Defence manipulation by parasitoid wasps and the problem of assessing host specificity. Fourth International Hymenoptera Conference, held in Canberra, Australia, in January 1999. Hymenoptera: evolution, biodiversity and biological control 29 - 37.
- Hou, M.L., Wan, F.H. and Wang, F.L. (2002).** Population dynamics of *Helicoverpa assulta* and *Myzus persicae* and their natural enemies in tobacco fields of Shandong Province. Chinese Journal of Biological Control 18 (2):54 - 57.
- Hovemeyer, K. (1992).** Population studies of *Cheilosia fasciata* (Diptera: Syrphidae), a leaf miner of *Allium ursinum*. Ecological Entomology 17 (4): 331 - 337.
- Hovemeyer, K. (1995).** Trophic links, nutrient fluxes, and natural history in the *Allium ursinum* food web, with particular reference to life history traits of two hoverfly herbivores (Diptera: Syrphidae). Oecologia 102 (1): 86 - 94.

- Hu, C., Ye, G.Y., Zhu, J.Q. and Shang, J.N. (1993).** Analysis of tea geometrid (*Ectropis obliqua* Prout) parasite communities and their dynamics. *Journal of Tea Science* 13 (2): 87 - 91.
- Hu, C., Ye, G.Y. and Wang, X.M. (1992).** Investigation on the parasitoids of wild mulberry silkworms. *Acta Phytophylacica Sinica* 19 (4): 337 - 344.
- Hu, C.X., Li, C.D., Wang, Z.Y., Yue, S.K., Liu, K.Y. and Han, S.I. (2000).** Study on the parasitoids of *Pissodes nitidus* Roel. *Journal of Forestry Research* 11 (3): 213 - 214.
- Hu, G.Y., Mitchell, E.R., Sieglaff, D.H. and Okine, J.S. (1998).** Field production of two species of parasitoids of the diamondback moth (Lepidoptera: Plutellidae). *Florida Entomologist* 81 (4): 526 - 534.
- Hu, G.Y., Mitchell, E.R. and Okine, J.S. (1997).** Diamondback moth (Lepidoptera: Plutellidae) on cabbage influence of initial immigration sites on a population distribution, density and larval parasitism. *Journal of Entomological Science* 32 (1): 56 - 71.
- Hu, J.S. and Vinson, S.B. (1998).** The in vitro development from egg to prepupa of *Campoletis sonorensis* (Hymenoptera: Ichneumonidae) in an artificial medium: importance of physical factors. *Journal of Insect Physiology* 44 (5-6): 455 - 461.
- Hu, X.Q. and Wu, S.X. (1987).** Observations on the control effect of parasitic natural enemies on *Cnaphalocrocis medinalis*. *Natural Enemies of Insects* 9 (4): 187 - 189, 198.
- Hua, L. and Shen, B.C. (1992).** A preliminary study on the peach fruitborer (*Carposina niponensis*) infesting wild jujube. *Plant Protection* 18 (2): 15 - 17.
- Huang, Y.Q., Zhang, X.J., Wei, H., Hu, Q.Y. and Zhan, Z.X. (2000).** Studies on the *Dichocrocis punctiferalis* Guenee and its enemies. *Acta Agriculturae Universitatis Jiangxiensis* 22 (4): 523 - 525.
- Huang-Hui', E.N., Fu, X.Y. and Huang, H.E. (2002).** Biological control of sugarcane pests in India. *Sugarcane* 9 (1): 24 - 32.
- Hubbard, S.F., Harvey, I.F. and Fletcher, J.P. (1999).** Avoidance of superparasitism: a matter of learning? *Animal Behaviour* 57 (6): 1193 - 1197.
- Huber, J.T., Eveleigh, E., Pollock, S. and McCarthy, P. (1996).** The chalcidoid parasitoids and hyperparasitoids (Hymenoptera: Chalcidoidea) of *Choristoneura* species (Lepidoptera: Tortricidae) in America north of Mexico. *Canadian Entomologist* 128 (6): 1167 - 1220.
- Hudson, I.R. (1988).** Some further notes on species of Ichneumonidae reared as ectoparasites of spiders. *British Journal of Entomology and Natural History* 1 (2): 77 - 78.
- Hudson, P. and Greenman, J. (1998).** Competition mediated by parasites: biological and theoretical progress. *Trends in Ecology and Evolution*. 13 (10): 387 - 390.
- Hughes, J.P., Harvey, I.F. and Hubbard, S.F. (1994).** Host-searching behavior of *Venturia canescens* (Grav.) (Hymenoptera: Ichneumonidae): superparasitism. *Journal of Insect Behavior* 7 (4): 455 - 464.
- Hughes, J.P., Harvey, I.F. and Hubbard, S.F. (1994).** Host-searching behavior of *Venturia canescens* (Grav.) (Hymenoptera: Ichneumonidae): interference - the effect of mature egg load and prior behavior. *Journal of Insect Behavior* 7 (4): 433 - 454.



- Hull, L.A., Biddinger, D.J. and Felland, C.M. (1992).** Tufted apple bud moth parasitoid diversity, relative effectiveness and temporal distribution in conventional and pheromone disruption apple orchards in Pennsylvania. *Pennsylvania Fruit News* 73 (2): 19 - 23.
- Humala, A.E. (1997).** Oxytorinae from Karelia new to Russia with description of a new genus and two new species (Hymenoptera: Ichneumonidae). *Zoosystematica Rossica* 5: 297-300.
- Humala, A.E. (2002).** A review of the ichneumon wasp genera *Cylloceria* Schiodte, 1838 and *Allomacrus* Forster, 1868 (Hymenoptera: Ichneumonidae) of the Russian fauna. *Entomologicheskoe Obozrenie* 81 (2): 370 - 385.
- Humble, L.M., Shepherd, R.F., Maher, T.F., Alfaro, R.I. and Glover, S.G. (1989).** Biology, outbreak characteristics and damage caused by the black army cutworm (Lepidoptera: Noctuidae). *Insects affecting reforestation: biology and damage* 82 - 88.
- Husberg, G.B. and Hokkanen, H.M.T. (2001).** Effects of *Metarhizium anisopliae* on the pollen beetle *Meligethes aeneus* and its parasitoids *Phradis morionellus* and *Diospilus capito*. *BioControl* 46 (3): 261 - 273.
- Hyden, N. (1990).** *Baptria tibiale* and *Eupithecia actaeata* (Lep.: Geometridae) in NW Europe - distribution, biology, oviposition and preferred habitats. *Entomologisk Tidskrift* 111 (1-2): 1 - 15.

## I

- Idris, A.B. (1999).** Catalogue of Pimplinae (Hymenoptera: Ichneumonidae) from Peninsular Malaysia. *Pan Pacific Entomologist* 75 (2): 73 - 81.
- Idris, A.B. and Grafius, E. (1993).** Differential toxicity of pesticides to *Diadegma insulare* (Hymenoptera: Ichneumonidae) and its host, the diamondback moth (Lepidoptera: Plutellidae). *Journal of Economic Entomology* 86 (2): 529 - 536.
- Idris, A.B. and Grafius, E. (1994).** The potential of using *Barbarea vulgaris* in insecticide-resistant diamondback moth management. *Resistant Pest Management*. 6 (2): 7 - 8.
- Idris, A.B. and Grafius, E. (1995).** Wildflowers as nectar sources for *Diadegma insulare* (Hymenoptera: Ichneumonidae), a parasitoid of diamondback moth (Lepidoptera: Yponomeutidae). *Environmental Entomology* 24 (6): 1726 - 1735.
- Idris, A.B. and Grafius, E. (1996).** Effects of wild and cultivated host plants on oviposition, survival, and development of diamondback moth (Lepidoptera: Plutellidae) and its parasitoid *Diadegma insulare* (Hymenoptera: Ichneumonidae). *Environmental Entomology* 25 (4): 825 - 833.
- Idris, A.B. and Grafius, E. (1997).** Nectar-collecting behavior of *Diadegma insulare* (Hymenoptera: Ichneumonidae), a parasitoid of diamondback moth (Lepidoptera: Plutellidae). *Environmental Entomology* 26 (1): 114 - 120.
- Idris, A.B. and Grafius, E. (1998).** Diurnal flight activity of *Diadegma insulare* (Hymenoptera: Ichneumonidae), a parasitoid of the diamondback moth (Lepidoptera: Plutellidae), in the field. *Environmental Entomology* 27 (2): 406 - 414.

- Idris, A.B. and Grafius, E. (2001).** Effects of plant density on abundance of diamondback moth (Lepidoptera: Plutellidae) and *Diadegma insulare* (Cresson) (Hymenoptera: Ichneumonidae). International Journal of Pest Management 47 (2): 103 - 107.
- Idris, A.B. and Kee, S.S. (2002).** Horizontal and vertical diversity of ichneumonid wasps (Hymenoptera: Ichneumonidae) in the Sungkai Wildlife Forest Reserve in Perak, Malaysia. Journal of Asia Pacific Entomology 5 (1): 85 - 89.
- Iga, M. (1992).** Reproduction of an introduced ichneumonid parasitoid, *Diadegma semiclausum* on larvae of *Plutella xylostella* L. Proceedings of the Kanto Tosan Plant Protection Society 39: 201 - 202.
- Iga, M. (1995).** Copulation, sex ratio and reproductive rate on mass rearing of *Diadegma semiclausum*, an introduced ichneumonid, on larvae of the diamondback moth, *Plutella xylostella*. Proceedings of the Kanto Tosan Plant Protection Society 42: 209 - 211.
- Iga, M. (1996).** Possibility of establishing an introduced ichneumonid parasitoid, *Diadegma semiclausum* [*D. semiclausum*] and how to use the parasitoid on larvae of the diamondback moth, *Plutella xylostella*. Proceedings of the Kanto Tosan Plant Protection Society 43: 207 - 208.
- Igbinosa, I.B. (1992).** Field and laboratory techniques for assessing infestations of the nettle caterpillar, *Latoia viridissima* Holland (Lepidoptera: Limacodidae). Insect Science and its Application 13 (3): 389 - 398.
- Ionaitis, V. (1991).** Records and descriptions of insect species new and rare for Lithuania, 1991. Institute of Ecology, Lithuanian Entomological Society; Vilnius; Lithuania, 70 pp.
- Iizuka, T. and Takasu, K. (1998).** Olfactory associative learning of the pupal parasitoid *Pimpla luctuosa* Smith (Hymenoptera: Ichneumonidae). Journal of Insect Behavior 11 (5): 743 - 760.
- Iizuka, T. and Takasu, K. (1999).** Balancing between host- and food-foraging by the host-feeding pupal parasitoid *Pimpla luctuosa* Smith (Hymenoptera: Ichneumonidae). Entomological Science 2 (1): 67 - 73.
- Ikincisoy, Y., Kornosor, S. and Sertkaya, E. (1994).** Investigation of the natural enemies of *Acantholeucania loreyi* Duponchel (Lepidoptera: Noctuidae) larvae in Cukurova region. Turkiye III. Biyolojik Mucadele Kongresi Bildirileeri, 25-28 Ocak 1994, Ege Universitesi Ziraat Fakultesi, Bitki Koruma Bolumu, Izmir 555 - 564.
- Ionaitis, V. and Jonaitis, V. (1992).** Structure and distribution of trophic relations of insect communities in the main agricultural crops. Acta Entomologica Lituanica (10): 3 - 16.
- Ionaitis, V.P. (1990).** Resources, formation and functioning of host-parasite insect complexes in ecosystems. Mokslas; Vilnius; Lithuania 231 pp.
- Ionaitis, V.P. (1991).** 110 species of ichneumonids (Hymenoptera: Ichneumonidae) new for the Lithuanian fauna found in 1990. Novye i Redkie Dlya Litovskoi SSR Vidy Nasekomykh, Soobshcheniya i opisaniya 1991: 30 - 41.
- Iqbal, M., Ismail, F. and Wright, D.J. (1996).** Loss of residual activity of abamectin on foliage against adult hymenopteran parasitoids. Entomophaga 41 (1): 117 - 124.

- Iqbal, M. and Wright, D.J. (1996).** Host resistance to insecticides can confer protection to endo-larval parasitoids. *Bulletin of Entomological Research* 86 (6): 721 - 723.
- Isenhour, D.J. and Wiseman, B.R. (1989).** Parasitism of the fall armyworm (Lepidoptera: Noctuidae) by *Campoletis sonorensis* (Hymenoptera: Ichneumonidae) as affected by host feeding on silks of *Zea mays* L. cv. Zapalote Chico. *Environmental Entomology* 18 (3): 394 - 397.
- Ishikawa, H. (1989).** Biochemical and molecular aspects of endosymbiosis in insects. *International Review of Cytology* 116: 1 - 45.
- Ishitsuka, T., Watanabe, Y. and Ohsawa, S. (1999).** Parasitoids of the diamondback moth *Plutella xylostella* (L.) collected from broccoli fields in Fukushima Prefecture [Japan]. *Annual Report of the Society of Plant Protection of North Japan* 50: 190 - 192.
- Izquierdo, J.I., Solans, P. and Vitalle, J. (1994).** Parasitoids and predators of *Helicoverpa armigera* (Hubner) on table tomato crops. *Boletín de Sanidad Vegetal, Plagas* 20 (2): 521 - 530.

## J

- Jackson, D.M. and Sisson, V.A. (1998).** Potential of *Nicotiana kawakamii* (Solanaceae) as a trap crop for protecting flue-cured tobacco from damage by *Heliothis virescens* (Lepidoptera: Noctuidae) larvae. *Journal of Economic Entomology* 91 (3): 759 - 766.
- Jacob, H.S. and Evans, E.W. (1998).** Effects of sugar spray and aphid honeydew on field populations of the parasitoid *Bathyplectes curculionis* (Hymenoptera: Ichneumonidae). *Environmental Entomology* 27 (6): 1563 - 1568.
- Jacob, H.S. and Evans, E.W. (2000).** Influence of experience on the response of *Bathyplectes curculionis* (Hymenoptera: Ichneumonidae), a nonaphidophagous parasitoid, to aphid odor. *Biological Control* 19 (3): 237 - 244.
- Jacob, H.S. and Evans, E.W. (2001).** Influence of food deprivation on foraging decisions of the parasitoid *Bathyplectes curculionis* (Hymenoptera: Ichneumonidae). *Annals of the Entomological Society of America* 94 (4): 605 - 611.
- Jager, C.R. and Verduijn, M. (1995).** Genetic differences between populations of the endoparasitoid *Herpestomus brunnicornis* related to their *Yponomeuta* host species. *Proceedings of the- Section Experimental and Applied Entomology of the Netherlands Entomological Society* 6: 147 - 152.
- Jahromi, K.T., Omid, M. and Sabahi, G. (2002).** A study of the three methods for alfalfa weevil control with special reference to flaming of plant stubble and debris. *Iranian Journal of Agricultural Sciences* 33 (2): 325 - 332.
- Jaiswal, A.K., Sharma, K.K., Sushil, S.N., Bhattacharya, A. and Mishra, Y.D. (1998).** Lac associated insect fauna during storage. *Shashpa* 5 (2): 133 - 136.
- Jaiswal, A.K., Sharma, K.K. and Agarwal, S.C. (1999).** A modified and upgraded device of insect-separation for managing the insect pests of lac. *National Academy Science Letters* 22 (5-6): 106 - 110.

- Jalali, S.K., Singh, S.P., Kumar, P. and Ballal, C.R. (1989).** New record of hyperparasitoids on *Campoletis chloridae* Uchida and *Eucelatoria bryani* Sabroskwy parasitizing *Heliothis armigera* (Hubner) on tomato. *Current Science* 58 (6): 326 - 327.
- Jalali, S.K. and Singh, S.P. (2001).** Life table studies on the natural enemies of *Chilo partellus* (Swinhoe) (Lepidoptera: Pyralidae). *Journal of Biological Control* 15 (2): 113 - 117.
- Jalali, S.K. and Singh, S.P. (2002).** Seasonal activity of stem borers and their natural enemies on fodder maize. *Entomon* 27 (2): 137 - 146.
- Jalaluddin, M. and Mohanasundaram, M. (1989).** Residual toxicity of four insecticides recommended for control of coconut coccids on the parasitoid fauna of *Opisina arenosella* Wlk. *Entomon* 14 (3-4): 199 - 202.
- Jamieson, W. and Resh, V.H. (1998).** Biology of *Tanychela pilosa* (Hymenoptera: Ichneumonidae), a parasitoid of the aquatic moth *Petrophila confusalis* (Lepidoptera: Pyralidae). *Entomological News* 109 (5): 329 - 338.
- Jansson, R.K., Leibe, G.L., Sanchez, C.A. and Lecrone, S.H. (1991).** Effects of nitrogen and foliar biomass on population parameters of cabbage insects. *Entomologia Experimentalis et Applicata* 61 (1): 7 - 16.
- Janzen, D.H., Sharkey, M.J. and Burns, J.M. (1998).** Parasitization biology of a new species of Braconidae (Hymenoptera) feeding on larvae of Costa Rican dry forest skippers (Lepidoptera: Hesperidae: Pyrginae). *Tropical Lepidoptera* 9: Supplement 2, 33 - 41.
- Jaworska, M. (1987).** Observations on *Lathrolestes marginatus* (Thompson), a parasitoid of the apple sawfly - *Hoplocampa testudinea* (Klug.) (Hymenoptera: Tenthredinidae). *Polskie Pismo Entomologiczne* 57 (3): 553 - 567.
- Jaworska, M. (1992).** Biological control of *Hoplocampa testudinea* Klug. (Hymenoptera: Tenthredinidae). *Acta Phytopathologica et Entomologica Hungarica* 27 (1-4): 311 - 315.
- Jaworska, M., Rópek, D. and Pruszyński, S. (1995).** The influence of entomopathogenic nematodes on insect natural enemies. Proceedings of the XXXV Scientific Meeting of the Institute of Plant Protection. Part II - posters. *Materialy Sesji Instytutu Ochrony Roslin* 35 (2): 434 - 436.
- Jervis, M.A. (1990).** Predation of *Lissonota coracinus* (Gmelin) (Hymenoptera: Ichneumonidae) by *Dolichonabis limbatus* (Dahlbom) (Hemiptera: Nabidae). *Entomologist's Gazette* 41 (4): 231 - 233.
- Jewett, D.K. and Carpenter, J.E. (1998).** Chemical communication between males and females of a pupal parasitoid *Diapetimorpha introita* (Hymenoptera: Ichneumonidae). *Annals of the Entomological Society of America* 91 (5): 748 - 753.
- Jewett, D.K. and Carpenter, J.E. (1999).** Chemically-mediated attraction of *Ichneumon* (= *Pterocormus*) *promissorius* (Hymenoptera: Ichneumonidae) males by females. *Environmental Entomology* 28 (4): 551 - 556.

- Jewett, D.K. and Carpenter, J.E. (2001).** Seasonal abundance of a pupal parasitoid, *Diapetimorpha introita* (Hymenoptera: Ichneumonidae). Armyworm Symposium 2000. Florida Entomologist 84 (1): 50 - 54.
- Jimenez, J. and Fernandez, R. (1985).** Behaviour of *Diadegma* sp. parasitism of *Heliothis virescens* caterpillars in field plots with different microbiological and chemical insecticide treatments. Ciencia-y Tecnica en la Agricultura, Proteccion de Plantas 8 (3): 67 - 79.
- Jimenez, M.L. (1987).** Biological relations between spiders and wasps. Folia Entomologica Mexicana, publ. 1988, 73: 173 - 183.
- Jimenez, R., Falco, J.V., Gimeno, C., Luna, F., Moreno, J., Selfa, J. and Serrano, C. (1992).** Biological control of *Chilo suppressalis* Walker (Lep.: Pyralidae) I. Study of the parasitoid complex in rice fields in Valencia. Boletin de Sanidad Vegetal, Plagas 18 (1): 201 - 212.
- Jin, L.Y., Alfaro, R.I. and Glover, S.G. (1989).** *Pissodes nitidus* Roelofs, the yellow-spotted pine weevil (Coleoptera: Curculionidae): a serious pest of Korean pine plantations in northeast China. Insects affecting reforestation: biology and damage 186 - 193.
- Johanowicz, D.L. and Mitchell, E.R. (2000).** A novel method to rear *Diadegma insulare* (Hymenoptera: Ichneumonidae), a parasitoid of the diamondback moth (Lepidoptera: Plutellidae). Florida Entomologist 83 (3): 377 - 379.
- Johanowicz, D.L. and Mitchell, E.R. (2000).** Effects of sweet alyssum flowers on the longevity of the parasitoid wasps *Cotesia marginiventris* (Hymenoptera: Braconidae) and *Diadegma insulare* (Hymenoptera: Ichneumonidae). Florida Entomologist 83 (1): 41 - 47.
- Johnson, J.A., Valero, K.A., Hannel, M.M. and Gill, R.F. (2000).** Seasonal occurrence of postharvest dried fruit insects and their parasitoids in a culled fig warehouse. Journal of Economic Entomology 93 (4): 1380 - 1390.
- Johnson, M.T. (1997).** Interaction of resistant plants and wasp parasitoids of tobacco budworm (Lepidoptera: Noctuidae). Environmental Entomology 26 (2): 207 - 214.
- Johnson, M.T., Gould, F. and Kennedy, G.G. (1997).** Effects of natural enemies on relative fitness of *Heliothis virescens* genotypes adapted and not adapted to resistant host plants. Entomologia Experimentalis et Applicata 82 (2): 219 - 230.
- Johnson, M.T. and Gould, F. (1992).** Interaction of genetically engineered host plant resistance and natural enemies of *Heliothis virescens* (Lepidoptera: Noctuidae) in tobacco. Environmental Entomology 21 (3): 586 - 597.
- Johnson, M.W., Mitchell, W.C., Robin, M.R., Cushing, N.L. and Rethwisch, M.D. (1988).** Parasitization of the diamondback moth, *Plutella xylostella* (L.) (Lepidoptera: Plutellidae) in Hawaii. Proceedings of the Hawaiian Entomological Society 28: 197 - 203.
- Jonaitis, V. (1991).** Insect species new and rare for Lithuania. Records and descriptions of 1991. Novye i Redkie Dlya Litovskoi SSR Vidy Nasekomykh, Soobshcheniya I opisaniya 1991: 70 pp.
- Jonaitis, V. (1992).** Records and descriptions of insect species new and rare for Lithuania 1992. Institute of Ecology, Lithuanian Entomological Society; Vilnius; Lithuania, 115 pp.

- Jonaitis, V. (1993).** New and rare for Lithuania insect species. Records and descriptions for 1993. Records and Descriptions of Insect Species New and Rare for Lithuania, 65 pp.
- Jonaitis, V. (1997).** Some aspects of forest ecosystems' relations. Miskinkyste, publ. 1998, 1: 39, 47 - 57.
- Jonaitis, V. and Jonaitis, V. (1996).** Some patterns of long-term dynamics of population density within a territory of various formations of ermine moths and their parasitoids. Lietuvos entomologu darbai Lietuvos entomologu draugijos 30 meciui 103 - 134.
- Jonaitis, V. and Rimsaite, J. (2002).** Fauna of Orthocentrinae (S.S.) (Hymenoptera: Ichneumonidae) in Lithuania. Acta Zoologica Lituanica 12 (2): 167 - 179.
- Jonathan, J.K. (1971).** A new Mesostenine genus *Rupa* from India. Oriental Insects 5: 149-152.
- Jonathan, J.K. (1980).** The *Isotima*-Complex (Hymenoptera: Ichneumonidae). Records of the Zoological Survey of India. Miscellaneous Publications. Occasional Paper 17: 1-146.
- Jonathan, J.K. (1982).** A new Mesostenine genus *Anupama* (Hymenoptera: Ichneumonidae) from India. Oriental Insects 15: 281-285.
- Jonathan, J.K. (1987).** A new Mesostenine genus *Pseudotricapus* from India (Hymenoptera: Ichneumonidae). Bulletin of the Zoological Survey of India 8: 57-60.
- Jonathan, J.K. and Gupta, V.K. (1973).** Ichneumonologia Orientalis, or the Ichneumonidae of the Oriental Region. Part III. The *Goryphus*-Complex (Hymenoptera: Ichneumonidae). Oriental Insects Monographs 3: 203.
- A new Mesostenine genus *Rupa* from India. Oriental Insects 5: 149-152.
- Jonathan, J.K. (1980).** The *Isotima*-Complex (Hymenoptera: Ichneumonidae). Records of the Zoological Survey of India. Miscellaneous Publications. Occasional Paper 17: 1-146.
- Jonathan, J.K. (1982).** A new Mesostenine genus *Anupama* (Hymenoptera: Ichneumonidae) from India. Oriental Insects 15: 281-285.
- Jonathan, J.K. (1987).** A new Mesostenine genus *Pseudotricapus* from India (Hymenoptera: Ichneumonidae). Bulletin of the Zoological Survey of India 8: 57-60.
- Jonathan, J.K. and Gupta, V.K. (1973).** Ichneumonologia Orientalis, or the Ichneumonidae of the Oriental Region. Part III. The *Goryphus*-Complex (Hymenoptera: Ichneumonidae). Oriental Insects Monographs 3: 203.
- A new Mesostenine genus *Rupa* from India. Oriental Insects 5: 149-152.
- Jonathan, J.K. 1980. The *Isotima*-Complex (Hymenoptera: Ichneumonidae). Records of the Zoological Survey of India. Miscellaneous Publications. Occasional Paper 17: 1-146.
- Jonathan, J.K. (1982).** A new Mesostenine genus *Anupama* (Hymenoptera: Ichneumonidae) from India. Oriental Insects 15: 281-285.
- Jonathan, J.K. (1987).** A new Mesostenine genus *Pseudotricapus* from India (Hymenoptera: Ichneumonidae). Bulletin of the Zoological Survey of India 8: 57-60.
- Jonathan, J.K. and Gupta, V.K. (1973).** Ichneumonologia Orientalis, or the Ichneumonidae of the Oriental Region. Part III. The *Goryphus*-Complex (Hymenoptera: Ichneumonidae). Oriental Insects Monographs 3: 203.

- Jordan, T. (1998).** Egg structure and life history of two species groups of *Grypocentrus* (Hymenoptera: Ichneumonidae), monophagous parasitoids of spring leaf-mining Eriocrania-moths on birch (Lepidoptera: Eriocraniidae). *Entomologia Generalis* 22 (3-4): 223 - 231.
- Jordan, T. (1995).** Life history and parasitoid community of the oak-leaf-mining moth *Tischeria ekebladella* (Bjerkander, 1795) (Lep.: Tischeriidae) in Northern Germany. *Journal of Applied Entomology* 119 (7): 447 - 454.
- Joshi, K.C. and Agarwal, S.B. (1987).** Bionomics of the tent caterpillar, *Malacosoma indica* Wlk. *Bulletin of Entomology New Delhi* 28 (1): 1 - 11.
- Jungen, C. and Madel, G. (1993).** Humoral immunity in *Plutella xylostella* L. (Lep.: Yponomeutidae) - induced lysozyme activity. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 8 (4-6): 893 - 897.
- Jussila, R. (1979).** A revision of the genus *Atractodes* (Hymenoptera: Ichneumonidae) in the western Palearctic region. *Acta Entomologica Fennica* 34: 1-44.
- Jussila, R. (1987).** Revision of the genus *Stilpnus* (Hymenoptera: Ichneumonidae) of the western Palearctic region. *Annales Entomologici Fennici* 53: 1-16.
- Jussila, R. (1998).** The genera *Lissaspis* Townes and *Singalissaspis* gen. n. (Hymenoptera, Ichneumonidae). *Entomologica Fennica* 9: 103-114.
- Jussila, R. (1991).** Ichneumonidae new to Finland (Hymenoptera). I. *Entomologica Fennica* 2 (1): 37 - 42.
- Jussila, R. (1994).** Additions to the revision of the genus *Atractodes* (Hymenoptera: Ichneumonidae) of the western Palaearctic Region. II. *Entomologica Fennica* 5 (3): 129 - 134.
- Jussila, R. (1999).** Additions to the revisions of the genus *Stilpnus* (Hymenoptera: Ichneumonidae) of the Palaearctic region. I. *Entomologica Fennica* 10 (2): 107 - 112.
- Jussila, R. (1999).** Ichneumonidae (Hymenoptera) new to Finland. V. *Entomologica Fennica* 10 (3): 175 - 177.
- Jussila, R. (2000).** *Dyspetes praerogator* (Thomson, 1883), a new species and genus to Finland and Denmark (Hym.: Ichneumonidae: Tryphoninae). *Entomologica Fennica* 11 (2): 88.
- Jussila, R. (2001).** Additions to the revision of the genus *Atractodes* (Hymenoptera: Ichneumonidae) of the Palaearctic Region. III. *Entomologica Fennica* 12 (4): 193 - 216.

## K

- Kabre, G.B. and Petter, A.J. (1997).** *Camallanus polypteri* n. sp. (Nematoda: Camallanidae) in freshwater fishes from Burkina Faso. *Onderstepoort Journal of Veterinary Research* 64 (1): 33 - 37.
- Kaczmarzyk, M., Klukowski, Z. and Kelm, M. (2001).** Occurrence of pollen beetle (*Meligethes* spp.) and its parasitoids Tersilochinae (Ichneumonidae) on winter rape – spatio-temporal analysis. *Progress in Plant Protection* 41 (2): 454 - 458.

- Kadamshoev, M. (1996).** Cutworm moths as pests of *Calendula*. *Zashchita Rastanii Moskva* 1: 32 - 33.
- Kadlubowski, W. and Piekarska, H. (1984).** Contribution to the knowledge of the fauna of Ichneumonoidea (Hymenoptera: Parasitica), occurring in apple orchards in the Poznan region. *Roczniki Nauk Rolniczych, E Ochrona Roslin* 14 (1-2): 47 - 71.
- Kadlubowski, W. and Szmyt, J. (1985).** Data on the bionomics of the pear ermel moth *Swammerdamia pyrella* (De Villers, 1789) (Lep.: Yponomeutidae). *Roczniki Nauk Rolniczych, E Ochrona Roslin* 15 (1-2): 119 - 128.
- Kaitaniemi, P. and Ruohomaki, K. (1999).** Effects of autumn temperature and oviposition date on timing of larval development and risk of parasitism in a spring folivore. *Oikos* 84 (3): 435 - 442.
- Kakar, K.L. and Dogra, G.S. (1989).** Some natural enemies of lepidopterous pests in Himachal Pradesh. *Agricultural Science Digest Karnal* 9 (2): 74 - 76.
- Kakar, K.L. and Sharma, J.P. (1991).** Role of natural enemies on the mortality of cabbage butterfly, *Pieris brassicae* (L.) in cauliflower seed crop. *Indian Journal of Plant Protection* 19 (2): 210 - 212.
- Kalmes, R. and Rojas-Rousse, D. (2001).** Frequency and viability of diploid and haploid male offspring of mated females of solitary endoparasitoid *Diadromus pulchellus*. *Comptes Rendus de l'Academie des Sciences. Serie III, Sciences de la- Vie.* 324 (2): 149 - 158.
- Kalra, V.K. (1989).** Natural parasitism of sesame leaf webber and pod borer, *Antigastra catalaunalis* Duponchel. *Indian Journal of Plant Protection* 17 (1): 9 - 11.
- Kamijo, K. (1996).** Notes on the genus *Cleonymus* Latreille (Hymenoptera: Pteromalidae), with descriptions of four new species from Japan. *Japanese Journal of Entomology* 64 (4): 745 - 758.
- Kamin-Belsky, N., Wool, D., Brower, J.H., Belsky, N. Kamin., Donahaye, E. and Navarro, S. (1986).** Unplanned factors affecting population size in laboratory populations of the almond moth *Ephesia cautella* (Walker). *Proceedings of the Fourth International Working Conference on Stored Product Protection, Tel Aviv, Israel, 21 - 26 September,* 463 - 469.
- Kamm, J.A. (1990).** Biological observations of glassy cutworm (Lepidoptera: Noctuidae) in western Oregon. *Pan Pacific Entomologist* 66 (1): 66 - 70.
- Kanecka, P. (1995).** Parasitoids of false spruce webworm (*Cephalcia abietis* (L.)) - findings obtained during larvae dissection. *Lesnictvi Forestry* 41 (12): 545 - 554.
- Kaneko, J. (1993).** Parasitic wasps of the silver Y moth, *Autographa gamma* (L.) (Noctuidae: Plusiinae) and the Asiatic common looper, *A. nigrisigna* (Walker) in Hokkaido, Japan. *Japanese Journal of Applied Entomology and Zoology* 37 (1): 22 - 24.
- Kaneko, J.I. (1995).** Rate of parasitism of cocoons silver Y moth, *Autographia gamma* (L.) in grass fields at Sapporo, Japan and species composition of parasitoids. *Japanese Journal of Applied Entomology and Zoology* 39 (2): 162 - 164.



- Kanhekar, L.J. and Nikam, P.K. (1989).** A new species of genus *Mesostenus* Gravenhorst (Hymenoptera: Ichneumonidae) from India. *Entomon* 14 (1-2): 15 - 18.
- Kansu, I.A. and Uguk, A. (1990).** Influence of food on fecundity and longevity of adults of *Pimpla turionellae* (L.) (Hymenoptera: Ichneumonidae). *Proceedings of the Second Turkish National Congress of Biological Control* 61 - 65.
- Kapadia, M.N. and Puri, S.N. (1991).** Parasitization of *Chrysoperla carnea* (Stephens) in cotton field under Parbhani conditions. *Journal of Maharashtra Agricultural Universities* 16 (3): 453 - 454.
- Kares, E.A. and Zarif, G. (1992).** Effect of cutting and weather factors on the population of *Hypera brunneipennis* Boh. and its parasitoid *Bathyplectes curculionis* Thoms. in alfalfa at Nubaria. *Annals of Agricultural Science, Moshtohor* 30 (2): 1117 - 1128.
- Kartosuwondo, U. and Sunjaya. (1991).** Potential role of wild crucifers in the preservation of *Diadegma eucerophaga* Horstman (Hymenoptera: Ichneumonidae), a parasitoid of the diamondback moth, *Plutella xylostella* Linnaeus (Lepidoptera: Plutellidae). *Biotropia* 4: 31 - 40.
- Kasparyan, D.R. (1969).** Palearctic species of the genus *Tryphon* Fallén (Hymenoptera, Ichneumonidae), I. (in Russian) *Éntomologicheskoe Obozrenie* 48: 639-662.
- Kasparyan, D.R. (1970).** A new genus of ichneumon flies, *Aridella* gen. n. (Hymenoptera, Ichneumonidae), from Middle Asia. (in Russian) *Zoologicheskii Zhurnal* 49: 154-157.
- Kasparyan, D.R. (1971).** Genus *Aderaeon* Townes (Hymenoptera, Ichneumonidae) new for Palearctic. (in Russian) *Zoologicheskii Zhurnal* 50: 1586-1589.
- Kasparyan, D.R. (1976).** Review of the Ichneumonids of the tribe Polysphinctini and Poemeniini (Hymenoptera, Ichneumonidae) of the Far East. (in Russian). *Trudy Zoologicheskogo Instituta* 67: 68-89.
- Kasparyan, D.R. (1976).** New ichneumonids of the tribe Cteniscini (Hymenoptera, Ichneumonidae) from the USSR and the Mongolian People's Republic. *Schelocentrus* gen. n., *Smicroplectrus* Thomson. (in Russian) *Nasekomye Mongol.* 4: 282-299.
- Kasparyan, D.R. (1977).** A revision of the genus *Eclytus* Holmgren (Hymenoptera, Ichneumonidae). (in Russian). *Éntomologicheskoe Obozrenie* 56: 156-170.
- Kasparyan, D.R. (1981).** Ichneumonidae (Subfamily Tryphoninae), tribe Tryphonini. *In: Strelkov, A.A. (ed.) Fauna of the USSR: Hymenoptera* 3(1). New Delhi: Amerind Publishing Co. 414 pp.
- Kasparyan, D.R. (1983).** *Ledora* gen. n. (Hymenoptera, Ichneumonidae) from the Soviet Far East. (in Russian) *Zoologicheskii Zhurnal* 62: 1749-1751.
- Kasparyan, D.R. (1985).** A new genus of the tribe Exenterini (Hymenoptera: Ichneumonidae) from Middle Asia. (in Russian) *Zoologicheskii Zhurnal* 64: 628-630.
- Kasparyan, D.R. (1986).** Contribution to the revision of ichneumonids of the genus *Adelognathus* Holmgren (Hymenoptera: Ichneumonidae). *Trudy Zoologicheskogo Instituta, Akademiya Nauk SSSR* 159: 38 - 56.

- Kasparyan, D.R. (1987).** Characteristics of the parasitoid fauna of sawflies (Hymenoptera: Symphyta). *Entomologicheskoe Obozrenie* 66 (3): 521 - 528.
- Kasparyan, D.R. (1988).** New taxa of ichneumonids of the family Paxylommatidae (Hymenoptera, Ichneumonoidea) from the Baltic Amber. (in Russian) *Trudy Vesesoyuznogo éntomologicheskogo obshchestva* 70: 125-131.
- Kasparyan, D.R. (1988).** A new subfamily and two new genera of ichneumonids (Hymenoptera, Ichneumonidae) from Baltic Amber (in Russian). *Trudy Zoologicheskogo instituta, Akademiya nauk SSSR* 175: 38-43.
- Kasparyan, D.R. (1989).** Fauna of the USSR. Hymenoptera. Volume III, Number 1. Ichneumonidae (subfamily Tryphoninae). E.J. Brill; Leiden; Netherlands, 414 pp, 22 pp.
- Kasparyan, D.R. (1990).** Fauna of the USSR. Insecta Hymenoptera. Vol. III, No. 2. Parasitic ichneumonids (Ichneumonidae) subfamily Tryphoninae: tribe Exenterini, subfamily Adelognathinae. "Nauka", Leningradskoe Otdelenie; Leningrad; USSR, 340 pp.
- Kasparyan, D.R. (1993).** Review of Palaearctic species of ichneumonid wasps of the genus *Phytodietus* Grav. (Hymenoptera: Ichneumonidae). *Entomologicheskoe Obozrenie* 72 (4): 869 - 890.
- Kasparyan, D.R. (1993).** Townesioninae, a new ichneumonid subfamily from the eastern Palearctic (Hymenoptera: Ichneumonidae). *Zoosystematica rossica* 1: 86-88.
- Kasparyan, D.R. (1993).** *Zambion*, a new Tryphonine genus from Africa (Hymenoptera: Ichneumonidae). *Zoosystematica Rossica* 1: 86-88.
- Kasparyan, D.R. (1993).** Townesioninae, a new ichneumonid subfamily from the Eastern Palearctic (Hymenoptera: Ichneumonidae). *Zoosystematica Rossica* 2 (1): 155 - 159.
- Kasparyan, D.R. (1994).** Review of Ichneumon flies of the Townesitinae subfam. nov. (Hymenoptera, Ichneumonidae) from the Baltic ambers. (in Russian) *Paleontologicheskii Zhurnal* 4: 86-96.
- Kasparyan, D.R. (1995).** A new genus and two new species of ichneumonid wasps of the subfamily Neorhacodinae (Hymenoptera, Ichneumonidae) from deserts of Turkarmenia. (in Russian) *Entomologicheskoye Obozreniye* 74: 669-672.
- Kasparyan, D.R. (1998).** Taxonomic notes on the species of *Mesoleius* s. l., *Hyperbatus* and *Phaestus* in the museums of Stockholm, Lund and Munich (Hymenoptera: Ichneumonidae, Ctenopelmatinae). *Zoosystematica Rossica* 7 (10): 181 - 183.
- Kasparyan, D.R. (1999).** New species of ichneumonid wasps of the subfamily Stilbopinae (Hymenoptera: Ichneumonidae) of the Old World. *Entomologicheskoe Obozrenie* 78 (1): 163 - 173.
- Kasparyan, D.R. (2000).** Palaearctic ichneumon wasps of the genus *Mesoleius* Holmgren (s. str.) (Hymenoptera: Ichneumonidae). I. *Entomologicheskoe Obozrenie* 79 (1): 150 - 179.
- Kasparyan, D.R. (2001).** Palaearctic ichneumonid wasps of the genus *Mesoleius* Holmgren (s. str.) (Hymenoptera: Ichneumonidae). II. *Entomologicheskoe Obozrenie* 80 (3): 706 - 733.

- Kasparyan, D.R. (2002).** A new species of *Megarhyssa* Ashmead, 1900 (Hymenoptera: Ichneumonidae) from Mexico. Russian Entomological Journal 11 (3): 305 - 306.
- Kasparyan, D.R. and Manukyan, A.R. (1987).** A new genus of parasitic wasps (Hymenoptera: Ichneumonidae: Diplazontinae) from the eastern Palearctic region. (in Russian) Éntomologicheskoe Obozrenie 66: 841-844.
- Kasparyan, D.R. and Gultekin, L. (2002).** First records of two ichneumonid wasps from Northeastern Turkey (Hymenoptera: Ichneumonidae). Zoosystematica Rossica 11 (1): 218.
- Kasparyan, D.R. and Humala, A.E. (1995).** A new Palearctic species of the genus *Cushmania* Dasch, 1992 (Hymenoptera: Ichneumonidae: Microleptinae). Zoosystematica Rossica. publ. 1996, 4 (1): 163 - 165.
- Kasparyan, D.R. and Ruiz-Cancino, E. (2001).** Review of the Mexican species of the genus *Corsoncus* Townes, 1971 (Hymenoptera: Ichneumonidae: Anomaloninae). Russian Entomological Journal 10 (2): 159 - 162.
- Kasparyan, D.R. and Shaw, M.R. (2003).** A preliminary key to European species of the genus *Saotis* Forster, 1869, with a list of British species (Ichneumonidae: Ctenopelmatinae: Mesoleiini). Zoosystematica Rossica 11(2): 351-355.
- Katiyar, K.M. and Sharma, V. (1987).** Seasonal mortality of *Prolatus aciculata* Sharma, a larval parasitoid of *Sinoxylon anale*, due to hyperparasitoids. Transactions of Indian Society of Desert Technology 12 (1): 27 - 28.
- Kauffman, W.C. and Kennedy, G.G. (1989).** Inhibition of *Campoletis sonorensis* parasitism of *Heliothis zea* and of parasitoid development by 2-tridecanone-mediated insect resistance of wild tomato. Journal of Chemical Ecology 15 (6): 1919 - 1930.
- Kauffman, W.C. and Kennedy, G.G. (1989).** Toxicity of allelochemicals from wild insect-resistant tomato *Lycopersicon hirsutum* f. *glabratum* to *Campoletis sonorensis*, a parasitoid of *Heliothis zea*. Journal of Chemical Ecology 15 (7): 2051 - 2060.
- Kaur, R. (1988).** A revision of the genus *Litochila* (Hymenoptera: Ichneumonidae). Oriental Insects 22: 359 - 376.
- Kaur, R. and J.K. Jonathan. (1979).** Ichneumonologia Orientalis. Part VIII. The tribe Phytodietini from India (Hymenoptera: Ichneumonidae). Oriental Insects Monographs 9: 1-276.
- Kenis, M. and Mills, N.J. (1994).** Parasitoids of European species of the genus *Pissodes* (Col.: Curculionidae) and their potential for the biological control of *Pissodes strobi* (Peck) in Canada. Biological Control 4 (1): 14 -21.
- Kerslake, J.E., Kruuk, L.E.B., Hartley, S.E. and Woodin, S.J. (1996).** Winter moth (*Operophtera brumata* (Lepidoptera: Geometridae)) outbreaks on Scottish heather moorlands: effects of host plant and parasitoids on larval survival and development. Bulletin of Entomological Research 86 (2): 155 - 164.
- Kester, K. and Tuttle, J. (1999).** Host identified for *Rhynchophion flammipennis* (Hymenoptera: Ichneumonidae). Southwestern Naturalist 44 (1): 87 - 88.
- Kfir, A.R. (1991).** Non-chemical control of grain stalk borers. Plant Protection News 23: 4.

- Kfir, R. (1988).** Hibernation by the lepidopteran stalk borers, *Busseola fusca* and *Chilo partellus* on grain sorghum. *Entomologia Experimentalis et Applicata* 48 (1): 31 - 36.
- Kfir, R. (1990).** Parasites of the spotted stalk borer, *Chilo partellus* (Lepidoptera: Pyralidae) in South Africa. *Entomophaga* 35 (3): 403 - 410.
- Kfir, R. (1991).** Selecting parasites for biological control of lepidopterous stalk borers in summer grain crops in South Africa. *Redia* 74 (3): Appendix, 231 - 236.
- Kfir, R. (1992).** Seasonal abundance of the stem borer, *Chilo partellus* (Lepidoptera: Pyralidae) and its parasites on summer grain crops. *Journal of Economic Entomology* 85 (2): 518 - 529.
- Kfir, R. (1994).** Attempts at biological control of the stem borer *Chilo partellus* (Swinhoe) (Lepidoptera: Pyralidae) in South Africa. *African Entomology* 2 (1): 67 - 68.
- Kfir, R. (1995).** Parasitoids of the African stem borer, *Busseola fusca* (Lepidoptera: Noctuidae), in South Africa. *Bulletin of Entomological Research* 85 (3): 369 - 377.
- Kfir, R. (1997).** Natural control of the cereal stemborers *Busseola fusca* and *Chilo partellus* in South Africa. *Insect Science and its Application* 17 (1): 61 - 67.
- Kfir, R. (1998).** Origin of the diamondback moth (Lepidoptera: Plutellidae). *Annals of the Entomological Society of America* 91 (2): 164 - 167.
- Kfir, R. (2001).** Prospects for biological control of *Chilo partellus* in grain crops in South Africa. *Insect Science and its Application* 21 (4): 275 - 280.
- Kfir, R. (2002).** Increase in cereal stem borer populations through partial elimination of natural enemies. *Entomologia Experimentalis et Applicata* 104 (2-3): 299 - 306.
- Kfir, R., Gouws, J. and Moore, S.D. (1993).** Biology of *Tetrastichus howardi* (Olliff) (Hymenoptera: Eulophidae): a facultative hyperparasitoid of stem borers. *Biocontrol Science and Technology* 3 (2): 149 - 159.
- Kfir, R. and Bell, R.A. (1993).** Intraseasonal changes in populations of the African stem borer *Busseola fusca* (Fuller) (Lepidoptera: Noctuidae) and its parasitoids in Natal, South Africa. *Revue de Zoologie Africaine* 107 (6): 543 - 553.
- Kfir, R. and Walters, H.S. (1996).** Transovarial transmission of a microsporidian pathogen in hymenopterous parasitoids of the stem borer *Chilo partellus* (Swinhoe) (Lepidoptera: Pyralidae). *African Entomology* 4 (2): 287 - 290.
- Khajuria, D.R., Sharma, J.P. and Dogra, G.S. (1992).** Role of natural mortality factors in suppression of apple fruit moth, *Argyresthia conjugella* Zeller. *Indian Journal of Plant Protection* 20 (1): 92 - 94.
- Khalaim, A.I. (2001).** A review of the species of the genus *Gelanes* (Hymenoptera, Ichneumonidae, Tersilochinae) of the Palearctic region. *Vestnik Zoologii* 36(6): 3-12.
- Khalaim, A.I. (2002).** A new species of the genus *Phradis* Forster, 1869 from the USA (Hymenoptera: Ichneumonidae: Tersilochinae). *Russian Entomological Journal* 11 (2): 221 - 222.

- Khalaim, A.I. (2002).** A review of the subgenera *Nanodiaparsis*, *Ischnobatis* and *Lanugoparsis* subgen. n., genus *Diaparsis* Forster (Hymenoptera: Ichneumonidae) with descriptions of new species. *Entomologicheskoe Obozrenie* 81 (2): 386-393.
- Khalaim, A.I. (2002).** Two new species of the genus *Barycnemis* Frster from Mexico (Hymenoptera: Ichneumonidae, Tersilochinae). *Zoosystematica Rossica* 11(1): 167-169.
- Khalaim, A.I. (2003).** Review of the Palearctic subgenus *Rugodiaparsis* Horstmann, 1971 of the genus *Probles* Forster, 1869 (Hymenoptera: Ichneumonidae, Tersilochinae). *Russian Entomol. J.* 12(1): 75-78.
- Khalaim, A.I. (2004).** A new species of the genus *Phradis* Forster with 10-11-segmented antenna from the Ukraine (Hymenoptera: Ichneumonidae, Tersilochinae). *Zoosystematica Rossica*, 12(2), 2003: 243-244.
- Khalaim, A.I. (2004).** A review of the Palearctic species of the genera *Barycnemis* Forst., *Epistathmus* Forst. and *Spinolochus* Horst. (Hymenoptera, Ichneumonidae, Tersilochinae). *Proceeding of the Russian Entomol. Soc. St. Petersburg* 2004, 75(1): 46-63.
- Khalaim, A.I. (2004).** New tersilochines from Australia and New Zealand (Hymenoptera: Ichneumonidae, Tersilochinae). *Zoosystematica Rossica* 13(1): 43-45.
- Khan, S.M. (1994).** Food consumption and utilization of *Agrotis segetum* larvae parasitized by *Meloboris collector*. *Sarhad Journal of Agriculture* 10 (2): 187 - 189.
- Khan, S.M. (1997).** Effectiveness of *Meloboris collector* (Hym.: Ichneumonidae) against its host *Agrotis segetum* and *A. ipsilon* (Lepidoptera: Noctuidae). *Sarhad Journal of Agriculture* 13 (1): 77.
- Kienzle, J., Zebitz, C.P.W., Brass, S. and Athanassov, A. (1997).** Abundance of different tortricid species and their parasitoid antagonists in ecological apple orchards in Southern Germany. *Entomological research in organic agriculture. Selected papers from the European Workshop, Austrian Federal Ministry of Science and Research, Vienna, Austria, 14-16 March, 1995. Biological Agriculture and Horticulture* 15 (1-4): 211 - 221.
- Kienzle, J. and Zebitz, C.P.W. (1997).** Extensive mulch system - a contribution towards increasing the diversity of beneficial arthropods in fruit production systems. *Contributions to the 4th Scientific Meeting on Ecological Agriculture, held on 3-4 March 1997, at the Rheinische Friedrich Wilhelms Universitat Bonn. Schriftenreihe Institut fur Organischen Landbau* 4: 192 - 198.
- Kim, M.K., Sisson, G. and Stoltz, D. (1996).** Ichnovirus infection of an established gypsy moth cell line. *Journal of General Virology* 77 (9): 2321 - 2328.
- Kimura, H. and Itoh, N. (1992).** Encapsulation of the eggs of *Bathyplectes curculionis* (Thomson) in larvae of *Hypera postica* (Gyllenhal) in Japan. *Research Bulletin of the Plant Protection Service, Japan* 28: 41 - 45.
- Kingsley, P.C., Bryan, M.D., Day, W.H., Burger, T.L., Dysart, R.J. and Schwalbe, C.P. (1993).** Alfalfa weevil (Coleoptera: Curculionidae) biological control: spreading the benefits. *Environmental Entomology* 22(6): 1234 - 1250.

- Kinuthia, W., Schmidt, O., Theopold, U. and Li, D.M. (1999).** Is the surface of endoparasitic wasp eggs and larvae covered by a limited coagulation reaction? *Journal of Insect Physiology* 45 (5): 501 - 506.
- Klukowski, Z. (1993).** Factors limiting the population of the antler moth, *Cerapteryx graminis* (L.) (Lepidoptera: Noctuidae) in the area of ecological devastation in the Sudety Mts. *Acta Entomologica Lituanica* 11: 187 - 192.
- Klukowski, Z. (2001).** Oilseed rape infestation by stem weevils (*Ceutorhynchus pallidactylus* Marsh, *C. napi* Gyll.) and the occurrence of their parasitoids *Tersilochus* spp. (Ichneumonidae) - spatio-temporal analysis. *Progress in Plant Protection* 41 (2): 459 - 463.
- Knight, G.A., Lavigne, R.J. and Pogue, M.G. (1991).** The parasitoid complex of forest tent caterpillar, *Malacosoma disstria* (Lepidoptera: Lasiocampidae), in eastern Wyoming shelterbelts. *Great Lakes Entomologist* 24 (4): 255 - 261.
- Kodomari, S. (1987).** Use of granulosis viruses for the control of two tea tortricid moths. *Extension Bulletin, ASPAC Food and Fertilizer Technology Center for the Asian and Pacific Region, Taiwan* 257: 22 - 23.
- Kok, L.T. and McAvoy, T.J. (1989).** Fall broccoli pests and their parasites in Virginia. *Journal of Entomological Science* 24 (2): 258 - 265.
- Kolarov, J.A. (1980).** A new cremastine genus and species from Bulgaria. *Acta Zoologica Bulgarica* 16: 90-92.
- Kolarov, J.A. (1986).** A revision of the Orthocentrinae of Bulgaria (Hymenoptera: Ichneumonidae). *Annales Historico-Naturales Musei Nationalis Hungarici* 78: 255-264.
- Kolarov, J.A. (1987).** A new Ctenopelmatinae genus and species from Bulgaria (Hymenoptera, Ichneumonidae). *Entomofauna* 8: 69-76.
- Kolarov, J. (1987).** Taxonomic and faunistic studies on Bulgarian Cremastinae (Hymenoptera: Ichneumonidae). I. *Folia Entomologica Hungarica* 48: 87 - 89.
- Kolarov, J. (1989).** Ichneumonidae (Hymenoptera) from Balkan peninsula and some adjacent regions. II. *Lissonotinae, Ctenopelmatinae, Tersilochinae, Cremastinae and Campopleginae.* *Turkiye Entomoloji Dergisi* 13 (2): 67 - 84.
- Kolarov, J., Ozbek, H. and Yildirim, E. (1999).** New distributional data of the Turkish Ichneumonidae (Hymenoptera). I. *Pimplinae and Tryphoninae.* *Journal of the Entomological Research Society* 1 (2): 9 - 15.
- Kolarov, J. and Beyarslan, A. (1994).** Notes on the Turkish Ichneumonidae (Hymenoptera) III. *Banchinae, Ctenopelmatinae and Tersilonchinae.* *Turkiye III. Biyolojik Mucadele Kongresi Bildirileeri, 25-28 Ocak 1994, Ege Universitesi Ziraat Fakultesi, Bitki Koruma Bolumu, Izmir* 93 - 100.
- Kolarov, J. and Glavendekic, M. (1998).** A review of the genus *Netelia* Gray (Hymenoptera: Ichneumonidae: Tryphoninae) in former Yugoslavia. *Acta Entomologica Serbica* 3 (1-2): 117 - 125.

- Kolarov, J.A. (1992).** New and little known Cremastinae species in Bulgarian fauna and a key for identification of the Western Palearctic genera (Hymenoptera: Ichneumonidae). *Acta Zoologica Bulgarica* 44: 75-78.
- Kolarov, Ya., Stancheva, I. and Sredkov, I. (1987).** A new pest of cherry in Bulgaria - *Pemphredon inornatus* Say (Hymenoptera: Sphecidae). *Pochvoznanie, Agrokhimiya I Rastitelna Zashchita* 22 (5): 95 - 98.
- Kolarov, Ya.A. (1988).** A new parasitoid of the cereal leaf beetle. *Rasteniye i Zashchita* 25 (2): 108 - 110.
- Kolarov, Ya.A. (1989).** Studies on the genus *Diadegma* Foerster, 1868 (Hymenoptera: Ichneumonidae) in Bulgaria. *Acta Zoologica Bulgarica* 37: 67 - 73.
- Kolarov, Ya.A. (1990).** Ecological notes on the species of the subfamily Tersilochinae (Hymenoptera: Ichneumonidae) in Bulgaria. *Ekologiya, Bulgaria* 23: 79 - 83.
- Kolarov, Ya.A. (1990).** Ichneumonidae (Hymenoptera) from Bulgaria in foreign collections. *Acta Zoologica Bulgarica* 39: 25 - 31.
- Kolarov, Ya.A. (1990).** Species of the subfamily Campopleginae (Hymenoptera: Ichneumonidae) new and little known to the Bulgarian fauna. *Acta Zoologica Bulgarica* 39: 32 - 36.
- Kolomiets, N.G. (1986).** Ichneumonids and tachinid flies - parasites of pests of forest in western Siberia. *Sibirskii Biologicheskii Zhurnal* 1: 50 - 53.
- Konishi, K. (1986).** A new subgenus and species of the genus *Netelia* (Hymenoptera, Ichneumonidae) from New Guinea. *Kontyu* 54: 415-449.
- Konishi, K. (1991).** A revision of the subgenus *Prosthodocis* Enderlein of the genus *Netelia* Gray of Japan (Hymenoptera: Ichneumonidae), I. *Japanese Journal of Entomology* 59 (4): 775 - 788.
- Konishi, K. (1992).** A revision of the subgenus *Prosthodocis* Enderlein of the genus *Netelia* Gray of Japan (Hymenoptera: Ichneumonidae), II. *Japanese Journal of Entomology* 60 (1): 39 - 53.
- Konishi, K. (1996).** A revision of the subgenus *Paropheltes* Cameron of the genus *Netelia* Gray (Hymenoptera: Ichneumonidae) of Japan. *Japanese Journal of Entomology* 64 (1): 163 - 187.
- Konishi, K. (1996).** Study on the subgenus *Toxochiloides* Tolkantitz of the genus *Netelia* Gray (Hymenoptera: Ichneumonidae) of Japan. *Japanese Journal of Entomology* 64 (3): 473 - 481.
- Kopelke, J.P. (1986).** *Adelognathus cubiceps* Roman 1924 (Ichneumonidae: Adelognathinae) - an unusual parasitoid of gall-forming *Pontania* species (Tenthredinidae: Nematinae) (Insecta: Hymenoptera). *Senckenbergiana Biologica* 67 (4-6): 253 - 259.
- Koponen, M. (1991).** Contributions to the knowledge of the Braconidae of Finland (Hymenoptera). *Entomologia Fennica* 2 (4): 193 - 209.
- Koptur, S. and Lawton, J.H. (1988).** Interactions among vetches bearing extrafloral nectaries, their biotic protective agents, and herbivores. *Ecology, USA* 69 (1): 278 - 283.

- Kotlabi, H. and Miller, R. (1990).** A survey of wheat stem borer, *Oria musculosa* Hubner (Lepidoptera: Noctuidae), in Syria (1983 - 1987). Arab Journal of Plant Protection 8 (2): 130 - 127.
- Krishnamoorthy, A. and Mani, M. (1998).** New record of parasitoid *Diadegma apostata* (G.) on brinjal shoot and fruit borer. Insect Environment 4 (3): 87.
- Kronstedt, T. (1990).** Separation of two species standing as *Alopecosa aculeata* (Clerck) by morphological, behavioural and ecological characters, with remarks on related species in the pulverulenta group (Araneae: Lycosidae). Zoologica Scripta 19 (2): 203 - 225.
- Kroschel, J. and Koch, W. (1994).** Studies on the population dynamics of the potato tuber moth (*Phthorimaea operculella* Zell. (Lep.: Gelechiidae) in the Republic of Yemen. Journal of Applied Entomology 118 (4-5): 327 - 341.
- Krotova, I.G. (1993).** Parasites of aphidophagous Syrphidae (Diptera) from cereals of western Siberia. Zoologicheskii Zhurnal 72 (12): 58 - 62.
- Kruger, O. and McGavin, G.C. (1998).** The influence of ants on the guild structure of Acacia insect communities in Mkomazi game reserve, north-east Tanzania. African Journal of Ecology 36 (3): 213 - 220.
- Kuhar, T.P., Youngman, R.R. and Laub, C.A. (1999).** Alfalfa weevil (Coleoptera: Curculionidae) pest status and incidence of *Bathyplectes* spp. (Hymenoptera: Ichneumonidae) and *Zoophthora phytonomi* (Zygomycetes: Entomophthorales) in Virginia. Journal of Economic Entomology 92 (5): 1184 - 1189.
- Kuhlmann, U. (1996).** Biology and ecology of *Herpestomus brunnicornis* (Hymenoptera: Ichneumonidae), a potential biological control agent of the apple ermine moth (Lepidoptera: Yponomeutidae). International Journal of Pest Management 42 (2): 131 - 138.
- Kuhlmann, U., Polesny, F., Muller, W. and Olszak, R.W. (1996).** Measuring and understanding the impact of natural enemies on apple pest populations: life tables for the apple ermine moth, *Yponomeuta malinellus*. International conference on integrated fruit production, at Cedzyna, Poland, 28 August - 2 September 1995. Bulletin OILB SROP 19 (4): 197 - 202.
- Kumagai, A.F. (2002).** The Ichneumonidae (Hymenoptera) of the Estacao Ecologica of the Universidade Federal de Minas Gerais, Belo Horizonte, with emphasis on the Pimplinae species. Revista Brasileira de Entomologia 46 (2): 189 - 194.
- Kumagai, A.F. and Graf, V. (2002).** Biodiversity of Ichneumonidae (Hymenoptera) and monitoring of the species of Pimplinae and Poemeniinae from Capao da Imbuia, Curitiba, Parana. Revista Brasileira de Zoologia 19 (2): 445 - 452.
- Kumar, A., Kapoor, V.C. and Mahal, M.S. (1989).** Pupal population of aphidophagous syrphids and their hyperparasitism on Brassica juncea Coss. Journal of Insect Science 2 (2): 114 - 118.
- Kumar, N., Kumar, A. and Tripathi, C.P.M. (1994).** Functional response of *Campoletis chloridae* Uchida (Hymenoptera: Ichneumonidae), a parasitoid of *Heliothis armigera*



- (Hubner) (Lepidoptera: Noctuidae) in an enclosed experimental system. *Biological Agriculture and Horticulture* 10 (4): 287 - 295.
- Kumar, P., Singh, S.P., Chandish., Ballal, R. and Jalali, S.K. (1988).** Relationship between the host age and the fitness components of *Hyposoter didymator* Thunb. (Hymenoptera: Ichneumonidae). *Journal of Biological Control* 2 (2):69 - 71.
- Kumar, P. and Ballal, C.R. (1992).** The effect of parasitism by *Hyposoter didymator* (Hym.: Ichneumonidae) on food consumption and utilization by *Spodoptera litura* (Lep.: Noctuidae). *Entomophaga* 37 (2): 197 - 203.
- Kumar, S.P., Kaushal, P. and Chauhan, U. (1999).** Natural enemies of bud borer *Helicoverpa armigera* (Hubner) in carnation. *Insect Environment* 5 (2): 52 - 53.
- Kuppusamy, A. and Kannan, S. (1993).** Life-history of *Eumeta cramerii* (Westwood) (Lepidoptera: Psychidae) and its natural enemy *Sinophorus psycheae* Sonan (Hymenoptera: Ichneumonidae). *Phytophaga Madras* 5 (2): 109 - 121.
- Kusigemati, K. (1982).** A new porizontine genus *Breviterebra* from Japan (Hymenoptera, Ichneumonidae). *Kontyu* 50: 621-625.
- Kusigemati, K. (1988).** Descriptions of two new ichneumonflies (Hymenoptera) parasitic on economic hosts from Nepal. *Kontyu* 56 (4): 812 - 816.
- Kusigemati, K. (1988).** New host records of Ichneumonidae (Hymenoptera), with descriptions of two new species from Japan (VI). *Kontyu* 56 (3): 573 - 580.
- Kusigemati, K. (1989).** Descriptions of three new ichneumonflies parasitic on lycaenid butterflies in Japan (Hymenoptera: Ichneumonidae). *Memoirs of the Faculty of Agriculture, Kagoshima University* 25: 83 - 89.
- Kusigemati, K. (1990).** Descriptions of two new ichneumon flies (Hymenoptera) parasitic on alfalfa weevil, *Hypera postica* (Gyllenhal) (Coleoptera: Curculionidae) from Japan. *Japanese Journal of Entomology* 58 (3): 619 - 624.
- Kusigemati, K. (1993).** Description of a new ichneumonfly (Hymenoptera) parasitic on *Amphipoea* spp. (Lepidoptera: Noctuidae) from Japan. *Memoirs of the Faculty of Agriculture, Kagoshima University* 29: 89 - 91.
- Kusigemati, K. (1993).** Descriptions of three new Ichneumonflies (Hymenoptera) parasitic on economic hosts from Japan. *Japanese Journal of Entomology* 61 (2): 235 - 242.
- Kusigemati, K. (1993).** Descriptions of two new ichneumonflies (Hymenoptera) parasitic on diamondback moth, *Plutella xylostella* L. (Lepidoptera: Yponomeutidae) from Japan. *Japanese Journal of Entomology* 61 (1): 101 - 107.
- Kusigemati, K. and Tanaka, A. (1992).** New host records of Ichneumonidae (Hymenoptera), with a description of a new Eriborus species from Japan (VIII). *Memoirs of the Faculty of Agriculture, Kagoshima University* 28: 83 - 88.
- Kuslitzky, (1973).** Ichneumon-flies of the genera *Teleutaea* Foerst., *Cephaloglypta* Obrt., and *Zygoglypta* Mom. (Hymenoptera, Ichneumonidae) in the Soviet fauna. (in Russian) *Éntomologicheskoe Obozrenie* 52: 886-898.

- Kuslitskii, V.S., Livshits, I.Z. and Mitrofanov, V.I. (1987).** Hymenopterous parasitoids of pests of fruit orchards. *Zashchita Rastenii Moskva* 9: 49 - 52.
- Kuslitskii, V.S., Livshits, I.Z. and Mitrofanov, V.I. (1987).** Hymenopterous parasitoids of pests of fruit orchards. *Zashchita Rastenii Moskva* 11: 49 - 53.
- Kuznetsova, Yu.I. and Kozhechkin, O.A. (1986).** The formation of an entomophage complex. *Zashchita Rastenii* 6; 40 - 41.

## L

- Laba, I.W. and Kartohardjono, A. (1998).** Conservation of parasitoid and predator within plant pest management. *Jurnal Penelitian and Pengembangan Pertanian* 17 (4): 122 - 129.
- Labeyrie, V. (1991).** Influence of polymorphism on trophic relations of *Diadromus pulchellus* (Ichneumonidae). *Redia* 74 (3): Appendix, 211 - 215.
- Lagowska, B. and Winiarska, W. (1982).** Parasitic Hymenoptera reared from the larvae of the apple weevil, *Anthonomus pomorum* L., collected in the Lublin region. *Roczniki Nauk Rolniczych, E Ochrona Roslin*, publ. 1986, 12 (1-2): 121 - 126.
- Lale, N.E.S. and Igwebuiké, J.U. (2002).** Field infestation of *Faidherbia (Acacia) albida* (Del.) A. Chew. pods by stored product Coleoptera in the Nigerian savanna and effect of infestation on nutrient quality. *Journal of Arid Environments* 51 (1): 103 - 112.
- Landis, D.A. and Haas, M.J. (1992).** Influence of landscape structure on abundance and within-field distribution of European corn borer (Lepidoptera: Pyralidae) larval parasitoids in Michigan. *Environmental Entomology* 21 (2): 409 - 416.
- Lane, S.D. and Mills, N.J. (2003).** Intraspecific competition and density dependence in an *Ephestia kuehniella* - *Venturia canescens* laboratory system. *Oikos* 101 (3): 578 - 590.
- Lanfranco, D. and Cerda, L. (1986).** *Coccygomimus [Pimpla] fuscipes*, an indigenous ichneumon parasite of the budworm *Rhyacionia buoliana*. *Bosque* 7 (1): 36 - 37.
- Lanfranco, D.M., Aguilar, A.M., Horcos, R.A. and Buchner, J.F. (1994).** Native parasitoids in the control of pine shoot moth (*Rhyacionia buoliana*): a functional complex? *Bosque* 15 (1): 15 - 26.
- Lanfranco, D.M., Buchner, J.F., Aguilar, A.M. and Horcos, R.A. (1991).** Native parasitoids in control of the pine shoot moth *Rhyacionia buoliana*: progress in identifying the complex and its development strategies. *Bosque* 12 (1): 69 - 74.
- Langor, D.W., Digweed, S.C., Williams, D.J.M., Spence, J.R. and Saunders, C. (2000).** Establishment and spread of two introduced parasitoids (Ichneumonidae) of the birch leafminer, *Fenusa pusilla* (Tenthredinidae). *BioControl* 45 (4): 415 - 423.
- Lannic, J.Le, and Nenon, J.P. (1999).** Functional morphology of the ovipositor in *Megarhyssa atrata* (Hymenoptera: Ichneumonidae) and its penetration into wood. *Zoomorphology* 119 (2): 73 - 79.
- Lapa, A.M. and Tkachev, V.M. (1992).** Ecological aspects of an integrated pest management system in orchards. *Acta Phytopathologica et Entomologica Hungarica* 27 (1-4): 401 - 403.

- Laulier, M., Pradier, E., Bigot, Y. and Periquet, G. (1995).** An easy method for preserving nucleic acids in field samples for later molecular and genetic studies without refrigerating. *Journal of Evolutionary Biology* 8 (5): 657 - 663.
- Laurenne, N.M., Belshaw, R., Broad, G. and Quicke, D.L.J. (2000).** Molecular confirmation of host records for ichneumonoid parasitoids of wood-boring beetle larvae. *Journal of Hymenoptera Research* 9 (2): 241 - 245.
- Lavine, M.D. and Beckage, N.E. (1995).** Polydnviruses: potent mediators of host insect immune dysfunction. *Parasitology Today* 11 (10): 368 - 378.
- Leclercq, M. (1996).** On the entomofauna of a wild boar carcass. *Bulletin and Annales de la Societe Royale Belge d'Entomologie* 132 (4): 417 - 422.
- Lecomte, C., and Thibout, E. (1991).** Learning and conditioning in a specialist parasitoid *Diadromus pulchellus* (Ichneumonidae). *Redia* 74 (3): Appendix, 305 - 307.
- Lecomte, C. and Thibout, E. (1993).** Pre- and post-imaginal experience in a specialist parasitoid *Diadromus pulchellus* (Hym.: Ichneumonidae). *Entomophaga* 38 (2): 175 - 184.
- Lee, J.W. and Kim, C.W. (1984).** Studies on the Ichneumonidae from Korea. V. A new genus *Pseudagrypon* of Anomaloninae (Hymenoptera: Ichneumonidae). *Korean Journal of Entomology* 14: 31-34.
- Lee, J.H., Reed, D.K., Lee, H.P. and Carlson, R.W. (1987).** The parasite complex of gypsy moth (*Lymantria dispar* L.) at Mt. Chonggye. *Korean Journal of Entomology* 17 (4): 245 - 246.
- Lee, J.W. (1991).** A revision of the genus *Cidaphus* (Hymenoptera: Ichneumonidae). *Contributions of the American Entomological Institute* 26 (5): 1 - 48.
- Lee, J.W. (1992).** A revision of species of the B-group of *Astiphromma* (Hymenoptera: Ichneumonidae: Mesochorinae). *Oriental Insects* 26: 213 - 239.
- Lee, J.W. (1992).** A revision of the genus *Plectochorus* (Hymenoptera: Ichneumonidae: Mesochorinae). *Oriental Insects* 26: 241 - 263.
- Lee, J.W., Cha, J.Y. and Suh, K.I. (1995).** Taxonomy and phylogeny of the subfamily Tryphoninae (Hymenoptera: Ichneumonidae). II. A review of tribe Exenterini. *Entomological Research Bulletin* 21: 1 - 18.
- Lee, J.W., Suh, K.I. and Cha, J.Y. (1994).** A systematic study of the Ichneumonidae (Hymenoptera) from Korea XVI. The genus *Astiphromma* Foerster (Mesochorinae). *Korean Journal of Entomology* 24 (1): 19 - 30.
- Lee, J.W. and Cha, J.Y. (1993).** A systematic study of the Ichneumonidae (Hymenoptera) from Korea XV. Review of the Tribe Tryphonini (Tryphoninae). *Entomological Research Bulletin* 19: 10 - 34.
- Lee, J.W. and Kim, K.B. (2002).** Taxonomical review of the subfamily Ophioninae (Hymenoptera: Ichneumonidae) I. Genus *Stauropoctonus* Brauns. *Korean Journal of Entomology* 32 (2): 81 - 86.

- Lee, J.W. and Suh, K.I. (1991).** A systematic study of the Ichneumonidae (Hymenoptera) from Korea XIII. Genus *Mesochorus* (Mesochorinae). Entomological Research Bulletin 17: 11 - 32.
- Lee, J.W. and Suh, K.I. (1993).** A systematic study of Ichneumonidae (Hymenoptera) from North Korea I. Subfamily Mesochorinae. Annales Historico Naturales Musei Nationalis Hungarici 85: 141 - 153.
- Lee, S.G., Yoo, J.K., Lee, J.O. and Hwang, C.Y. (1995).** The biological characteristics and DBM density suppression effect of *Diadegma semiclausum* Hellen. RDA Journal of Agricultural Science, Crop Protection 37 (1): 329 - 333.
- Lei, G.C., Vikberg, V., Nieminen, M. and Kuussaari, M. (1997).** The parasitoid complex attacking Finnish populations of the *Glanville fritillary Melitaea cinxia* (Lep.: Nymphalidae), an endangered butterfly. Journal of Natural History 31 (4): 635 - 648.
- Leluk, J., Schmidt, J. and Jones, D. (1989).** Comparative studies on the protein composition of hymenopteran venom reservoirs. Toxicon Oxford 27 (1): 105 - 114.
- Li, C.R., Fan, M.Z., Huang, B., Wang, S.B. and Li, Z.Z. (2002).** The genus *Cordyceps* and its allies from Anhui I. Mycosystema 21 (2): 167 - 171.
- Li, D., Blasevich, F., Theopold, U. and Schmidt, O. (2003).** Possible function of two insect phospholipid-hydroperoxide glutathione peroxidases. Journal of Insect Physiology 49 (1): 1 - 9.
- Li, D., Zhao, Z., Roberts, H., Schneider, M.V., Theopold, U. and Schmidt, O. (2003).** Genetic analysis of two distinct reproductive strategies in sexual and asexual field populations of an endoparasitic wasp, *Venturia canescens*. Heredity 90 (4): 291 - 297.
- Li, X., Liu, S.S. and Wang, D. (2002).** Effects of host plant on the host-selection behavior of *Diadegma semiclausum*. Chinese Journal of Biological Control 18 (4): 145 - 148.
- Li, X. and Liu, S.S. (2002).** The role of plant volatiles and oviposition experience in the foraging behaviour of *Diadegma semiclausum* (Hym.: Ichneumonidae). Journal of Zhejiang University Agriculture and Life Sciences 28 (3): 293 - 297.
- Lim, G.S. (1986).** The effects of parasitoids on *Plutella xylostella* (L.) in the Cameron highlands. Biological control in the tropics. Regional Symposium on Biological Control, Serdang (Malaysia). 163-172.
- Lin, T.C. (1990).** The abundance and diversity of hymenopterans in Ulu Kinchin, Pahang, Malaysia. Malayan Nature Journal 43 (4): 274 - 284.
- Lind, P. (1998).** Solving yellowjacket problems. Journal of Pesticide Reform 18: (1): 22 - 23.
- Link, D., Costa, E.C. and Correa-Costa, E. (1988).** Biological aspects of wheat armyworms. Revista do Centro de Ciencias Rurais, Universidade Federal de Santa Maria 18 (2): 99 - 105.
- Link, D. and Costa, E.C. (1989).** Effectiveness of light traps for the control of the cucurbit borer *Diaphania* spp. (Lepidoptera: Pyralidae) in Santa Maria, RS. Revista do Centro de Ciencias Rurais 19 (4): 311 - 315.

- Littlewood, S.C. (1988).** Notes on the overwintering *Apanteles-Tetrastichus-Lysibia* complex (Hymenoptera: Parasitica), parasitoids of the large white butterfly. *Journal of Natural History* 22 (4): 883 - 895.
- Liu, S.S., Wang, X.G., Guo, S.J., He, J.H. and Shi, Z.H. (2000).** Seasonal abundance of the parasitoid complex associated with the diamondback moth, *Plutella xylostella* (Lepidoptera: Plutellidae) in Hangzhou, China. *Bulletin of Entomological Research* 90 (3): 221 - 231.
- Lobaton, M.C.S. (1990).** On the dispersion of muscicolous nematodes by insects. *Boletim da Sociedade Portuguesa de Entomologia* 120: 209 - 221.
- Lockard, E. I. (1995).** Biodiversity and geographic distributions of parasitic Hymenoptera (Ichneumonidae: Campopleginae and Ichneumoninae) along a latitudinal gradient in eastern North America. M. S. thesis, Univ. of Georgia, Athens, 233pp.
- Lockley, T.C. and Young, O.P. (1993).** Survivability of overwintering *Argiope aurantia* (Araneidae) egg cases, with an annotated list of associated arthropods. *Journal of Arachnology* 21 (1): 50 - 54.
- Logan, J.A. and Hain, F.P. (1991).** Chaos and insect ecology. Information Series Virginia Polytechnic Institute and State University, College of Agriculture and Life Sciences 91-3: 108 pp.
- Long, R.F., Corbett, A., Lamb, C., Reberg-Horton, C., Chandler, J. and Stimmann, M. (1998).** Beneficial insects move from flowering plants to nearby crops. *California Agriculture* 52 (5): 23 - 26.
- Lopez, M., Rojas, J.C., Vandame, R. and Williams, T. (2002).** Parasitoid-mediated transmission of an iridescent virus. *Journal of Invertebrate Pathology* 80 (3): 160 - 170.
- Losey, J.E., Song, P.Z., Schmidt, D.M., Calvin, D.D. and Liewehr, D.J. (1992).** Larval parasitoids collected from overwintering European corn borer (Lepidoptera: Pyralidae) in Pennsylvania. *Journal of the Kansas Entomological Society* 65 (1): 87 - 90.
- Lourencao, A.L., De Carvalho, L.O., Lasca, D.H.C. and De Campos-Lasca, D.H. (1989).** *Anacraga citrinopsis* Dyar (Lepidoptera: Dalceridae) on castor in the state of Sao Paulo. *Bragantia* 48 (1): 109 - 112.
- Lovalo, N., McPherson, B.A. and Cox-Foster, D.L. (2002).** Effects of the polydnavirus of *Cotesia congregata* on the immune system and development of non-habitual hosts of the parasitoid. *Journal of Insect Physiology* 48 (5): 517 - 526.
- Lozzia, G.C. and Rigamonti, I.E. (1991).** Observations on strategies of biological and integrated control of grape moths in northern Italy. *Vignevini* 18 (11): 33 - 37.
- Luckhart, S. and Webb, B.A. (1996).** Interaction of a wasp ovarian protein and Polydnavirus in host immune suppression. *Developmental and Comparative Immunology* 20 (1): 1 - 21.
- Lugojja, F., Ogenga-Latigo, M.W. and Smit, N.E.J.M. (2001).** Species diversity and activity of parasitoids of the sweetpotato butterfly, *Acraea acerata*, in Uganda. *Fifth*

Triennial Congress of the African Potato Association, 28 May - 2 June 2000, Kampala, Uganda. African Crop Science Journal 9 (1): 157 - 163.

**Luhman, J.C. (1990).** A taxonomic revision of Nearctic *Endasys* Foerster, 1868 (Hymenoptera: Ichneumonidae: Gelinae). University of California Publications in Entomology 109: 185 pp.

**Luhman, J.C. (1991).** A revision of the world *Amphibulus* Kriechbaumer (Hymenoptera: Ichneumonidae: Phygadeuontinae). Insecta Mundi 5 (3-4): 129 - 152.

**Lukas, J., Zd'arkova, E., Wakefield, M., Lukas, J. and Hubert, J. (2002).** Parasitoids occurring in food-processing factories and grain stores. COST Action 842, 1999 - 2004. Biological control of pest insects and mites with special reference to entomophthorales. Proceedings of the Second Meeting of Working Group 4:-Bio control of arthropod pests in the stored products, Prague, Czech Republic, 30 - 31<sup>st</sup> May 2002. 83 - 86.

**Luo, W.D. and Qin, D.Y. (1993).** A new species of *Pleolophus* Townes from Beijing (Hymenoptera: Ichneumonidae: Gelinae). Entomotaxonomia 15 (1): 65 - 68.

**Luo, W.D. and Qin, D.Y. (1995).** Two new species of *Pleolophus* Townes (Hymenoptera: Ichneumonidae: Gelinae) from Beijing, China. Entomotaxonomia 17 (2):147 - 151.

**Lyons, D.B. (1999).** Phenology of the native parasitoid *Sinophorus megalodontis* (Hymenoptera: Ichneumonidae) relative to its introduced host, the pine false webworm (Hymenoptera: Pamphiliidae). Canadian Entomologist 131 (6): 787 - 800.

**Lysyk, T.J. and Nealis, V.G. (1988).** Temperature requirements for development of the Jack pine budworm (Lepidoptera: Tortricidae) and two of its parasitoids (Hymenoptera). Journal of Economic Entomology 81 (4): 1045 - 1051.

## M

**Ma, Q., Wang, D.Y. and Wang, S.L. (1995).** Three new species of genus *Diplazon* Viereck (Hymenoptera: Ichneumonidae: Diplazontinae) from Xinjiang, China. Entomotaxonomia 17 (1): 51 - 58.

**Ma, Q., Ma, Q., Wang, D.Y. and Wang, S.L. (1995).** Three new species of genus *Diplazon* Viereck (Hymenoptera: Ichneumonidae: Diplazontinae) from Xinjiang, China. Entomotaxonomia 17 (1): 51 - 58.

**Ma, R.Z., Swedenborg, P.D. and Jones, R.L. (1992).** Host-seeking behaviour of *Eriborus terebrans* (Hymenoptera: Ichneumonidae) toward the European corn borer and the role of chemical stimuli. Annals of the Entomological Society of America 85 (1): 72 - 79.

**Ma, W.Y., Peng, J.W., Wang, X.L. and Zuo, Y.X. (1989).** Biology and population fluctuations of *Casinarina nigripes* Gravenhorst. Acta Ecologica Sinica 9 (1): 27 - 35.

**Ma, W.Y., Peng, J.W., Wang, X.L. and Zuo, Y.X. (1989).** Biology and population fluctuation of *Casinarina nigripes* Gravenhorst. Acta Ecologica Sinica 9 (1): 27 - 34.

**Machuca, J.R., Arretz, P. and Araya, J. (1989).** Presence of *Campoletis sonorensis* (Cameron, 1886) (Hymenoptera: Ichneumonidae) in Chile. Acta Entomologica Chilena 15: 269 - 270.

- Machuca, J.R.L., Arretz, P.V. and Araya, J.E.C. (1988).** Parasitism of noctuids in cultivated artichokes in the metropolitan region: identification and preliminary observations of the parasites. *Revista Chilena de Entomologia* 16: 83 - 87.
- Mackay, A.I. and Kring, T.J. (1998).** Acceptance and utilization of diapausing *Helicoverpa zea* (Lepidoptera: Noctuidae) pupae by *Ichneumon promissorius* (Hymenoptera: Ichneumonidae). *Environmental Entomology* 27 (4): 1006 - 1009.
- Madrigal, C.A. (1989).** A critical view of forest pest management in Colombia. *Cronica Forestal y del Medio Ambiente* 7: 43 - 57.
- Maier, K. (1990).** Contribution to the biology of primary and secondary parasitoids of *Lymantria dispar* L. (Lep.: Lymantriidae). *Journal of Applied Entomology* 110 (2): 167 - 182.
- Maier, K. and Bogenschutz, H. (1988).** The importance of caterpillar flies (Diptera: Tachinidae) in the collapse of an outbreak of *Lymantria dispar* L. (Lep.: Lymantriidae). *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 6 (1-3): 164 - 168.
- Maier, K. and Bogenschutz, H. (1990).** Population dynamics of *Lymantria dispar* L. (Lep.: Lymantriidae) and the influence of parasitoids during an outbreak in southwest Germany 1984-86. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 97 (4): 381 - 393.
- Maier, K.J. (1995).** The influence of parasitoids on *Lymantria dispar* (Lepidoptera: Lymantriidae) in woodlands with different population densities and fluctuations. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 10 (1-6): 129 - 134.
- Maingay, H.M., Bugg, R.L., Carlson, R.W. and Davidson, N.A. (1991).** Predatory and parasitic wasps (Hymenoptera) feeding at flowers of sweet fennel (*Foeniculum vulgare* Miller var. *dulce* Battandier & Trabut, Apiaceae) and spearmint (*Mentha spicata* L., Lamiaceae) in Massachusetts. *Biological Agriculture and Horticulture* 7 (4): 363 - 383.
- Maini, S. and Burgio, G. (1990).** The parasitoids of *Ostrinia nubilalis* (Hb.) in Emilia Romagna. *Informatore Fitopatologico* 40 (9): 19 - 28.
- Maleque, M.A., Islam, M.N., Kundu, R. and Islam, M.S. (1999).** Impact of chemical insecticides on the natural enemies of brinjal shoot and fruit borer. *Bangladesh Journal of Entomology* 9 (1-2): 53 - 60.
- Mallik, S.N., Kumar, M., Sinha, A.N. and Karn, B.P. (1989).** *Trathala flavo-orbitalis* Cameron (Ichneumonidae) - a parasite of *Leucinodes orbonalis* Guen. from Bihar. *Current Science* 58: 19, 1098 - 1099.
- Malmberg, T., Beukeboom, L.W., Driessen, G., Van Alphen, J.J.M., Sommeijer, M.J. and Meeuwsen, F.J.A.J. (2000).** Distribution of a VLP-protein polymorphism in sexual and asexual *Venturia canescens* populations (Hymenoptera). *Proceedings of the 11th meeting of experimental and applied entomologists in the Netherlands, Wageningen, Netherlands, 17 December 1999. Proceedings of the Section Experimental and Applied Entomology of the Netherlands Entomological Society* 11: 89 - 93.

- Mamedov, Z.M. (1988).** Parasitoids of the brown-tail moth (*Euproctis chrysorrhoea* L.) and the gypsy moth (*Lymantria dispar* L.). in orchards of Azerbaidzhan. Izvestiya Akademii Nauk Azerbaidzhanskoi SSR, Biologicheskikh Nauk 4: 75 - 77.
- Man, J., Bai, L.Y. and Chen, Y.N. (2000).** Studies on the bionomics and thermal constant of beet armyworm *Spodoptera exigua* (Hubner). Acta Phytophylacica Sinica 27 (3): 215 - 220.
- Mangum, W.A. (2001).** Honey bee biology. American Bee Journal 141 (10): 715 - 717.
- Mani, M., Gopalakrishnan, C. and Krishnamoorthy, A. (2001).** Natural enemies of ber hairy caterpillar *Thiacidas postica* Walker (Lepidoptera: Noctuidae) in Karnataka. Entomon 26 (3-4): 313 - 315.
- Mannion, C.M., Carpenter, J.E., Wiseman, B.R. and Gross, H.R. (1994).** Host corn earworm (Lepidoptera: Noctuidae) reared on meridic diet containing silks from a resistant corn genotype on *Archytas marmoratus* (Diptera: Tachinidae) and *Ichneumon promissorius* (Hymenoptera: Ichneumonidae). Environmental Entomology 23 (4): 837 - 845.
- Manojlovic, B. (1989).** Distribution and importance of parasitoids of the corn borer *Ostrinia nubilalis* Hbn. (Lepidoptera: Pyralidae) in maize hybrids of different FAO maturity groups. Zastita Bilja 40 (2): 115 - 130.
- Manojlovic, B., Sivcev, I., Draganic, M. and Baca, F. (1994).** Investigations so far on European corn borer entomophages (*Ostrinia nubilalis* Hbn., Lepidoptera: Pyralidae) in Yugoslavia. Zastita Bilja 45 (2): 81 - 90.
- Mao, H.X. and Kunimi, Y. (1991).** Pupal mortality of the oriental tea tortrix, *Homona magnanima* Diakonoff (Lepidoptera: Tortricidae), caused by parasitoids and pathogens. Japanese Journal of Applied Entomology and Zoology 35 (3): 241 - 245.
- Marchesini, E., Tosi, L. and Galbero, G. (1998).** Natural enemies of *Hyphantria cunea* in Veneto. Informatore Agrario 54: 15, 115 - 119.
- Marchesini, E. and Monta, L.D. (1994).** Observations on natural enemies of *Lobesia botrana* (Den. & Schiff.) (Lepidoptera: Tortricidae) in Venetian vineyards. Bollettino di Zoologia Agraria e di Bachicoltura 26 (2): 201 - 230.
- Mariau, D. (1997).** Role of parasitoids in controlling oil palm leaf-eating insects. Plantations, Recherche, Developpement 4 (5): 297 - 303.
- Markov, V.A. (2000).** On the migration of *Tortrix viridana* L. (Lepidoptera: Tortricidae): analytical investigation. Zhurnal Obshechi Biologii 61(2): 206 - 224.
- Marris, G.C., Bell, H.A., Naylor, J.M. and Edwards, J.P. (1999).** The role of *Pimpla hypochondriaca* venom in the suppression of pupal noctuid host immunity. Entomologia Experimentalis et Applicata 93 (3): 291 - 298.
- Marris, G.C., Hubbard, S.F. and Scrimgeour, C. (1996).** The perception of genetic similarity by the solitary parthenogenetic parasitoid *Venturia canescens*, and its effects on the occurrence of superparasitism. Entomologia Experimentalis et Applicata 78 (2): 167 - 174.



- Marris, G.C., Naylor, J., Parkinson, N. and Edwards, J.P. (1994).** Wasp-induced infections in a parasitoid - host association. Proceedings - Brighton Crop Protection Conference, Pests and Diseases, 1994, vol 3: 1145 - 1150.
- Martinek, V. (1988).** Parasites of web-spinning sawflies of the genus *Cephalcia* in the Czech Socialist Republic. II. Voltinism of the ichneumonids *Olesicampe monticola* and *Sinophorus crassifemur*. Lesnictvi 34 (6): 547 - 573.
- Martinek, V. (1989).** Parasites of the sawflies of the genus *Cephalcia* spp. in the Czech Republic IV. Voltinism of the ichneumonid wasps of the genus *Homaspis*, and the species *Notopygus nigricornis*. Lesnictvi 35 (9): 783 - 806.
- Martinek, V. (1989).** Parasites of web-spinning sawflies of the genus *Cephalcia* in the Czech Socialist Republic. III. Voltinism of the ichneumon wasps *Xenoschesis fulvipes* and *Ctenopelma nigrum*. Lesnictvi 35 (3): 193 - 212.
- Maruya, W.I., De Pinto, S.A. and Gravena, S. (2001).** Parasitism and *Nomuraea rileyi* (Farlow) Samson on defoliator caterpillar (Lepidoptera) in soyabean crop. Boletín de Sanidad Vegetal, Plagas 27 (4): 561 - 567.
- Mason, W.R.M. (1962).** Some new Asiatic species of Exenterini (Hymenoptera: Ichneumonidae) with remarks on generic limits. Canadian Entomologist 94: 1273-1296.
- Mason, W.R.M. (1975).** A new genus of Pimplini from Jamaica (Hymenoptera: Ichneumonidae). Proceedings of the Entomological Society of Washington 77: 225-227.
- Mason, C.E., Romig, R.F., Wendel, L.E. and Wood, L.A. (1994).** Distribution and abundance of larval parasitoids of European corn borer (Lepidoptera: Pyralidae) in the East Central United States. Environmental Entomology 23 (2): 521 - 531.
- Mason, P.G., Erlandson, M.A. and Youngs, B.J. (2001).** Effects of parasitism by *Banchus flavescens* (Hymenoptera: Ichneumonidae) and *Microplitis mediator* (Hymenoptera: Braconidae) on the bertha armyworm, *Mamestra configurata* (Lepidoptera: Noctuidae). Journal of Hymenoptera Research 10 (1): 81 - 90.
- Mason, W.R.M. and Ross, H.H. (1990).** Cubitus posterior in Hymenoptera. Proceedings of the Entomological Society of Washington 92 (1): 93 - 97.
- Masumoto, T., Shimizu, I. and Konishi, K. (2002).** The first record of parasitism on *Neoscona scylloides* (Araneae: Araneidae) by *Eriostethus rufus* (Hymenoptera: Ichneumonidae). Acta Arachnologica 51 (1): 5 - 6.
- Matama-Kauma, T., Kyamanywa, S., Ogwang, J.A., Omwega, C.O. and Willson, H.R. (2001).** Cereal stemborer species complex and establishment of *Cotesia flavipes* Cameron in Eastern Uganda. Insect Science and its Application 21 (4): 317 - 325.
- Mateias, M.C. and Danulescu, M. (1990).** Observations on *Oligia strigilis* L. (Lepidoptera: Noctuidae) on perennial Gramineae. Analele, Institutului de Cercetari pentru Cereale si Plante Tehnice, Fundulea 58: 303 - 308.
- Mathew, G. (1989).** Natural enemies of the bagworm, *Pteroma plagiophleps* Hampson (Lepidoptera: Psychidae) in Kerala (India). Entomon 14 (3-4): 335 - 338.

- Mathur, K.K. and Singh, Y. (1991).** Parasitisation of *Heliothis armigera* (Hb.) on Bengal gram in Delhi. *Indian Journal of Entomology* 53 (1): 128 - 133.
- Matsumoto, R. and Saigusa, T. (1999).** Biology of *Pseudopimpla glabripropodeum* He et Chen (Hymenoptera: Ichneumonidae: Pimplinae) with description of final instar larva and redescription of adults. *Entomological Science* 2 (4): 467 - 471.
- Maund, C.M. and Hsiao, T.H. (1991).** Differential encapsulation of two *Bathyplectes* parasitoids among alfalfa weevil strains, *Hypera postica* (Gyllenhal). *Canadian Entomologist* 123 (1): 197 - 203.
- Mayo-Jimenez, P.A.P. (1985).** Preliminary biological control trials with the parasitic wasp *Endasys subclavatus* (Hymenoptera: Ichneumonidae) in prepupae of *Zadiprion vallicola* [a pest of pines] in the state of Michoacan. *Boletin Tecnico, Instituto Nacional de Investigaciones Forestales, Mexico* 134: 23 pp.
- Mazanec, Z. (1990).** Immature stages and life history of *Enytus* sp. (Hymenoptera: Ichneumonidae), a parasitoid of *Perthida glyphopa* Common (Lepidoptera: Incurvariidae). *Journal of the Australian Entomological Society* 29 (1): 57 - 66.
- McAuslane, H.J., Vinson, S.B. and Williams, H.J. (1990).** Effect of host diet on flight behavior of the parasitoid *Campoletis sonorensis* (Hymenoptera: Ichneumonidae). *Journal of Entomological Science* 25 (4): 562 - 570.
- McAuslane, H.J., Vinson, S.B. and Williams, H.J. (1990).** Influence of host plant on mate location by the parasitoid *Campoletis sonorensis* (Hymenoptera: Ichneumonidae). *Environmental Entomology* 19 (1): 26 - 31.
- McAuslane, H.J., Vinson, S.B. and Williams, H.J. (1991).** Influence of adult experience on host microhabitat location by the generalist parasitoid, *Campoletis sonorensis* (Hymenoptera: Ichneumonidae). *Journal of Insect Behavior* 4 (1): 101 - 113.
- McCloskey, C., Arnason, J.T., Donskov, N., Chenier, R., Kaminski, J. and Philogene, B.J.R. (1993).** Third trophic level effects of azadirachtin. *Canadian Entomologist* 125 (1): 163 - 165.
- McDonald, R.C. and Kok, L.T. (1991).** Hyperparasites attacking *Cotesia glomerata* (L.) and *Cotesia rubecula* (Marshall) (Hymenoptera: Braconidae) in Southwestern Virginia. *Biological Control* 1 (2): 170 - 175.
- McDonald, R.C. and Kok, L.T. (1992).** Colonization and hyperparasitism of *Cotesia rubecula* (Hym.: Braconidae), a newly introduced parasite of *Pieris rapae*, in Virginia. *Entomophaga* 37 (2): 223 - 228.
- McDougall, C., Philogene, B.J.R., Arnason, J.T. and Donskov, N. (1988).** Comparative effects of two plant secondary metabolites on host-parasitoid association. *Journal of Chemical Ecology* 14 (4): 1239 - 1252.
- McGeoch, M.A. (1993).** The Microlepidoptera associated with a fungus gall on *Acacia karroo* Hayne in South Africa. *African Entomology* 1 (1): 49 - 56.

- McKay, T. and Galloway, T.D. (1999).** Survey and release of parasitoids (Hymenoptera) attacking house and stable flies (Diptera: Muscidae) in dairy operations. *Canadian Entomologist* 131 (6): 743 - 756.
- McNeil, J.N. and Turgeon, J.J. (1988).** Primary parasitoids of the true armyworm, *Pseudaletia unipuncta* (Lepidoptera: Noctuidae), in Quebec. *Revue d'Entomologie du Quebec* 33 (1-2): 1 - 8.
- Megahed, M.M., Abou-Zeid, N.A. and Abbas, M.S.T. (1990).** Biology of *Diadegma molliplum* Hlmgrn., a larval parasite of the potato-tuber moth, *Phthorimaea operculella* Zell. *Agricultural Research Review* 68 (1): 147 - 156.
- Mehlhorn, H., Schmahl, G. and Haberkorn, A. (1988).** Toltrazuril effective against a broad spectrum of protozoan parasites. *Parasitology Research* 75 (1): 64 - 66.
- Mehrnejad, M.R., Batlle, I., Hormaza, I. and Espiau, M.T. (2002).** The natural parasitism ratio of the pistachio twig borer moth, *Kermania pistaciella*, in Iran. *Proceedings of the 3rd International Symposium on Pistachios and Almonds, 12th Colloquium of GREMPA, Zaragoza, Spain, 20-24 May, 2001. Acta Horticulturae* 591: 541 - 544.
- Mehto, D.N., Singh, K.M. and Singh, R.N. (1986).** Natural enemy complex on insect pest complex in chickpea *Cicer arietinum* Linn. *Bulletin of Entomology New Delhi*, publ. 1989, 27 (1): 1 - 12.
- Meindl, P., Kromp, B., Bartl, B., Ioannidou, E., Borgemeister, C. and Poehling, H.M. (2001).** Arthropod natural enemies of the cereal leaf beetle (*Oulema melanopus* L.) in organic winter wheat fields in Vienna, Eastern Austria. *Proceedings of the IOBC-WPRS Working Group "Integrated Control in Cereal Crops" at Godollo, Hungary, 9-12 September, 1999. Bulletin OILB SROP* 24 (6): 79 - 86.
- Men, U.B. and Thakre, H.S. (1998).** Field parasitization of sunflower semilooper, *Thysanoplusia orichalcea* F. by various parasitoids. *PKV Research Journal* 22 (2): 219 - 221.
- Men, U.B. and Thakre, H.S. (1998).** Parasitoids, predators and pathogens recorded on sunflower semilooper, *Thysanoplusia orichalcea* F. in Vidarbha. *PKV Research Journal* 22 (2): 218 - 219.
- Menalled, F.D., Marino, P.C., Gage, S.H. and Landis, D.A. (1999).** Does agricultural landscape structure affect parasitism and parasitoid diversity? *Ecological Applications* 9 (2): 634 - 641.
- Mescheloff, E. and Rosen, D. (1993).** Biosystematic studies on the Aphidiidae of Israel (Hymenoptera: Ichneumonoidea). 5. The genera *Trioxys* and *Binodoxys*. *Israel Journal of Entomology* 27: 31 - 47.
- Miczulski, B. (1987).** Incidence of parasitic Hymenoptera in the population dynamics of the cereal leaf beetles, *Oulema* spp. (Coleoptera: Chrysomelidae). *Ekologia Polska* 35 (3-4): 741 - 754.

- Miczulski, B. (1988).** The occurrence of parasitic Hymenoptera (Hymenoptera, Parasitica) in fields of winter wheat and spring barley. *Roczniki Nauk Rolniczych, E Ochrona Roslin* 17 (1): 89 - 95.
- Miller, G.L. and Cave, R.D. (1987).** Bionomics of *Micromus posticus* (Walker) (Neuroptera: Hemerobiidae) with descriptions of the immature stages. *Proceedings of the Entomological Society of Washington* 89 (4): 776 - 789.
- Miller, R.H., El Masir, S. and Al Jundi, K. (1992).** Incidence of wheat stem sawflies and their natural enemies on wheat and barley in northern Syria. *Arab Journal of Plant Protection* 10 (1): 30 - 25.
- Mills, N.J. (1993).** Observations of the parasitoid complexes of budmoths (Lepidoptera: Tortricidae) on larch in Europe. *Bulletin of Entomological Research* 83 (1): 103 - 112.
- Mills, N.J. and Kenis, M. (1991).** A study of the parasitoid complex of the European fir budworm, *Choristoneura murinana* (Lepidoptera: Tortricidae), and its relevance for biological control of related hosts. *Bulletin of Entomological Research* 81 (4): 429 - 436.
- Minami, T., Ishii, M. and Temma, K. (1999).** Difference in parasitoid complex of gypsy moth, *Lymantria dispar* L. (Lepidoptera: Lymantriidae), between mountain coppice and urban greenery in Osaka, Japan. *Japanese Journal of Applied Entomology and Zoology* 43 (4): 169 - 174.
- Mirchev, P., Georgiev, G.T. and Tsankov, G. (2001).** Studies on the parasitoids of *Gelechia senticetella* (Stgr.) (Lepidoptera: Gelechiidae) in Bulgaria. *Anzeiger für Schadlingskunde* 74 (4): 94 - 96.
- Mishra, A.K. and Shrivastava, S.K. (2000).** Natural enemy complex of *Helicoverpa armigera* Hubn. in Durg. *Agricultural Science Digest* 20 (1): 17 - 20.
- Mishra, B.K., Mandal, S.M.A. and Tunga, T.K. (1992).** Seasonal activity of parasitoids of *Helicoverpa armigera* Hubner in the Eastern Ghat High Land Zone of Orissa. *Orissa Journal of Agricultural Research* 5 (3-4): 170 - 173.
- Mitchell, E.R., Hu, G.Y., Okine, J.S. and Carpenter, J.E. (1999).** Parasitism of diamondback moth (Lepidoptera: Plutellidae) larvae by *Cotesia plutellae* (Hymenoptera: Braconidae) and *Diadegma insulare* (Hymenoptera: Ichneumonidae) in cabbage fields after inundative releases of *C. plutellae*. *Journal of Entomological Science* 34 (1): 101 - 112.
- Mitchell, E.R., Hu, G.Y. and Okine, J.S. (1997).** Diamondback moth (Lepidoptera: Plutellidae) infestation and parasitism by *Diadegma insulare* (Hymenoptera: Ichneumonidae) in collards and adjacent cabbage fields. *Florida Entomologist* 80 (1): 54 - 71.
- Mizell, R.F. and Schiffhauer, D.E. (1991).** Biology, effect on hosts, and control of the azalea leafminer (Lepidoptera: Gracillariidae) on nursery stock. *Environmental Entomology* 20 (2): 597 - 602.
- Modarres Awal, M. (1997).** List of Agricultural Pests and their Natural Enemies in Iran. Ferdowsi University Press, 429 pp (P. 275).

- Mohammadi, M. (1993).** Complementary investigation on the biology of *Rhyacionia buoliana* Schiff. (Lep.: Tortricidae) in northern part of Iran. Proc. XI. Plant Protec. Cong. Iran; Univ. Gilan, P. 241.
- Mohammed, I.M., Badrul, A.B., Anowar, C. and Chowdhury, A. (1999).** A preliminary list of the parasitic wasps (Hymenoptera: Ichneumonidae) of Bangladesh. Bangladesh Journal of Entomology 9 (1-2): 61 - 67.
- Mohanadas, K. (1996).** New records of some natural enemies of the teak defoliator, *Hyblaea puera* Cramer (Lepidoptera: Hyblaeidae) from Kerala, India. Entomon 21 (3-4): 251 - 253.
- Molina-Ochoa, J., Hamm, J.J., Lezama-Gutierrez, R., Lopez-Edwards, M., Gonzalez-Ramirez, M. and Pescador-Rubio, A. (2001).** A survey of fall armyworm (Lepidoptera: Noctuidae) parasitoids in the Mexican states of Michoacan, Colima, Jalisco, and Tamaulipas. Armyworm Symposium 2000. Florida Entomologist 84 (1): 31 - 36.
- Moller, K. (2002).** Ichneumonidae as important parasitoids of *Panolis flammea* Schiff. (Hym., Lep., Noctuidae). Entomologische Nachrichten und Berichte 46 (3): 191 - 192.
- Momoi, S. (1993).** Ergebnisse der zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei. 331. Einige mongolischen Arten der Unterfamilien Ephialtinae und Xoridinae (Hymenoptera: Ichneumonidae). Folia Entomologica Hungarica 26 (Suppl.): 241-250.
- Monajemi, N. (1975).** Contribution a l'étude des hymenopteres parasites de *Hypera postica* (Col.: Curculionidae) en Iran. J. Entomol. Soc. Iran 2 (2): 85-99.
- Monnerat, R.G. and Bordat, D. (1998).** Influence of HD-1 *Bacillus thuringiensis* spp. *kurstaki* on the developmental stages of *Diadegma* sp. (Hym.: Ichneumonidae) parasitoid of diamondback moth (Lep.: Yponomeutidae). J. Appl. Entomol. 122: 49-51.
- Monnerat, R.G., Kirk, A.A. and Bordat, D. (2002).** Biology of *Diadegma* sp. (Hymenoptera: Ichneumonidae), a parasitoid of *Plutella xylostella* (L.) (Lepidoptera: Yponomeutidae), from Reunion Island. Neotropical Entomology 31 (2): 271 - 274.
- Moore, S. and Kfir, R. (1995).** Host preference of the facultative hyperparasitoid *Tetrastichus howardi* (Hym.: Eulophidae). Entomophaga 40 (1): 69 - 76.
- Moore, S.D. and Kfir, R. (1996).** Biological studies of *Xanthopimpla stemmator* (Thunberg) (Hymenoptera: Ichneumonidae), a parasitoid of lepidopteran stem borers. African Entomology 4 (2): 131 - 136.
- Moraes, R.R. Loeck, A.E. and Belarmino, L.C. (1991).** Natural enemies of *Rachiplusia nu* (Guenee, 1852) and *Pseudoplusia includens* (Walker, 1857) (Lepidoptera: Noctuidae) on soyabean in Rio Grande do Sul. Pesquisa Agropecuaria Brasileira 26 (1): 57 - 64.
- Morena, J., Falco, J.V. and Jimenez, R. (1990).** Study of the parasitoid complex of *Prays citri* Mill. (Lep.: Hyponomeutidae) in the south of the Alicante Province. Boletin de Sanidad Vegetal, Plagas 16 (2): 515 - 522.
- Mori, P. (1988).** Parasitoids of insect and mite pests. Vita in Campagna 6 (2): 26 - 27.
- Morris, M.J., Roberts, S.J., Maddox, J.V. and Armbrust, E.J. (1996).** Epizootiology of the fungal pathogen, *Zoophthora phytonomi* (Zygomycetes: Entomophthorales) in field

- populations of alfalfa weevil (Coleoptera: Curculionidae) larvae in Illinois. *Great Lakes Entomologist* 29 (3): 129 - 140.
- Morse, D.H. (1994).** The effect of host size on sex ratio in the ichneumonid wasp, *Trychosis cyperia*, a spider parasitoid. *American Midland Naturalist* 131 (2): 281 - 287.
- Mowat, D.J. and Clawson, S. (1995).** Natural control of the small ermine moth *Yponomeuta padella* (L.). *Agriculture, Ecosystems and Environment* 52 (2-3): 93 - 102.
- Muckenfuss, A.E. and Shepard, B.M. (1994).** Seasonal abundance and response of diamondback moth, *Plutella xylostella* (L.) (Lepidoptera: Plutellidae), and natural enemies to esfenvalerate and *Bacillus thuringiensis* subsp. *kurstaki* Berliner in coastal South Carolina. *Journal of Agricultural Entomology* 11 (4): 361 - 373.
- Mukunthan, N. (1989).** Life tables of sugar-cane top borer, *Scirpophaga excerptalis* Wlk. *Insect Science and its Application* 10 (3): 269 - 276.
- Munoz, L.E., Serrano, P.A., Pulido, J.I. and De La Cruz, L.J. (1994).** Life cycle, behaviour and natural enemies of *Neoleucinodes elegantalis* (Guenee, 1854) (Lepidoptera: Pyralidae) fruit borer of *Solanum quitoense* Lam. in Valle del Cauca. *Acta Agronomica, Universidad Nacional de Colombia* 1991, reed. 41 (1-4): 99 - 104.
- Munro, V.M.W. (1998).** A record of the releases and recoveries of the Australian parasitoids *Xanthopimpla rhopaloceros* Krieger (Hymenoptera: Ichneumonidae) and *Trigonospila brevifacies* Hardy (Diptera: Tachinidae) introduced into New Zealand for leafroller control. *New Zealand Entomologist* 21: 81 - 91.
- Munro, V.M.W. (1998).** A retrospective analysis of the establishment and dispersal of the introduced Australian parasitoids *Xanthopimpla rhopaloceros* (Krieger) (Hymenoptera: Ichneumonidae) and *Trigonospila brevifacies* (Hardy) (Diptera: Tachinidae) within New Zealand. *Biocontrol Science and Technology* 8 (4): 559 - 571.
- Munro, V.M.W. and Henderson, I.M. (2002).** Nontarget effect of entomophagous biocontrol: shared parasitism between native lepidopteran parasitoids and the biocontrol agent *Trigonospila brevifacies* (Diptera: Tachinidae) in forest habitats. *Environmental Entomology* 31 (2): 388 - 396.
- Munster-Swendsen, M. (1989).** Phenology and natural mortalities of the fir needleminer, *Epinotia fraternana* (Hw.) (Lepidoptera: Tortricidae). *Entomologiske Meddelelser* 57 (3): 111 - 120.
- Munster-Swendsen, M. (1994).** Pseudoparasitism: detection and ecological significance in *Epinotia tedella* (Cl.) (Tortricidae). *Norwegian Journal of Agricultural Sciences SUPP16*: 329 - 335.
- Muraleedharan, N., Selvasundaram, R. and Radhakrishnan, B. (1988).** Natural enemies of certain tea pests occurring in southern India. *Insect Science and its Application* 9 (5): 647 - 654.
- Muraleedharan, N. and Selvasundaram, R. (1991).** Bio-ecology of *Phytodietus spinipes* (Cameron) a parasitoid of *Homona coffearia* Nietner, the tea tortrix. *Journal of Plantation Crops* 19 (1): 26 - 32.

- Murchie, A.K., Williams, I.H., Paul, V.H., Kohr, K., Thomas, J. and Dupprich, P.D. (1998).** A bibliography of the parasitoids of the cabbage seed weevil (*Ceutorhynchus assimilis* PAYK.). Integrated Control in Oilseed Crops. Proceedings of a meeting held at Poznan, Poland, 10-12, April 1997. Bulletin OILB SROP 21 (5): 163 - 169.
- Murray, D.A., Kriegel, R.D., Johnson, J.W. and Howitt, A.J. (1996).** Natural enemies of cranberry fruitworm, *Acrobasis vaccinii*, (Lepidoptera: Pyralidae) in Michigan highbush blueberries. Great Lakes Entomologist 29 (2): 81 - 86.
- Murray, D.A.H., Monsour, C.J., Teakle, R.E., Rynne, K.P. and Bean, J.A. (1995).** Interactions between nuclear polyhedrosis virus and three larval parasitoids of *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae). Journal of the Australian Entomological Society 34 (4): 319 - 322.
- Murray, D.A.H., Rynne, K.P., Winterton, S.L., Bean, J.A. and Lloyd, R.J. (1995).** Effect of host plant on parasitism of *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) by *Hyposoter didymator* Thunberg (Hymenoptera: Ichneumonidae) and *Cotesia kazak* (Telenga) (Hymenoptera: Braconidae). Journal of the Australian Entomological Society 34 (1): 71 - 73.
- Murugan, K., Kumar, N.S., Jeyabalan, D., Nathan, S.S., Sivaramakrishnan, S. and Swamiappan, M. (2000).** Influence of *Helicoverpa armigera* (Hubner) diet on its parasitoid *Campoletis chloridae* Uchida. Insect Science and its Application 20 (1): 23 - 31.
- Mushtaque, M. (1989).** Control of large cabbage butterfly in Sialkot by local parasites. Progressive Farming 9 (4): 38 - 40.
- Mushtaque, M., Irshad, M. and Mohyuddin, A.I. (1993).** Studies on mortality factors in immature stages of *Pieris brassicae* (L.) (Pieridae: Lepidoptera) and effect of Bactospeine on its larvae and parasites. Pakistan Journal of Zoology 25 (2): 165 - 167.
- Mustata, G. (2001).** The role of parasitoid and hyperparasitoid complex in the population of diamond back moth in Romania. Analele Stiintifice ale Universitatii 'Al I Cuza' din Iasi. Serie Noua Sectiunea I Biologie Animala 47: 7 - 15.

## N

- Nabli, H., Bailey, W.C. and Necibi, S. (1999).** Beneficial insect attraction to light traps with different wavelengths. Biological Control 16 (2): 185 - 188.
- Nagalingam, B. (1988).** Studies on the biology of *Psorosticha zizyphi* Stn. on acid lime (*Citrus aurantifolia* Swingle). Mysore Journal of Agricultural Sciences 22 (1): 48 - 52.
- Nagalingam, B. (1991).** Hymenopterous parasitoids of citrus leaf roller. Current- Research - University of Agricultural Sciences Bangalore 20 (7): 139 - 140.
- Naganagoud, A. and Kulkarni, K.A. (1997).** Surveillance of sorghum armyworm *Mythimna separata* (Walker) and its natural enemies in transitional region of Dharwad. Journal of Biological Control, publ. 1998, 11 (1-2): 65 - 68.

- Nakahara, Y., Hiraoka, T. and Iwabuchi, K. (1999).** Effects of lipophorin and 20-hydroxyecdysone on in vitro development of the larval endoparasitoid *Venturia canescens* (Hymenoptera: Ichneumonidae). *Journal of Insect Physiology* 45 (5): 453 - 460.
- Nakahara, Y., Hiraoka, T. and Iwabuchi, K. (2000).** Growth-promoting effects of ecdysteroids and juvenile hormone on in vitro development of the larval endoparasitoid, *Venturia canescens* (Hymenoptera: Ichneumonidae). *Journal of Insect Physiology* 46 (4): 467 - 476.
- Nakahara, Y., Iwabuchi, K. and Mitsuhashi, J. (1997).** In vitro rearing of the larval endoparasitoid, *Venturia canescens* (Gravenhorst) (Hymenoptera: Ichneumonidae). III Growth promoting ability of extract of *Galleria mellonella* pupae. *Applied Entomology and Zoology* 32 (1): 91 - 99.
- Nakahara, Y., Nakamura, M., Hiraoka, T. and Iwabuchi, K. (2002).** Insect lipophorin and vertebrate lipoproteins support larval development of the endoparasitoid *Venturia canescens* (Hymenoptera: Ichneumonidae) as dietary lipid sources. *Applied Entomology and Zoology* 37 (4): 637 - 643.
- Nakahara, Y. and Iwabuchi, K. (2000).** In vitro rearing of the larval endoparasitoid, *Venturia canescens* (Gravenhorst) (Hymenoptera: Ichneumonidae). IV. Effects of osmotic pressure and pH on larval development. *Applied Entomology and Zoology* 35 (3): 381 - 388.
- Nakahara, Y. and Iwabuchi, K. (2000).** Investigation of low thermal threshold for development of the larval endoparasitoid, *Venturia canescens* (Hymenoptera: Ichneumonidae) using in vitro culture technique. *Entomological Science* 3 (1): 19 - 23.
- Nakanishi, A. (1978).** A new genus and two new species of the Diplazontinae (Hymenoptera: Ichneumonidae) from Japan. *Kontyu* 46: 429-432.
- Nandihalli, B.S. and Lee, J.H. (1995).** Effect of host food plants on the biology of the host, *Helicoverpa assulta* (Guenee), and its parasitoid, *Campoletis chlorideae* Uchida. *Advances in Agricultural Research in India* 3: 22 - 32.
- Nandihalli, B.S. and Lee, J.H. (1995).** Seasonal occurrence of *Campoletis chlorideae* Uchida and its control efficacy on the oriental tobacco budworm, *Helicoverpa assulta* (Guenee), in tobacco fields in Suwon. *Korean Journal of Applied Entomology* 34 (2): 147 - 153.
- Nanni, C. (1991).** Activity of indigenous entomophagous species against *Hyphantria cunea*. *Informatore Fitopatologico* 41 (7-8): 15 - 21.
- Napiorkowska, J. (1991).** Parasitic insects reared from larvae and pupae of *Archips rosanus* L. (Lepidoptera: Tortricidae) collected on ornamental shrubs of Lublin. *Wiadomosci Entomologiczne* 10 (3): 163 - 167.
- Napiorkowska, J. and Filipowich, A. (1995).** The ichneumon wasp *Exetastes atrator* Forster (Hymenoptera: Ichneumonidae) as a specific parasitoid of the cabbage moth - *Mamestra brassicae* (L.) (Lepidoptera: Noctuidae). *Materialy ogólnopolskiej konferencji naukowej*



- Nauka Praktyce Ogrodniczej z okazji XXV lecia Wydzialu Ogrodniczego Akademii Rolniczej w Lublinie 399 - 402.
- Narendran, T.C. and Singh, J. (2000).** A new species of *Campoplex gravenhorst* (Hymenoptera: Ichneumonidae) from Sikkim, India. *Oriental Insects* 34: 289 - 292.
- Naresh, J.S., Malik, V.S., Balan, J.S. and Khokhar, K.S. (1986).** A new record of *Trathala* sp., a larval endoparasite attacking brinjal fruit borer, *Leucinodes orbonalis* Guenee. *Bulletin of Entomology New Delhi*, publ. 1989, 27 (1): 74.
- Naresh, J.S., Malik, V.S. and Balan, J.S. (1986).** Estimation of fruit damage and larval population of brinjal fruit borer, *Leucinodes orbonalis* Guenee and its parasitism by *Trathala* sp. on brinjal. *Bulletin of Entomology New Delhi*, publ. 1989, 27 (1): 44 - 47.
- Narolsky, N.B. (1990).** New data on the cremastine wasps (Hymenoptera: Ichneumonidae) from Kazackstan and Central Asia. (in Russian) *Zoologicheskii Zhurnal* 7: 63-68.
- Narolsky, N.B. (2002).** A review of some type specimens of the Cremastinae (Hymenoptera: Ichneumonidae) wasps described by G. Szepliget in the collection of the Hungarian National Museum. *Vestnik Zoologii* 36 (2): 23 - 30.
- Narolsky, N.B. (2002).** *Tersoakus* gen. nov., a new genus of cremastine wasps from the Russian Far East (Hymenoptera: Ichneumonidae: Cremastinae). *Zoologische Mededelingen* 76: 1 - 16, 97 - 102.
- Nath, P. and Rai, R. (1999).** Study of key mortality factors in the population dynamics of chickpea pod borer, *Helicoverpa armigera* (Hubner) (Noctuidae: Lepidoptera) infesting chickpea, *Cicer arietinum* L. *Tropical Ecology* 40 (2): 281 - 288.
- Ndungu, A.W. (1994).** New natural enemies of the coffee berry moth, *Prophantis smaragdina* (Butler) in Kenya. *Kenya Coffee*. 59: 688, 1725 - 1726.
- Nealis, V.G., Roden, P.M. and Ortiz, D.A. (1999).** Natural mortality of the gypsy moth along a gradient of infestation. *Canadian Entomologist* 131 (4): 507 - 519.
- Nealis, V.G. and Lysyk, T.J. (1980).** Sampling overwintering jack pine budworm, *Choristoneura pinus pinus* Free. (Lepidoptera: Tortricidae), and two of its parasitoids (Hymenoptera). *Canadian Entomologist* 120 (12): 1101 - 1111.
- Neel, C.S., Weaver, J.E., Baniecki, J.F. and Wallbrown, R.M. (1992).** A survey of biological agents of the alfalfa weevil in Mason and Preston Counties, West Virginia, for 1992. *Circular Agricultural and Forestry Experiment Station, West Virginia University* 158: 11 pp.
- Neira, C.M., Ruff, F.J. and Mundaca, B.N. (1989).** Natural enemies of *Pieris brassicae* L. (Lepidoptera: Pieridae) from cultivated crucifers in Valdivia. *Agro Ciencia* 5 (2): 5 - 10.
- Nelson, J.M. (1991).** The biology and early stages of the fly *Paraprosalpia sepiella* (Zetterstedt) (Diptera: Anthomyiidae). *Entomologist's Gazette* 42 (3): 185 - 187.
- Nemoto, T., Shibuya, M., Kuwahara, Y. and Suzuki, T. (1987).** New 2-acylcyclohexane-1,3-diones: kairomone components against a parasitic wasp, *Venturia canescens*, from feces of the almond moth, *Cadra cautella*, the Indian meal moth, *Plodia interpunctella*. *Agricultural and Biological Chemistry* 51 (7): 1805 - 1810.

- Nenon, J. (1993).** The *Megarhyssa* (Hymenoptera: Ichneumonidae) in the south of Quebec; simplified identification key and flight period. *Faberies* 18 (2): 49 - 58.
- Nenon, J.P. (1995).** Bioecology of *Megarhyssa atrata* (Hymenoptera: Ichneumonidae), an ectoparasitoid of *Tremex columba* (Hymenoptera: Siricidae) in Quebec. *Phytoprotection* 76 (3): 115 - 122.
- Nenon, J.P., Le Lannic, J., Kacem, N., Barbier, R. and Allo, M.R. (1995).** Micromorphology of the ovipositor in Hymenoptera and evolution from phytophagous Symphyta to parasitoids Apocrita. *Comptes Rendus de l'Academie des Sciences. Serie III, Sciences de la Vie* 318 (10): 1045 - 1051.
- Newton, B.L. and Sharkey, M.J. (2000).** A new species of *Bassus* (Hymenoptera: Braconidae: Agathidinae) parasitic on *Samea multiplicalis*, a natural control agent of waterlettuce. *Florida Entomologist* 83 (3): 284 - 289.
- Ngamo-Tinkeu, L.S., Delabye, G., Stilmant, D. and Hance, T. (1998).** Phenology of parasitism and its influence on the predatory efficiency of larvae of *Episyrphus balteatus* (Diptera: Syrphidae). Integrated control in cereal crops. Meeting held at Lleida, Spain, 13-14 March, 1997. *Bulletin OILB SROP* 21 (8): 133 - 139.
- Nielsen, G.C. and Holm, S. (1988).** The meadow quick grass gallmidge in the field in 1988. *Dansk Froavl* 71: 22, 400 - 402.
- Nierhaus-Wunderwald, D. (1998).** Biology and natural regulation of the Yponomeutidae. *Notice pour le Praticien* 29: 8 pp.
- Nikam, P.K. (1979).** A new genus and species of Anomaloninae (Hymenoptera: Ichneumonidae) from India. *Entomon* 7: 115-118.
- Nikam, P.K. (1990).** Ichneumonidae of Indian *Heliopsis* Ochsenheimer. *Journal of Entomological Research* 14 (1): 52 - 59.
- Nikam, P.K. and Gaikwad, A.M. (1989).** Role of hymenopterous parasitoids in the biological control of *Heliopsis armigera* (Hubn.) (Lepidoptera: Noctuidae) with special reference to *Campoletis chloridae* Uchida (Hymenoptera: Ichneumonidae) in India. *Journal of Entomological Research* 13 (1-2): 6 - 20.
- Nikam, P.K. and Gaikwad, A.M. (1991).** Effect of host larvae of *Helicoverpa armigera* Hubner on the parasitising ability of *Campoletis chloridae* Uchida. *Entomon* 16 (4): 301 - 303.
- Nikdel, M., Sadaghian, B. and Dordaei, A.A. (2004).** Report of two Ichneumonid species as a parasitoid of born-tail moth in Arasbaran. *Proceedings of 16<sup>th</sup> Iranian Plant Protection Congress*, p. 102.
- Nilsson, C. (1988).** The pollen beetle (*Meligethes aeneus* F.) in winter and spring rape at Alnarp 1976-1978. III. Mortality factors. *Vaxtskyddsnotiser*, publ. 1989, 52 (6): 145 - 150.
- Nilsson, C. (1994).** Pollen beetles (*Meligethes* spp.) in oil seed rape crops (*Brassica napus* L.): biological interactions and crop losses. Swedish University of Agricultural Sciences, Alnarp; Sweden, 308 pp.

- Nilsson, C. (1994).** Pollen beetle and rape: interactions and crop losses. *Vaxtskyddsnotiser* 58 (2): 56 - 57.
- Nitzsche, O. and Ulber, B. (1998).** Influence of different tillage treatments following the harvest of oilseed-rape on the mortality of pollen beetle (*Meligethes* spp.) parasitoids. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 105 (4): 417 - 421.
- Niwa, C.G. (1988).** Parasites and predators associated with the ponderosa pine tip moth, *Rhyacionia zozana* (Kearfott) (Lepidoptera: Tortricidae), in California and Oregon. *Canadian Entomologist* 120 (10): 881 - 886.
- Niwa, C.G. and Daterman, G.E. (1989).** Pheromone mating disruption of *Rhyacionia zozana* (Lepidoptera: Tortricidae): Influence on the associated parasite complex. *Environmental Entomology* 18 (4): 570 - 574.
- Niyibigira, E.I., Abdallah, Z.S., Overholt, W.A., Lada, V.Y. and Van Huis, A. (2001).** Distribution and abundance, in maize and sorghum, of lepidopteran stemborers and associated indigenous parasitoids in Zanzibar. *Insect Science and its Application* 21 (4): 335 - 346.
- Noda, T. (2000).** Detection of diploid males and estimation of sex determination system in the parasitic wasp *Diadegma semiclausum* (Hellen) (Hymenoptera: Ichneumonidae) using an allozyme as a genetic marker. *Applied Entomology and Zoology* 35 (1): 41 - 44.
- Noda, T., Miyai, S., Yamada, S. and Konishi, K. (1996).** Larval and pupal parasitoids of diamondback moth, *Plutella xylostella* (L.) in cabbage fields in Morioka, Japan. *Japanese Journal of Applied Entomology and Zoology* 40 (2): 164 - 167.
- Noda, T., Miyai, S.I., Takashino, K. and Nakamura, A. (2000).** Density suppression of *Plutella xylostella* (Lepidoptera: Yponomeutidae) by multiple releases of *Diadegma semiclausum* (Hymenoptera: Ichneumonidae) in cabbage fields in Iwate, northern Japan. *Applied Entomology and Zoology* 35 (4): 557 - 563.
- Norten, E., Linde, A., Happe, M. and Madel, G. (1988).** Infection and pathology of a microsporidiosis of *Diadegma semiclausum* Hellen (Hym.: Ichneumonidae). *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 6 (1-3): 284 - 285.
- Notton, D.G. and Shaw, M.R. (1998).** A review of the Palaearctic Neorhacodinae (Hymenoptera: Ichneumonidae) with *Eremura* Kasparyan, 1995 new to the west Palaearctic. *Bulletin of the British Museum Natural History, Entomology Series* 67 (2): 209 - 218.
- Noyes, J.S. (1989).** The diversity of Hymenoptera in the tropics with special reference to Parasitica in Sulawesi. *Ecological Entomology* 14 (2): 197 - 207.
- Nuorteva, P., Tulisalo, E., Li, T.S. and Hong, P.Z. (1999).** Metal pollution in relation to mass outbreaks of *Dendrolimus punctatus* Walker on *Pinus massoniana* Lamb. in China. *Journal of Environmental Sciences* 11 (4): 498 - 503.
- Nurindah, A.A.A.G. and Indrayani, I.G.A.A. (1989).** New records of natural enemies of *Heliothis armigera* on cotton in Indonesia. *Industrial Crops Research Journal* 2 (1): 1 - 5.

- Nurullahoglu, Z.U. and Aksoylar, M.Y. (1997).** The effects of low temperature on the total lipid, total fatty acid and fatty acid composition in female pupae and adults of *Pimpla turionellae* L. (Hymenoptera: Ichneumonidae). Turkish Journal of Zoology 21 (3): 295 - 301.
- Nwanze, K.F. and Nwilene, FE. (1998).** Interactions of host plant resistance and biological control of stemborers in sorghum. Insect Science and its Application 18 (3): 261 - 266.
- Nwilene, F.E., Nwanze, K.F. and Reddy, Y.V.R. (1998).** Effect of sorghum ecosystem diversification and sowing date on shoot fly, stem borer and associated parasitoids. Crop Research Hisar 16 (2): 239 - 245.

## O

- Obukhov, M.L. (1993).** Baculiform viruses. Biology Bulletin of the Russian Academy of Sciences 19 (3): 205 - 213.
- Ochoa, R., Carballo, M., Rutilio-Quezada, J. and Quezada, JR. (1989).** Some aspects of the biology and behaviour of *Plutella xylostella* (Lepidoptera: Plutellidae) and its parasitoid *Diadegma insulare* (Hymenoptera: Ichneumonidae). Manejo Integrado de Plagas 11: 21 - 30.
- O'Connor, J.P. and Shaw, M.R. (2004).** *Afrephialtes cicatricose* (Ratzeburg) (Hymenoptera: Ichneumonidae, Pimplinae), an unexpected addition to the Irish fauna. Entomologist's Gazette 55: 119-120.
- Oh, M.R., Kim, S.S., Park, JD., Paik, JC. and Kim, DI. (1997).** Biological characteristics of *Cotesia plutellae* (Hymenoptera: Braconidae), a larval parasitoid of *Plutella xylostella* (Lepidoptera: Yponomeutidae). Korean Journal of Entomology 27 (3): 79 - 84.
- Ohbayashi, T., Iwabuchi, K. and Mitsuhashi, J. (1994).** In vitro rearing of a larval endoparasitoid, *Venturia canescens* (Gravenhorst) (Hymenoptera: Ichneumonidae). I. Embryonic development. Applied Entomology and Zoology 29 (1): 123 - 126.
- Okada, T. (1989).** Parasitoids of the diamondback moth, *Plutella xylostella* (L.) (Lepidoptera: Yponomeutidae): species and seasonal changes of parasitism in cabbage fields. Japanese Journal of Applied Entomology and Zoology 33 (1): 17 - 23.
- Okada, T. (1991).** Seasonal changes in population density of the diamondback moth, *Plutella xylostella* (L.) and parasitism by parasitoids in cabbage fields. Proceedings of the Kansai Plant Protection Society 33: 39 - 44.
- Okine, J.S., Mitchell, E.R. and Hu, G.Y. (1996).** Low temperature effect on viability of *Diadegma insulare* (Hymenoptera: Ichneumonidae) pupae and effect of this parasitoid on feeding rate of diamondback moth larvae (Lepidoptera: Plutellidae). Florida Entomologist 79 (4): 503 - 509.
- Olckers, T. and Hulley, P.E. (1991).** Notes on some insect galls associated with Solanum plants in South Africa. South African Journal of Zoology 26 (2): 55 - 61.
- Olofsson, E. (1994).** Biology and outbreaks of *Microdiprion pallipes* (Hymenoptera: Diprionidae) in Sweden. Studia Forestalia Suecica 193: 20 pp.

- Oloo, G.W. and Ogeda, K. (1990).** The incidence of *Chilo partellus* (Swinh.) (Pyralidae) and the contribution of natural enemies to its mortality under intercropping system in Kenya. *Tropical Pest Management* 36 (3): 244 - 248.
- Oloumi-Sadeghi, H., Steffey, K.L., Roberts, S.J., Maddox, J.V. and Armbrust, E.J. (1993).** Distribution and abundance of two alfalfa weevil (Coleoptera: Curculionidae) larval parasitoids in Illinois. *Environmental Entomology* 22 (1): 220 - 225.
- Omata, K. (1989).** Control mechanisms of larval diapause and quiescence in *Trogus mactator*, an endoparasite of papilionids. *Entomologia Experimentalis et Applicata* 53 (1): 31 - 37.
- O'Neill, K.M., Kemp, W.P., Seibert, C., Rolston, M.G., Bess, J.A. and Philips, T.K. (2001).** Natural enemy assemblages on native and reseeded grasslands in Southwestern Montana: a family-level analysis. *Western North American Naturalist* 61: 21, 195 - 203.
- Ooi, P.A.C., Kadir, A.A.S.A. and Barlow, H.S. (1992).** Trends in biological control in Malaysia. *Pest management and the environment in 2000*. 341 - 347.
- Ortega, G. (1989).** The genus *Syzeuctus* Foerster, 1868 in the Canary Islands, with the description of a new species: *S. baezi* n. sp. (Ichneumonidae: Banchinae). *Boletín de la Asociación Española de Entomología* 13: 269 - 276.
- Ortel, J. (1991).** Effects of lead and cadmium on chemical composition and total water content of the pupal parasitoid, *Pimpla turionellae*. *Entomologia Experimentalis et Applicata* 59 (1): 93 - 100.
- Ortel, J. (1995).** Accumulation of Cd and Pb in successive stages of *Galleria mellonella* and metal transfer to the pupal parasitoid *Pimpla turionellae*. *Entomologia Experimentalis et Applicata* 77 (1): 89 - 97.
- Ortel, J. and Vogel, W.R. (1989).** Effects of lead and cadmium on oxygen consumption and life expectancy of the pupal parasitoid, *Pimpla turionellae*. *Entomologia Experimentalis et Applicata* 52 (1): 83 - 88.
- Ortiz, F.M., Brenes-Blanco, J. and Perez-Jiron, H. (1999).** Rearing of the parasitoids *Cotesia plutellae*, *Microplitis plutellae* and *Diadegma insulare* under laboratory conditions. II part. *Manejo Integrado de Plagas* 52: 100 - 102.
- Ortu, S. and Floris, I. (1989).** Preliminary study on the control of *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae) on potato crops in Sardinia. *Difesa- delle Piante*. 12 (1-2): 81-88; Paper presented at the conference 'Pests and diseases of horticultural field crops and methods of their control' held in Siracusa, Sicily, on 22 - 24 February 1988.
- Otten, H., Wackers, F., Battini, M. and Dorn, S. (2001).** Efficiency of vibrational sounding in the parasitoid *Pimpla turionellae* is affected by female size. *Animal Behaviour* 61 (3): 671 - 677.
- Otten, H., Wackers, F.L., Dorn, S., Sommeijer, M.J. and Francke, P.J. (1998).** Variation in vibrations produced by *Pimpla turionellae* (Hymenoptera: Ichneumonidae). *Proceedings of the 9th Meeting of Experimental and Applied Entomologists, Leiden, Netherlands, 19*

- December 1997. Proceedings of the Section Experimental and Applied Entomology of the Netherlands Entomological Society 9: 73 - 74.
- Overgaard, S.M. (1989).** Some parasitic wasps reared from Danish burnet moths (Lepidoptera: Zygaenidae: *Zygaena* Fabr. 1775, Hymenoptera: Braconidae, Ichneumonidae, Pteromalidae and Eupelmidae). Entomologiske Meddelelser 57 (3): 143 - 149.
- Ozalp, P. and Emre, I. (1992).** Effects of water-soluble vitamins on egg production and hatchability of adult *Pimpla turionellae* L. Doga, Turk Zooloji Dergisi 16 (1): 78 - 83.
- Ozalp, P. and Emre, I. (2001).** The effects of carbohydrates upon the survival and reproduction of adult female *Pimpla turionellae* L. (Hym.: Ichneumonidae). Journal of Applied Entomology 125 (4): 177 - 180.
- Ozbek, H., Guclu, S. and Tozlu, G. (1999).** Biology and natural enemies of *Diplolepis mayri* Schld. (Hymenoptera: Cynipidae), a pest of *Rosa canina* L. in Erzurum province. Turkiye Entomoloji Dergisi 23 (1): 39 - 50.
- Ozbek, H., Pekel, S. and Kolarov, J. (2000).** New distributional data of the Turkish Ichneumonidae (Hymenoptera) II. Ctenopelmatinae and Campopleginae. Journal of the Entomological Research Society 2 (1):17 - 23.
- Ozdemir, Y. (1994).** Taxonomic studies on ichneumonid parasitoids of species of yponomeutids in Central Anatolia. Turkiye III. Biyolojik Mucadele Kongresi Bildirileeri, 25 - 28 Ocak 1994, Ege Universitesi Ziraat Fakultesi, Bitki Koruma Bolumu, Izmir 101 - 108.
- Ozdemir, Y. (1996).** Banchinae and Ichneumoninae (Hym.: Ichneumonidae) in Central Anatolia. Bitki Koruma Bulteni 36 (3-4): 91 - 104.
- Ozdemir, Y. (1996).** Taxonomic work on the Banchinae and Ichneumoninae (Hym.: Ichneumonidae) in the Central Anatolian region. Ziraat Mucadele Arastirma Yilligi 28 - 29: 223 - 224.
- Ozdemir, Y. and Kilincer, N. (1990).** Species of Pimplinae and Ophioninae (Hym.: Ichneumonidae) from Central Anatolia. Proceedings of the Second Turkish National Congress of Biological Control 309 - 318.
- Ozkan, A. and Yanikolu, A. (1999).** Effects of 2,4-D and maleic hydrazide on the glycogen level in the embryonic development of *Pimpla turionellae* (L.) (Hym.: Ichneumonidae). Journal of Applied Entomology 123 (4): 211 - 216.
- Ozkan, F. and Emre, I. (1997).** Effects of orally administrated malathion, an organophosphate insecticide, on longevity, egg production and hatchability of adult female *Pimpla turionellae* L. Turkish Journal of Zoology 21 (3): 309 - 313.
- Ozlu, A. (1993).** Change in density of the neurosecretory material in the median neurosecretory cells of *Pimpla turionellae* L. (Hymenoptera: Ichneumonidae) during the oviposition period. Doga, Turk Zooloji Dergisi 17 (3): 319 - 324.

## P

- Pagliano, G. and Scaramozzino, P. 1990 (1989).** Elenco dei generi di Hymenoptera del mondo. Memorie della Società Entomologica Italiana 122: 1-210.
- Pair, S.D. (1995).** Biology and rearing of *Diapetimorpha introita* (Cresson) (Hymenoptera: Ichneumonidae) on host and non-host noctuid pupae. Journal of Entomological Science 30 (4): 468 - 480.
- Pair, S.D. and Gross, H.R. (1989).** Seasonal incidence of fall armyworm (Lepidoptera: Noctuidae) pupal parasitism in corn by *Diapetimorpha introita* and *Cryptus albitarsis* (Hymenoptera: Ichneumonidae). Journal of Entomological Science 24 (3): 339 - 343.
- Palacio, E. (1999).** A curious case of parasitoids of the genus *Acrotaphus* (Hymenoptera: Ichneumonidae) in a nest of *Trigonopsis violascens* (Dalla Torre) (Hymenoptera: Sphecidae) [in Colombia]. Entomologo 27: 88, 7 - 8.
- Pandey, K.C. and Faruqi, S.A. (1987).** Observations on parasitization of lucerne weevil, *Hypera postica* Gyllenhal. Indian Journal of Entomology, publ. 1989, 49 (4): 552 - 554.
- Pantaleoni, R.A. (1987).** Studies on the parasitoids (Hym.: Ichneumonidae et Pteromalidae) of *Chrysoperla carnea* (St.) (Neur.: Chrysopidae) that search for and attack the stages enclosed in a cocoon. Bollettino dell'Istituto di Entomologia della Università degli Studi di Bologna 41: 241 - 255.
- Parakhin, G. (1991).** *Ephialtes manifestator*, a threat to buprestids and cerambycids. Zashchita Rastenii 10: 34 - 35.
- Parkinson, N., Conyers, C. and Smith, I. (2002).** A venom protein from the endoparasitoid wasp *Pimpla hyondriaca* is similar to snake venom reprotolysin-type metalloproteases. Journal of Invertebrate Pathology 79 (2): 129 - 131.
- Parkinson, N., Richards, E.H., Conyers, C., Smith, I. and Edwards, J.P. (2002).** Analysis of venom constituents from the parasitoid wasp *Pimpla hypochondriaca* and cloning of a cDNA encoding a venom protein. Insect Biochemistry and Molecular Biology 32 (7): 729 - 735.
- Parkinson, N., Smith, I., Audsley, N. and Edwards, J.P. (2002).** Purification of pimplin, a paralytic heterodimeric polypeptide from venom of the parasitoid wasp *Pimpla hypochondriaca*, and cloning of the cDNA encoding one of the subunits. Insect Biochemistry and Molecular Biology 32 (12): 1769 - 1773.
- Parkinson, N., Smith, I., Weaver, R. and Edwards, J.P. (2001).** A new form of arthropod phenoloxidase is abundant in venom of the parasitoid wasp *Pimpla hypochondriaca*. Insect Biochemistry and Molecular Biology 31 (1): 57 - 63.
- Parkinson, N.M., Conyers, C.M., Keen, J.N., MacNicoll, A.D., Smith, I., and Weaver, R.J. (2003).** cDNAs encoding large venom proteins from the parasitoid wasp *Pimpla hypochondriaca* identified by random sequence analysis. Comparative Biochemistry and Physiology. C, Toxicology and Pharmacology 134 (4): 513 - 520.

- Parkinson, N.M. and Weaver, R.J. (1999).** Noxious components of venom from the pupa-specific parasitoid *Pimpla hypochondriaca*. *Journal of Invertebrate Pathology* 73 (1): 74 - 83.
- Parr, J.C., Pass, B.C. and Nordin, G.L. (1993).** Compatibility of *Zoophthora phytonomi* (Entomophthorales: Entomophthoraceae) and *Bathyplectes anurus* (Hymenoptera: Ichneumonidae) in Kentucky alfalfa fields. *Environmental Entomology* 22 (3): 674 - 678.
- Patel, A.A. and Bhalani, P.A. (1989).** A new record of *Diadegma* sp. as a larval parasite of the sesamum leafroller in Gujarat. *Indian Journal of Entomology*, publ. 1990, 51 (4): 474.
- Patel, A.G., Yadav, D.N. and Patel, R.C. (1988).** Effect of low temperature storage on *Campoletis chlorideae* Uchida (Hymenoptera: Ichneumonidae) an important endo-larval parasite of *Heliothis armigera* Hubner (Lepidoptera: Noctuidae). *Gujarat Agricultural University Research Journal* 14 (1): 79 - 80.
- Patel, A.G., Yadav, D.N. and Patel, R.C. (1991).** Natural enemies of *Agrotis* spp. from Gujarat. *Gujarat Agricultural University Research Journal* 17 (1): 130 - 131.
- Patel, H.M. and Yadav, D.N. (1991).** Occurrence of mottled emigrant *Catopsilia pyranthe* (Lepidoptera: Pieridae) and its natural enemies at Anand. *Indian Journal of Agricultural Sciences* 61 (10): 789 - 791.
- Patel, I.S. and Dodia, D.A. (1997).** First record of *Diplazon laetatoris* Fab. (Ichneumonidae: Hymenoptera) as a pupal parasite of syrphid fly, *Xanthogramma scutellarge* Fab. (Syrphidae: Diptera) in Gujarat State. *Gujarat Agricultural University Research Journal* 23 (1): 106 - 107.
- Patel, P.N. and Habib, M.E.M. (1993).** *Microcharops anticarsiae* (Hym.: Ichneumonidae), parasitoid of *Anticarsia gemmatalis* (Lep.: Noctuidae): host age preference, sex ratio effects and functional response. *Entomophaga* 38 (4): 511 - 517.
- Patel, P.N. and Habib, M.E.M. (1998).** Development of *Microcharops anticarsiae* Gupta (Hym.: Ichneumonidae), an endoparasitoid of the velvetbean caterpillar, *Anticarsia gemmatalis* (Lep.: Noctuidae) and morphology of its immature stages. *Journal of Applied Entomology* 122 (7): 369 - 373.
- Patil, R.P. and Nikam, P.K. (1993).** On a new species of *Theلودon* Townes (Hymenoptera: Ichneumonidae) in India. *Indian Journal of Parasitology* 17 (1): 67 - 70.
- Patil, R.P. and Nikam, P.K. (1995).** On a new species of *Diaglyptidea* Viereck (Hymenoptera: Ichneumonidae) from India. *Oriental Insects* 29: 381 - 384.
- Patil, S.U. and Naik, M.I. (1998).** Natural enemies of teak defoliator, *Hyblaea puera* Cramer and their seasonal incidence. *Indian Journal of Forestry* 21 (3): 253 - 255.
- Patnaik, H.P., Rath, L.K., Senapati, B. and Behera, P.K. (1991).** Incidence of *Helicoverpa armigera* Hub. on chickpea and its population phenology in north central plateau zone of Orissa. *Orissa Journal of Agricultural Research* 4 (3-4): 137 - 142.
- Pavenstadt-Grupp, I. and Ruthmann, A. (1989).** Microsporidian infection in *Pimpla turionellae* (Ichneumonidae: Hymenoptera): characteristics and reaction with *Calcofluor White*. *Parasitology Research* 76 (1):74 - 79.



- Pavuk, D.M., Williams, C.E. and Taylor, D.H. (1995).** Parasitism of *Plathypena scabra* (Lepidoptera: Noctuidae) by *Sinophorus teratis* (Hymenoptera: Ichneumonidae). Great Lakes Entomologist 28 (3-4): 205 - 206.
- Pavuk, D.M. and Hughes, L.L. (1998).** The parasitoid complex of first generation *Ostrinia nubilalis* (Lepidoptera: Pyralidae) larvae in northwest Ohio. Great Lakes Entomologist 31 (3-4): 169 - 172.
- Pavuk, D.M. and Stinner, B.R. (1991).** New Lepidoptera-parasitoid associations in weedy corn plantings: a potential alternate host for *Ostrinia nubilalis* (Lepidoptera: Pyralidae) parasitoids. Great Lakes Entomologist 24 (4): 219 - 223.
- Pavuk, D.M. and Williams, R.N. (1988).** A sesiid host record for *Pterocormus chasmodops* (Hymenoptera: Ichneumonidae). Proceedings of the Entomological Society of Washington 90 (1): 105.
- Pawar, C.S., Bhatnagar, V.S. and Jadhav, D.R. (1989).** *Campoletis chlorideae* Uchida (Hymenoptera: Ichneumonidae) as a parasite of *Helicoverpa armigera* (Hub.) (Lepidoptera: Noctuidae) in southwest India. Proceedings of the Indian Academy of Sciences, Animal Sciences 98 (4): 259 - 265.
- Peigler, R.S. and Vinson, S.B. (1988).** Parasitoid and hostplant records for genus *Schinia* (Noctuidae) in Texas. Journal of the Lepidopterists' Society, USA 42 (2): 144 - 145.
- Pekel, S., Kolarov, J. and Ozbek, H. (2000).** New records of the subfamily Banchinae (Hymenoptera: Ichneumonidae) from Turkey. Journal of the Entomological Research Society 2 (3): 1 - 4.
- Pekel, S. and Ozbek, H. (2000).** Faunistic and systematic study on the subfamily of Cremastinae (Hymenoptera: Ichneumonidae) in Erzurum province. Turkiye Entomoloji Dergisi 24 (3): 215 -228.
- Pemberton, R.W., Lee, J.H., Reed, D.K., Carlson, R.W. and Han, H.Y. (1993).** Natural enemies of the Asian gypsy moth (Lepidoptera: Lymantriidae) in South Korea. Annals of the Entomological Society of America 86 (4): 423 - 440.
- Pemberton, R.W. and Lee, J.H. (1996).** The influence of extrafloral nectaries on parasitism of an insect herbivore. American Journal of Botany 83 (9): 1187 - 1194.
- Pennington, A., Marris, G., Renn, N., Wakefield, M. and Edwards, J.P. (2000).** Nematode infection of lepidopteran pupae is facilitated by the presence of a parasitoid wasp. TheBCPC Conference: Pests and diseases, Volume 3. Proceedings of an international conference held at the Brighton Hilton Metropole Hotel, Brighton, UK, 13-16 November 2000. 1013 - 1018.
- Pereira, P., Adler, C. and Schoeller, M. (1998).** The use of pheromone traps for monitoring *Ephestia kuehniella* Zeller (Lepidoptera: Pyralidae) and detection of parasitoids in flour mills. Integrated protection of stored products. Proceedings of the meeting at Zurich, Switzerland, 31 August-2 September, 1997. Bulletin OILB SROP 21 (3): 111 - 117.
- Perez-Moreno, I., Saenz-de-Cabezón, F.J. and Marco, V. (2000).** Evaluation of natural parasitism on hibernate pupae of the European grape moth (*Lobesia botrana* Den. &

- Schiff.) in vineyard of La Rioja. Boletín de Sanidad Vegetal, Plagas 26 (4): Supplement, 715 - 721.
- Periquet, G., Hedderwick, M.P. and El Agoze, M. (1993).** Sex determination in the hymenopteran *Diadromus pulchellus* (Ichneumonidae): validation of the one-locus multi-allele model. Heredity 70 (4): 420 - 427.
- Perju, T. and Moldovan, I. (1991).** Entomofauna of *Centaurea* species with a view to biological control. Buletinul Institutului Agronomic Cluj Napoca. Seria Agricultura si Horticultura 45 (2): 73 - 84.
- Peter, C. and David, B.V. (1991).** Natural enemies of the pumpkin caterpillar *Diaphania indica* (Lepidoptera: Pyralidae) in Tamil Nadu. Entomophaga 36 (3): 391 - 394.
- Peter, C. and David, B.V. (1991).** Population dynamics of the pumpkin caterpillar, *Diaphania indica* (Saunders) (Lepidoptera: Pyralidae). Tropical Pest Management 37 (1): 75 - 79.
- Peter, C. and David, B.V. (1993).** Studies on the hyperparasites of *Diaphania indica* (Lepidoptera: Pyralidae) through *Apanteles taragamae* (Hymenoptera: Braconidae). Journal of the Bombay Natural History Society 90 (3): 412 - 416.
- Peters, T.M. and Van Driesche, R.G. (1987).** Potential for increased use of biological control agents against pests of forage crops in Massachusetts. Research Bulletin Massachusetts Agricultural Experiment Station 718: 75 - 86.
- Piegler, R.S. (1994).** Catalog of parasitoids of Saturniidae of the world. Journal of Research on the Lepidoptera 33: 121 pp.
- Piekarska, H. (1991).** Studies on the preimaginal development of the ichneumonid *Itopectis maculator* (Fabr.) (Hymenoptera: Ichneumonidae) I. Morphology of the preimaginal stages. Roczniki Akademii Rolniczej w Poznaniu, Ogrodnictwo 19: 93 - 100.
- Piekarska, H., Strzykala-Trybula, E. and Boniecki, P. (1991).** Studies on the preimaginal development of the ichneumonid *Itopectis maculator* (Fabr.) (Hymenoptera: Ichneumonidae) II. Duration of development. Roczniki Akademii Rolniczej w Poznaniu, Ogrodnictwo 19: 101 - 105.
- Piekarska, H. and Kusmierczak, B. (1990).** Contribution to knowledge of the fauna of ichneumonids (Hymenoptera: Ichneumonidae) parasitising apple leaf-rollers in orchards in the vicinity of Poznan. Roczniki Akademii Rolniczej w Poznaniu, Ogrodnictwo 18: 53 - 64.
- Piekarska-Boniecka, H. (1991).** *Itopectis maculator* (Fabr.) (Hymenoptera: Ichneumonidae) a hyperparasitoid in orchards in the vicinity of Poznan. Wiadomosci Entomologiczne 10 (3): 189.
- Piekarska-Boniecka, H. (1991).** Some information on the bionomics of the ichneumonid wasp *Itopectis maculator* (Fabr.) (Hymenoptera: Ichneumonidae). Wiadomosci Entomologiczne 10 (4): 227 - 232.

- Piekarska-Boniecka, H. (1994).** Contribution to knowledge of the leaf-rollers (Lepidoptera: Tortricidae) and their parasitoids (Hymenoptera: Ichneumonidae) occurring on red currant in the environs of Poznan. *Wiadomosci Entomologiczne* 13 (3): 185 - 190.
- Piekarska-Boniecka, H. (1997).** Pimplinae (Hymenoptera: Ichneumonidae) occurring in the fruit-growing environment in the environs of Poznan. Part 2. *Roczniki Akademii Rolniczej w Poznaniu, Ogrodnictwo* 25: 55 - 64.
- Piekarska-Boniecka, H. (1998).** Pimplinae (Hymenoptera: Ichneumonidae) occurring in the fruit-growing environment in the environs of Poznan. Part 3. *Roczniki Akademii Rolniczej w Poznaniu, Ogrodnictwo* 26: 91 - 96.
- Piekarska-Boniecka, H. (1999).** Pimplinae and Diacritinae (Hymenoptera: Ichneumonidae) of selected habitats of an agricultural landscape of Poznan environs. Part 1. *Wiadomosci Entomologiczne* 18 (2): 77 - 84.
- Piekarska- Boniecka, H., Wilkaniec, B., Borowiak-Sobkowiak, B. and Trzcinski, P. (2002).** A natural control of a fruit tree tortix (*Archips rosanus* (L.)) occurring in apple orchard and black currant plantation in Poznan Region. *Progress in Plant Protection* 42 (1): 223 - 227.
- Piekarska-Boniecka, H., Wilkaniec, B., Trzcinski, P. and Zawieja, M. (1999).** Parasitoids of Ichneumonidae family bred from larvae and pupae of leaf-rollers occurring in an apple orchard in the environs of Gostyn. *Progress in Plant Protection* 39 (2): 452 - 454.
- Piekarska-Boniecka, H. and Sawoniewicz, J. (1994).** Contribution to the knowledge of Pimplinae (Hymenoptera: Ichneumonidae) occurring in the Biebrza Valley Lower Basin. *Wiadomosci Entomologiczne* 13 (1): 41 - 53.
- Piekarska-Boniecka, H. and Suder-Byttner, A. (2002).** Pimplinae, Diacritinae and Poemeniinae (Hymenoptera: Ichneumonidae) occurring in a fruit-growing environment in Przybroda. *Journal of Plant Protection Research* 42 (3): 221 - 227.
- Piekarska-Boniecka, H. and Tian, M.Y. (1998).** The effects of ichneumonids (Hymenoptera) and other parasitoids on the rose leafroller, *Archips rosanus* (L.) (Lepidoptera: Tortricidae), from black currant. *Roczniki Akademii Rolniczej w Poznaniu, Ogrodnictwo* 26: 79-89.
- Piekarska-Boniecka, H. and Wilkaniec, B. (1998).** Parasitoids of the family Ichneumonidae reared from larvae of the apple blossom weevil (*Anthonomus pomorum* (L.)) in the environs of Poznan. *Progress in Plant Protection* 38 (2): 447 - 449.
- Pillai, G.B. and Nair, K.R. (1987).** *Brachycoryphus nursei* (Cameron) (Hymenoptera: Ichneumonidae), a parasitoid of *Opisina arenosella* Wlk. on coconut in Kerala, India. *Journal of Plantation Crops* 15 (1):9 - 14.
- Pillai, G.B. and Nair, K.R. (1989).** Observations of *Xanthopimpla punctata* punctata F. (Hymenoptera: Ichneumonidae), a pupal parasitoid of *Opisina arenosella*. *Journal of Plantation Crops* 16: Supplement, 173 - 177.

- Pillai, G.B. and Nair, K.R. (1990).** Role of ichneumonid parasitoids in the biological suppression of the coconut leaf eating caterpillar, *Opisina arenosella* Wlk. Indian Coconut Journal Cochin 21 (4): 3 - 10.
- Pillai, G.B. and Nair, K.R. (1992).** A note on the biology of *Xanthopimpla nana nana* Schulz. (Hymenoptera: Ichneumonidae), a pupal parasitoid of *Opisina arenosella* Wlk. on coconut. Indian Coconut Journal Cochin 23 (2): 2 - 4.
- Pimbert, M.P. and Srivastava, C.P. (1989).** Vegetation management and the biological control of *Helicoverpa armigera* in chickpea. International Chickpea Newsletter 21: 16 - 19.
- Pinzon-Cortes, J., Florez-Daza, E. and Palacio, E. (2000).** Report of *Tromatobia* sp. (Hymenoptera: Ichneumonidae) as a parasitoid on eggs of the spider *Araneus granadensis* (Araneae: Araneidae). Entomologo 28: 89, 2 - 4.
- Pisarska, R. (1995).** The succession of fauna in the Bialowieza primeval pine forest. II. Fragmenta Faunistica 38: 14 - 25, 333 - 494.
- Pitcairn, M.J. (1987).** The diapause phenology and population dynamics of the Egyptian alfalfa weevil [*Hypera postica*] and two of its parasitoids [*Bathyplectes curculionis* and *Tetrastichus incertus*] in California. Dissertation Abstracts International, B Sciences and Engineering 47 (7): 2749.
- Plaskota, E. and Dabrowski, Z.T. (1986).** Biological principles of leek moth (*Acrolepia assectella* Zeller, Lepidoptera: Plutellidae) control. I. Host plants, damage caused by leek moth larvae, and leek moth parasites. Annals of Warsaw Agricultural University SGGW AR, Horticulture 13: 27 - 34.
- Plaza, A.S., Leon, E.M., Fonseca, J.P. and De La Cruz, J. (1992).** Biology, behaviour and natural enemies of *Neoleucinodes elegantalis* (Guenee). Revista Colombiana de Entomologia 18 (1): 32 - 37.
- Polaszek, A., Delvare, G., Blary, D., Polaszek, A., Delvare, G. and Blary, D. (2000).** Cereal stem borers in Africa: economic importance, taxonomy, natural enemies and control methods. Les- foreurs des tiges de cereales en Afrique: importance economique, systematique, ennemis naturels et methodes de lutte. 534 pp., 41 pp.
- Polaszek, A., Fitton, M.G., Bianchi, G. and Huddleston, T. (1994).** The parasitoids of the African white rice borer, *Maliarpha separatella* Ragonot (Lepidoptera: Pyralidae). Bulletin of Entomological Research 84 (1): 65 - 90.
- Polaszek, A. (1998).** African cereal stem borers: economic importance, taxonomy, natural enemies and control. CAB INTERNATIONAL; Wallingford; UK, x + 530 pp.
- Ponder, B.M. and Seabrook, W.D. (1988).** Biology of the blueberry leaf-tier *Croesia curvalana* (Kearfott) (Tortricidae): a field and laboratory study. Journal of the Lepidopterists' Society, USA 42 (2): 120 - 131.
- Porcelli, F. (1988).** Morphology of the larval instars of *Habronyx heros* Wesmael (Ichneumonidae: Anomaloniinae) with notes on its biology. Entomologica 23: 171 - 189.

- Porter, C.C. (1973).** A new Argentine genus of the subtribe Mesostenina (Hymenoptera, Ichneumonidae). *Acta Zoologica Lilloana* 26: 301-311.
- Porter, C.C. (1976).** A new Argentine genus of Theriini (Hymenoptera, Ichneumonidae). *Psyche* 83: 185-191.
- Porter, C.C. (1985).** *Trachysphyrus* and the new genus *Aeliopotes* in the coastal desert of Peru and north Chile (Hymenoptera: Ichneumonidae). *Pysche* 92: 513-545.
- Porter, C.C. (1987).** A revision of the *Chilean Mesostenini* (Hymenoptera: Ichneumonidae). *Contributions of the American Entomological Institute* 23 (3): 1 - 164.
- Porter, C.C. (1989).** Una revision taxonomica de las Groteini Chilenos (Hymenoptera: Ichneumonidae). *Acta Entomologica Chilena* 15: 143-162.
- Porter, C.C. (1989).** New *Chilean Itamuton* (Hymenoptera: Ichneumonidae: Mesostenini) reared from *Elicura litigator* (Neuroptera: Myrmeleontidae). *Florida Entomologist* 72 (4): 660 - 664.
- Porter, C.C. (1991).** Biogeography of *Chilean Ichneumonidae* (Hymenoptera: Ichneumonidae). *Acta Entomologica Chilena* 16: 37 - 68.
- Porter, C.C. (1993).** First record of Phrudinae (Hymenoptera: Ichneumonidae) from South America with notice of a new genus and species from Chile. *Journal of the New York Entomological Society* 101 (1): 130 - 134.
- Porter, C.C. (1998).** A review of the genus *Myrmeleonostenus* (Hymenoptera: Ichneumonidae: Mesostenini) with description of two new species from Taiwan. *Journal of the Taiwan Museum* 51 (2): 93 - 105.
- Porter, C.C. (1998).** Guía de los géneros de Ichneumonidae en la región neantártica del sur de Sudamérica. *Opera Lilloana* 42: 1-234
- Porter, C.C. (1999).** First notice of *Therion* from South America with description of two new species from Argentina and Bolivia (Hymenoptera: Ichneumonidae). *Insecta Mundi* 13 (1-2): 85 - 91.
- Pramuk, J.B. and Fox, L.R. (1998).** Demography and larval development of the endoparasitoid *Patricloides montanus* Cresson (Hymenoptera: Ichneumonidae). *Journal of the Kansas Entomological Society* 71 (1): 44 - 50.
- Prasad, D., Chand, P., Deka, N.K. and Prasad, R. (1989).** Population dynamics of *Heliothis armigera* (Hub.) on chickpea. *Giornale Italiano di Entomologia* 4: 22, 223 - 228.
- Prasad, D., Chand, P. and Haque, M.F. (1987).** Parasitisation of *Heliothis armigera* (Hub.) in chickpea field as affected by pesticidal seed treatment. *Research and Development Reporter* 4 (2): 236 - 238.
- Prasad, D. and Chand, P. (1986).** *Campoletis chlorideae* Uchida a new parasitoid of *Heliothis armigera* (Hubner) in Ranchi, Bihar. *Indian Journal of Entomology* 48 (2): 231 - 232.
- Press, J.W. (1989).** Compatibility of *Xylocoris flavipes* (Hemiptera: Anthocoridae) and *Venturia canescens* (Hymenoptera: Ichneumonidae) for suppression of the almond moth,

- Cadra cautella* (Lepidoptera: Pyralidae). *Journal of Entomological Science* 24 (1): 156 - 160.
- Press, J.W. and Arbogast, R.T. (1991).** Effect of low temperature on survival of immatures of the parasite *Venturia canescens* (Gravenhorst) (Hymenoptera: Ichneumonidae). *Journal of the Kansas Entomological Society* 64 (3): 345 - 348.
- Prevost, G., Davies, D.H. and Vinson, S.B. (1990).** Evasion of encapsulation by parasitoid correlated with the extent of host hemocyte pathology. *Entomologia Experimentalis et Applicata* 55 (1): 1 - 10.
- Prevost, G. and Vinson, S.B. (1988).** Immunity relationships between the parasitoid *Campoletis sonorensis* and lepidopteran hosts: a hemocyte study. *Colloques de l'INRA* 48: 103 - 104.
- Price, E. and Price, H. (1993).** A gall or not a gall? *Cecidology* 8 (1): 27 - 28.
- Prins, A.H., Nell, H.W. and Klinkhamer, P.G.L. (1992).** Size-dependent root herbivory on *Cynoglossum officinale*. *Oikos* 65 (3): 409 - 413.
- Prozell, S., Scholler, M., Adler, C. and Schoeller, M. (1998).** Insect fauna of a bakery, processing organic grain and applying *Trichogramma evanescens* Westwood. Integrated protection of stored products. Proceedings of the meeting at Zurich, Switzerland, 31 August-2 September, 1997. *Bulletin OILB SROP* 21 (3): 39 - 44.
- Pulkkinen, M. (1989).** The distribution and ecology of the *Strobilomyia* flies (Diptera: Anthomyiidae) infesting larch seed and cones in Finland. *Annales Entomologici Fennici* 55 (2): 41 - 47.
- Purrington, F.F. and Stinner, D.H. (1988).** The bindweed plume moth, *Emmelina monodactyla* (Lepidoptera: Pterophoridae): first host record for *Phaeogenes vincibilis* (Hymenoptera: Ichneumonidae). *Great Lakes Entomologist* 21 (4): 157 - 158.
- Puvuk, D.M. and Stinner, B.R. (1992).** Influence of weed communities in corn plantings on parasitism of *Ostrinia nubilalis* (Lepidoptera: Pyralidae) by *Eriborus terebrans* (Hymenoptera Ichneumonidae). *Biological Control* 2 (4): 312 - 316.

## Q

- Qian, F. (1987).** A research on the biology of *Casinarina nigripes* Gravenhorst. *Journal of Nanjing Forestry University* 4: 32 - 38.
- Quednau, F.W. (1990).** Introduction, permanent establishment, and dispersal in Eastern Canada of *Olesicampe geniculatae* Quednau and Lim (Hymenoptera: Ichneumonidae), an important biological control agent of the mountain ash sawfly, *Pristiphora geniculata* (Hartig) (Hymenoptera: Tenthredinidae). *Canadian Entomologist* 122 (9-10): 921 - 934.
- Quicke, D.L.J. (1991).** Ovipositor mechanics of the braconine wasp genus *Zaglyptogastra* and the ichneumonid genus *Pristomerus*. *Journal of Natural History* 25 (4): 971 - 977.
- Quicke, D.L.J., Fitton, M.G., Notton, D.G., Broad, G.R., Dolphin, K., Austin, A.D. and Dowton, M. (2000).** Phylogeny of the subfamilies of Ichneumonidae (Hymenoptera): a simultaneous molecular and morphological analysis. *Fourth International Hymenoptera*

Conference, held in Canberra, Australia, in January 1999. Hymenoptera: evolution, biodiversity and biological control 74 - 83.

**Quicke, D.L.J., Fitton, M.G. and Ingram, S. (1992).** Phylogenetic implications of the structure and distribution of ovipositor valvelli in the Hymenoptera (Insecta). *Journal of Natural History* 26 (3): 587 - 608.

**Quicke, D.L.J. and Kruft, R.A. (1995).** Latitudinal gradients in North American braconid wasp species richness and biology. *Journal of Hymenoptera Research* 4: 194 - 203.

**Quicke, D.L.J., Basibuyuk, H. H., Fitton, M. G. and Rasnitsyn, A. P. (1999).** Morphological, palaeontological and molecular aspects of ichneumonoid phylogeny (Hymenoptera, Insecta). *Zool. Scripta* 28: 175-202.

**Quicke, D.L.J. and Shaw, M.R. (2004).** Cocoon silk chemistry in parasitic wasps (Hymenoptera, Ichneumonidea) and their hosts. *Biological Journal of the Linnean Society* 81: 161 – 170.

**Quinn, M.A., Kepner, R.L., Walgenbach, D.D., Foster, R.N., Bohls, R.A., Pooler, P.D., Reuter, K.C. and Swain, J.L. (1993).** Grasshopper stages of development as indicators of nontarget arthropod activity: implications for grasshopper management programs on mixed-grass rangeland. *Environmental Entomology* 22 (3): 532 - 540.

## R

**Radcliffe, E.B. and Flanders, K.L. (1998).** Biological control of alfalfa weevil in North America. *Integrated Pest Management Reviews* 3 (4): 225 - 242.

**Rafal'skii, V.Yu. (1989).** Ichneumonids - parasitoids of dendrophilous geometrids in the central forest-steppe of the Ukrainian SSR. *Vestnik Zoologii* 1: 81 - 83.

**Ramadhane, A.M. and Ihsan, S.I. (1999).** Biological characteristics related to the life cycle of the cabbage large butterfly *Pieris brassica* L. (Lepidoptera: Pieridae) and associated endo-parasitoids. *Arab Journal of Plant Protection* 17 (1): 45 - 48.

**Rao, S.N. and Nikam, P.K. (1983).** A new genus and species of Mesosteninae (Hymenoptera: Ichneumonidae) from India. *Entomon* 8: 179-182.

**Rao, G.M.V.P., Ramani, S. and Singh, S.P. (2001).** Field parasitisation of the maize stem borer, *Chilo partellus* (Swinhoe) (Lepidoptera: Pyralidae) in Bangalore. *Journal of Biological Control* 15 (2): 193 - 195.

**Rao, R.S.N., Sreedhar, U., Rao, S.G. and Satyavani, J.V.R. (1994).** Bio-environmental control of *Heliothis armigera* in FCV tobacco. *Tobacco Research* 20 (1): 36 - 39.

**Rasnitsyn, A.P. (1975).** Hymenoptera Apocrita of Mesozoic. (in Russian) *Trudy Paleontologicheskogo Instituta* 147: 1-132.

**Rasnitsyn, A.P. and Siitan, U.V. (1981).** Ichneumoninae. *In*: D.R. Kasparyan (ed.), [A guide to the identification of insects of the European part of the USSR. Vol. 3. Hymenoptera] (in Russian) . Moscow: Nauka Press. 668 pp.

**Rasnitsyn, A. P. (1983).** Ichneumonoidea (Hymenoptera) from the lower Cretaceous of Mongolia. *Contrib. Amer. Entomol. Inst.* 20: 259-265.

- Rasnitsyn, A. P. and Sharkey, M. J. (1988).** New Eoichneumonidae from the early Cretaceous of Siberia and Mongolia (Hymenoptera: Ichneumonoidea). Pp. 169-197 in: Gupta, V. K., ed., *Advances in Parasitic Hymenoptera Research*, E. J. Brill, Leiden.
- Rauleder, H. (2002).** Observations on the biology of the plum fruit moth (*Cydia funebrana*). *Gesunde Pflanzen* 54 (8): 241 - 248.
- Ravi, G. and Verma, S. (1997).** Seasonal incidence of chickpea pod borer *Helicoverpa armigera* and its larval parasitoid on chick pea crop. *Indian Journal of Entomology* 59 (4): 359 - 361.
- Read, P.E.C., Donovan, B.J. and Schroeder, N.C. (1990).** Rearing and distribution of the introduced wasp parasitoid *Sphecophaga vesparum* throughout New Zealand. *Proceedings of the Forty Third New Zealand Weed and Pest Control Conference* 191 - 194.
- Reed, D.J., Begon, M. and Thompson, D.J. (1996).** Differential cannibalism and population dynamics in a host-parasitoid system. *Oecologia* 105 (2): 189 - 193.
- Reineke, A., Asgari, S., Ma, G., Beck, M. and Schmidt, O. (2002).** Sequence analysis and expression of a virus-like particle protein, VLP2, from the parasitic wasp *Venturia canescens*. *Insect Molecular Biology* 11 (3): 233 - 239.
- Renault, S., Petit, A., Benedet, F., Bigot, S. and Bigot, Y. (2002).** Effects of the *Diadromus pulchellus* ascovirus, DpAV-4, on the hemocytic encapsulation response and capsule melanization of the leek-moth pupa, *Acrolepiopsis assectella*. *Journal of Insect Physiology* 48 (3): 297 - 302.
- Renault, S., Rouleux-Bonnin, F., Periquet, G. and Bigot, Y. (1999).** Satellite DNA transcription in *Diadromus pulchellus* (Hymenoptera). *Insect Biochemistry and Molecular Biology* 29 (2): 103 - 111.
- Rey Del Castillo, C. (1990).** Contribution to the knowledge of the Lissonotini of Spain II. Genus *Lissonota*, 1829 (in part) (Hym.: Ichneumonidae). *Eos Madrid* 65 (2): 209 - 250.
- Rey del Castillo, C. (1990).** Los Icneumonidos de la subfamilia Banchinae en las Islas Canarias (Hym., Ichneumonidae). *Vieraea* 18: 353-369.
- Rey Del Castillo, C. (1992).** Revision of western Palaearctic species of the subgenus *Loxonota* Aubert, 1978 (Hymenoptera: Ichneumonidae). *Annales de la Societe Entomologique de France* 28 (2): 133 - 156.
- Rey Del Castillo, C. (1989).** Palaearctic species of Banchini and Glyptini (Hym.: Ichneumonidae) represented in the Museo Nacional de Ciencias Naturales de Madrid. *Boletin de la Asociacion Espanola de Entomologia* 13: 183 - 193.
- Rey Del Castillo, C. (1989).** Contribution to the knowledge of the Lissonotini (Hym.: Ichneumonidae) in Spain: I. Genera *Alloplasta* Foerster, 1868, *Cryptopimpla* Taschenberg, 1863 and new data about *Syzeuctus* Foerster, 1868. *Boletin de la Asociacion Espanola de Entomologia* 13: 143 - 156.
- Rey Del Castillo, C. (1991).** First record of Adelognathinae, Neorhacodinae and Phrudinae in Spain (Hymenoptera: Ichneumonidae). *Boletin de la Asociacion Espanola de Entomologia* 15: 153 - 163.



- Reymonet, C., Gari, J.V.F. and Mari, J.M. (1993).** Survey of the parasitoids of the European sunflower moth *Homoeosoma nebulella* (Lep.: Pyralidae) in the Palearctic region. *Entomophaga* 38 (3): 355 - 358.
- Reznik, S.Ya. (1993).** Food specialization of insects (ecological, physiological, evolutionary aspects). *Trudy Zoologicheskogo Instituta, Rossiiskaya Akademiya Nauk* 193: 259 pp.
- Richards, E.H. and Parkinson, N.M. (2000).** Venom from the endoparasitic wasp *Pimpla hypochondriaca* adversely affects the morphology, viability, and immune function of hemocytes from larvae of the tomato moth, *Lacanobia oleracea*. *Journal of Invertebrate Pathology* 76 (1): 33 - 42.
- Richman, D.B., Thompson, D.C. and O'Mara, J. (1992).** Effects of leaf-tiers on broom snakeweed in central New Mexico. *Southwestern Entomologist* 17 (2): 187 - 189.
- Richmond, J.A., Werner, R.A. and Drooz, A.T. (1995).** Larch sawfly, *Pristiphora erichsonii* (Hymenoptera: Tenthredinidae) and its parasitoids from Alaska. *Journal of the Entomological Society of British Columbia* 92: 25 - 27.
- Riggin, T.M., Espelie, K.E., Wiseman, B.R. and Isenhour, D.J. (1993).** Distribution of fall armyworm (Lepidoptera: Noctuidae) parasitoids on five corn genotypes in south Georgia. *Florida Entomologist* 76 (2): 292 - 302.
- Riggin-Bucci, T.M. and Gould, F. (1997).** Impact of intraplot mixtures of toxic and nontoxic plants on population dynamics of diamondback moth (Lepidoptera: Plutellidae) and its natural enemies. *Journal of Economic Entomology* 90 (2): 241 - 251.
- Rijo, E., Jimenez, J., Vento, D., Caballero, S., Concepcion, E. and Barzagas, I. (1989).** Distribution of *Diadegma* sp. in the main tobacco growing areas of Cuba. *Ciencia Tecnica en la Agricultura, Proteccion de Plantas* 12 (1): 7 - 15.
- Rimsaite, J. and Jonaitis, V. (2002).** Distribution of parasitoids of fungi humificators in Lithuania. *Zemdirbyste, Mokslo Darbai* 79: 72 - 82.
- Rio-Mora Del, A. (1990).** Biology and chemical control of *Synanthedon cardinalis* in the Purepecha Mountains, Michoacan. *Ciencia Forestal*, publ. 1994, 15: 68, 3-27.
- Ritchie, J.M., Polaszek, A., Abeyasekera, S., Minja, E.M. and Mviha, P. (2000).** Insect pest incidence in seed pods of pigeonpea genotypes in on-farm trials in southern Malawi. *International Chickpea and Pigeonpea Newsletter* 7: 50 - 52.
- Robert, K. and Danielle, R.R. (1988).** The dynamics of a system of competitive solitary endoparasitic species of Hymenoptera: *Diadromus collaris* and *D. pulchellus*. *Acta Oecologica, Oecologia Generalis* 9 (1): 3 - 12.
- Roch, J.F. (1990).** Annotated list of Hymenoptera collected in Granby, census division of Shefford, Quebec. *Faberies* 15 (2): 48 - 52.
- Rodriguez Del Bosque, L.A. and Smith, J.W. Jr. (1990).** Parasitization of *Diatraea lineolata* pupae and diapausing larvae by several exotic parasites. *Florida Entomologist* 73 (4): 703 - 705.

- Rodriguez Del Bosque, L.A. and Smith, J.W. Jr. (1997).** Biological control of maize and sugarcane stemborers in Mexico: a review. *Insect Science and its Application* 17 (3-4): 305 - 314.
- Roland, J. (1994).** After the decline: what maintains low winter moth density after successful biological control? *Journal of Animal Ecology* 63 (2): 392 - 398.
- Rollard, C. (1988).** Note on the biology of ichneumonid wasps whose larvae feed on spider eggs. *Colloques de l'INRA* 48: 127 - 128.
- Rollard, C. (1990).** An ecological and biological approach to the spider/arachnophagous insect interaction through the example of *Argiope bruennichi* Ar., Argiopidae/*Tromatobia ornata* (Hym.: Ichneumonidae). *Bulletin de la Societe Zoologique de France* 115 (4): 379 - 385.
- Roshandel, S. and Ebrahimi, E. 2003.** Biology and seasonal parasitism of *Bathyplectes anarus* (Thomson), parasitoid of *Hypera postica* Gill. in Chahar Mahal Va Bakhtiari province. *Journal of Entomological Society of Iran*, 22(2): 1-15.
- Rossini, C., Hoebeke, E.R., Iyengar, V.K., Conner, W.E., Eisner, M. and Eisner, T. (2000).** Alkaloid content of parasitoids reared from pupae of an alkaloid-sequestering arctiid moth (*Utetheisa ornatrix*). *Entomological News* 111 (4): 287 - 290.
- Rotheray, G.E. (1988).** Larval morphology and feeding patterns of four *Cheilosia* species (Diptera: Syrphidae) associated with *Cirsium palustre* L. Scopoli (Compositae) in Scotland. *Journal of Natural History* 22 (1): 17 - 25.
- Rotheray, G.E. (1990).** A new species of *Bioblapsis* (Hymenoptera: Ichneumonidae) from Scotland parasitising a mycophagous hoverfly, *Cheilosia longula* (Diptera: Syrphidae). *Entomologica Scandinavica* 21 (3): 277 - 280.
- Rothman, L.D., Darling, D.C. and Goodall, D.W. (1990).** Parasitoids of the goldenrod gall moth: effects of scale on spatial density dependence. *Oecologia* 83 (1): 1 - 6.
- Rotrekl, J. (1994).** Effect of insecticides on the beneficial insect fauna in lucerne (*Medicago sativa* L.). *Ochrana Rostlin* 30 (1): 67 - 77.
- Rouleux-Bonnin, F., Renault, S., Rabouille, A., Periquet, G. and Bigot, Y. (1999).** Free serosal cells originating from the embryo of the wasp *Diadromus pulchellus* in the pupal body of parasitized leek-moth, *Acrolepiosis assectella*. Are these cells teratocyte-like? *Journal of Insect Physiology* 45 (5): 479 - 484.
- Rowshandel, S. (2000).** Biology and efficiency of *Bathyplectes anarus* in biological control of alfalfa weevil *Hypera postica* Gyll. *Proceedings of 14<sup>th</sup> Iranian Plant Protection Congress*, p. 28.
- Rubin, G. (2000).** Parasite complex of *Thaumetopoea pityocampa* Schiff. in the forest district Split. *Sumarski List* 124 (7-8): 363 - 372.
- Ryabchinskaya, T.A. and Kharchenko, G.L. (1995).** Parasitoids of leafrollers in apple orchards. *Zashchita Rastenii Moskva* 7: 16 - 17.

- Saaksjarvi, I.E. (1998).** One Cryptinae genus (Hymenoptera: Ichneumonidae) new to Finland. *Entomologica Fennica* 9 (1): 8
- Sabahi, Q., Talebi, K. and Kharrazi, A. (2002).** Effects of different doses of four insecticides on mortality and longevity of *Bathyplectes curculionis* (Thompson) (Hymenoptera: Ichneumonidae) the larval parasitoid of alfalfa weevil *Hypera postica* (Gyll.) (Coleoptera: Curculionidae). *Applied Entomology and Phytopathology* 69 (2): 69 - 77.
- Sabahi, Q., Talebi, Kh. and Radjabi, Gh. (2002).** Effect of four organophosphorus insecticides on mortality and fecundity of *Bathyplectes anurus* (Thompson) (Hymenoptera: Ichneumonidae) larval parasitoid of alfalfa weevil *Hypera postica* (Coleoptera: Curculionidae). *Proceedings of 15<sup>th</sup> Iranian Plant Protection Congress*, p. 27.
- Sacchetti, P. (1990).** Observations on the activity and bioethology of the entomophagous insects of *Prays oleae* (Bern.) in Tuscany. II. Incidence of parasitoids. *Redia* 73 (2): 405 - 421.
- Sachan, J.N. and Bhaumik, R. (1998).** Effect of insecticidal spray on parasitization level of *Campoletis chlorideae* on *Helicoverpa armigera* in chickpea ecosystem. *Indian Journal of Pulses Research* 11 (2): 70 - 75.
- Sachan, J.N. and Bhaumik, R. (1998).** Extent of parasitization of *Campoletis chlorideae* - a larval parasite of *Helicoverpa armigera* damaging chickpea. *Indian Journal of Pulses Research* 11 (2): 65 - 69.
- Saedi, K. (2002).** Preliminary studies on natural enemies of the Alfalfa weevil in Yasouj region. *Proceedings of 15<sup>th</sup> Iranian Plant Protection Congress*, p. 27.
- Saharaoui, L. (1994).** Survey and study of some bioecological aspects of entomophagous coccinellids (Coleoptera: Coccinellidae) in Algeria. *Journal of African Zoology* 108 (6): 537 - 546.
- Sait, S.M., Andreev, R.A., Begon, M., Thompson, D.J., Harvey, J.A. and Swain, R.D. (1995).** *Venturia canescens* parasitizing *Plodia interpunctella*: host vulnerability - a matter of degree. *Ecological Entomology* 20 (2): 199 - 201.
- Sait, S.M., Begon, M., Thompson, D.J., Harvey, J.A. and Hails, R.S. (1997).** Factors affecting host selection in an insect host-parasitoid interaction. *Ecological Entomology* 22 (2): 225 - 230.
- Sait, S.M., Begon, M., Thompson, D.J. and Harvey, J.A. (1996).** Parasitism of baculovirus-infected *Plodia interpunctella* by *Venturia canescens* and subsequent virus transmission. *Functional Ecology* 10 (5): 586 - 591.
- Sait, S.M., Thomas, M.B. and Kedwards, T. (1999).** Multi-species microcosms as intermediates between theory and the natural world. *Challenges in applied population biology*, Royal Holloway, University of London, UK, 8-9 July 1999. *Aspects of Applied Biology* 53: 65 - 72.

- Saito, O., Mizushima, S., Okuyama, S., Hanada, T., Torikura, H., Hachiya, K. and Sato, K. (1998).** Biology of the diamondback moth, *Plutella xylostella* (L.), in Hokkaido. Research Bulletin of the Hokkaido National Agricultural Experiment Station 167: 69 - 110.
- Sakanoshita, A., Ohta, M. and Uematsu, H. (1987).** Comparative study of ovogenesis in the parasitoids of the diamondback moth *Plutella xylostella*. Proceedings of the Association for Plant Protection of Kyushu 33: 139 - 141.
- Salas-Araiza, M.D. and Salazar-Solis, E. (1998).** Natural parasitism of Lepidoptera pests of broccoli in Bajío, Mexico. Manejo Integrado de Plagas 50: 34 - 41.
- Samoedi, D. (1988).** Seasonal population fluctuation of sugarcane top borer, *Tryporyza nivella intacta* Sn. (Pyralidae: Lepidoptera) at Jatitujuh Sugar Plantation. Majalah Perusahaan Gula 24 (4): 14 - 21.
- Samoedi, D. (1993).** Seasonal abundance of sugar cane top moth borer, *Tryporyza nivella intacta* Sn., and its natural enemies in South Sumatra. Bulletin Pusat Penelitian Perkebunan Gula Indonesia 139: 34 - 40.
- Sanborne, M. (1981).** *Ptychopsis phalara*, a new genus and species of South American Banchinae (Hymenoptera: Ichneumonidae). Canadian Entomologist 113: 519-521.
- Sanborne, M. (1981).** A revision of the genus *Sesioplex* (Hymenoptera: Ichneumonidae) in North America. Canadian Entomologist 113: 623-629.
- Sanborne, M. (1986).** On the correct application of *Rhimphoctona* Foerster (Hymenoptera: Ichneumonidae). Contributions of the American Entomological Institute 22 (6): 1-6.
- Sanborne, M. (1986).** Revision of the Nearctic species of *Xylophylax* Kriechbaumer (Hymenoptera: Ichneumonidae). Contributions of the American Entomological Institute 22 (8): 1-45.
- Sanborne, P.M. (1990).** Discovery of *Nemeritis* Holmgren (Hymenoptera: Ichneumonidae) in eastern North America. Canadian Entomologist 122 (11-12): 1267 - 1268.
- Sandanayake, W.R.M. and Edirisinghe, J.P. (1992).** *Trathala flavoorbitalis*: parasitization and development in relation to host-stage attacked. Insect Science and its Application 13 (3): 287 - 292.
- Sandanayake, W.R.M. and Edirisinghe, J.P. (1993).** Aspects of reproductive biology of *Trathala flavoorbitalis* (Cam.): a parasitoid of *Leucinodes orbonalis* (Guen.). Entomon 17 (3-4): 159 - 168.
- Sandhu, S.S., Singla, M.L. and Singh, J. (1990).** Ecology of *Heliothis armigera* (Hub.) on chick pea in Punjab. Journal of Insect Science 3 (1): 47 - 52.
- Sannino, L., Balbiani, A. and Espinosa, B. (1990).** Biology of *Athetis hospes* (Freyer, 1831), in Campania (Lepidoptera: Noctuidae: Amphipyridae). Bollettino della Societa Entomologica Italiana 122 (2): 141 - 148.
- Sannino, L., Balbiani, A. and Espinosa, B. (1993).** Biology of *Lacanobia oleracea* L. in Campania (Lepidoptera: Noctuidae: Hadeninae). Bollettino della Societa Entomologica Italiana 125 (1): 61 - 70.

- Sannino, L. and Espinosa, B. (1998).** Biological cycle of *Mamestra brassicae* and damage to horticultural crops in Campania. *Informatore Fitopatologico* 48 (5): 59 - 67.
- Sannino, L. and Espinosa, B. (1999).** Biology of *Mamestra brassicae* (Lepidoptera: Noctuidae) in Campania (South Italy). *Bollettino del Laboratorio di Entomologia Agraria "Filippo-Silvestri"* 55: 79 - 91.
- Santiago-Alvarez, C. and Caballero, P. (1990).** Susceptibility of parasitized *Agrotis segetum* larvae to a granulosis virus. *Journal of Invertebrate Pathology* 56 (1): 128 - 131.
- Santis De, L., Monetti, N.C. and Briano, De, A.E. (1987).** Natural enemies of *Rhinocyllus conicus* (Froelich, 1792) (Coleoptera: Curculionidae), a biological control agent of *Carduus thoermeri* Kazmi and *Carduus acanthoides* L. (Compositae) in Castelar, Buenos Aires Province, Argentina. *Revista de Investigacion Centro de Investigaciones para la Regulacion de Poblaciones de Organismos Nocivos* 5 (1-4): 27 - 35.
- Santis De, L. (1987).** Hymenopteran parasitoids and hyperparasitoids of *Anacraga* sp. (Lepidoptera: Dalceridae) in Brazil. *Revista Brasileira de Entomologia* 31 (1): 97 - 99.
- Santos Dos, B.B. (1989).** Parasitoids of *Mocis latipes* (Guenee, 1852) (Lepidoptera: Noctuidae). *Arquivos de Biologia e Tecnologia* 32 (3): 491 - 494.
- Santos, B.M., Sanchez, L., Reyes, M., Perez, Q. and Castillo, M. (2000).** Parasitoids of *Plutella xylostella* in the Dominican Republic. *Manejo Integrado de Plagas* 58: 76 - 77.
- Saparmamedova, N.K. (1988).** On the study of natural enemies of the plum moth *Grapholitha funebrana* Tr. (Lepidoptera: Tortricidae) in the Prikopetdag zone. *Izvestiya Akademii Nauk Turkmenskoi SSR. Seriya Biologicheskikh Nauk* 4: 40 - 43.
- Sastrosiswojo, S. (1996).** Biological control of the diamondback moth in IPM systems: case study from Asia (Indonesia). *Biological control introductions - opportunities for improved crop production. Proceedings of an International Symposium Brighton, UK, 18 November 1996.* 13 - 32.
- Sathe, T.V. (1990).** Impact of host food plant on parasitization behaviour in a larval parasitoid of *Heliothis armigera* (Hubner). *Proceedings of the Indian Academy of Sciences, Animal Sciences* 99 (3): 233 - 242.
- Sathe, T.V. and Gupta, V.K. (1988).** Intrinsic rates of increase and interspecific relationships between *Cotesia orientalis*, *Cotesia diurnii* and *Diadegma tarichoptilus* (Hymenoptera: Ichneumonidae), the larval parasitoids of *Exelastis atomosa*. *Advances in parasitic Hymenoptera research: Proceedings of the II. Conference on the Taxonomy and Biology of Parasitic Hymenoptera.* recd. 1993. 383 - 387.
- Sathe, T.V. and Nikam, P.K. (1986).** Life tables and intrinsic rate of increase of *Campoletis chlorideae* Uchida (Hymenoptera: Ichneumonidae), an internal larval parasitoid of *Heliothis armigera* (Hubn.). *Rivista di Parassitologia*, publ. 1988, 3: 47, 289 - 294.
- Sathe, T.V. and Santhakumar, M.V. (1988).** Factors responsible for host finding behaviour by *Campoletis chlorideae* Uchida (Hymenoptera: Ichneumonidae), a parasitoid of *Heliothis armigera* (Hubn.) (Lepidoptera: Noctuidae). *Rivista di Parassitologia*, publ. 1990, 5 (2): 233 - 240.

- Sato, H. (1995).** Comparison of community composition of parasitoids that attack leaf-mining moths (Lepidoptera: Gracillariidae). *Environmental Entomology* 24 (4): 879 - 888.
- Sato, M. and Takasu, K. (2000).** Food odor learning by both sexes of the pupal parasitoid *Pimpla alboannulatus* uchida (Hymenoptera: Ichneumonidae). *Journal of Insect Behavior* 13 (2):263 - 272.
- Saucke, H. (1996).** Biological and integrated control of *Plutella xylostella* (Lepidoptera: Yponomeutidae) and *Crociodolomia pavonana* (Lepidoptera: Pyralidae) in different cabbage-growing regions of Papua New Guinea. Justus Liebig Universitat, Fachbereich Veterinarmedizin; Giessen; Germany, 131 pp., 24 pp.
- Saucke, H., Dori, F. and Schmutterer, H. (2000).** Biological and integrated control of *Plutella xylostella* (Lep.: Yponomeutidae) and *Crociodolomia pavonana* (Lep.: Pyralidae) in brassica crops in Papua New Guinea. *Biocontrol Science and Technology* 10 (5): 595 - 606.
- Sawoniewicz, J. (1985).** Revision of European species of the subtribe Endaseina (Hymenoptera: Ichneumonidae), I. *Annales Zoologici* 39 (3): 131 - 146.
- Sawoniewicz, J. (1986).** Revision of some type-specimens of European Ichneumonidae (Hymenoptera), 2. *Annales Zoologici* 40 (5): 371 - 380.
- Sawoniewicz, J. (1986).** Structure of Ichneumonidae (Hymenoptera) communities in urban green areas of Warsaw. *Memorabilia Zoologica* 41: 103 - 139.
- Sawoniewicz, J. (1988).** Revision of some type-specimens of European Ichneumoninae (Hymenoptera), 3. *Annales Zoologici* 41 (15): 481 - 490.
- Sawoniewicz, J. and Buszko, J. (1994).** Ichneumonidae (Hymenoptera) reared from mining Lepidoptera in Poland. *Wiadomosci Entomologiczne* 13 (1): 55 - 61.
- Sawoniewicz, J. (1990).** Revision of some type-specimens of European Ichneumonidae (Hymenoptera), 5. *Annales Zoologici* 43 (12): 293 - 299.
- Saxena, H. (1992).** Field behaviour and natural parasitisation of *Mahasena graminivora* Hamps sugarcane bag-worm by *Paraphylax* sp. (Hymenoptera: Ichneumonidae). *Indian Journal of Entomology* 54 (4): 484.
- Sawoniewicz, J. (1996).** *Boleslawia burakowskii* gen. nov. et sp. nov. from Poland (Hymenoptera: Ichneumonidae: Cryptinae). *Genus* 7: 493-496.
- Saxena, K.N., Reddy, K.V.S., Omolo, E.O. and Ngode, L. (1990).** Non-pesticidal approach to integrated management of sorghum stem borers for resource-limited small-scale farmers in Africa. *Proceedings: Integrated Pest Management in Tropical and Subtropical Cropping Systems '89*. 1: 175 - 198.
- Saxena, R.R. and Bhatnagar, A. (1998).** Monitoring of *Helicoverpa armigera* through light and pheromone traps and seasonal activity of their natural enemies at Bastar plateau zone, India. *Annals of Plant Protection Sciences* 6 (2): 142 - 145.
- Sayama, K. and Konishi, K. (1996).** Biological notes on *Sphecophaga vesparum* (Curtis) (Hymenoptera: Ichneumonidae) from Japan. *Japanese Journal of Entomology* 64 (4): 889 - 890.

- Scaramozzino, P.L. and Pagliano, G. (1989).** Presence in Piedmont of *Pseudogonalos hahnii* (Spinola) and notes on Trigonalidae (Hymenoptera: Trigonalidae). *Fragmenta Entomologica* 21 (2): 275 - 282.
- Schabel, H.G., Schabel, A. and Msanga, H.P. (1988).** Bioecological aspects of the mahogany defoliator *Heteronygmia dissimilis* in Morogoro, Tanzania. *Insect Science and its Application* 9 (2): 179 - 184.
- Schaefer, P.W., Fuester, R.W., Chianese, R.J., Rhoads, L.D. and Tichenor, B. Jr. (1989).** Introduction and North American establishment of *Coccygomimus disparis* (Hymenoptera: Ichneumonidae), a polyphagous pupal parasite of Lepidoptera, including gypsy moth. *Environmental Entomology* 18 (6): 1117 - 1125.
- Schaefer, P.W., Fuester, R.W., Park, J.D., Lee, B.Y., Shin, C.H., Lee, J.H. and Lee, H.P. (2002).** Gypsy moth parasitoid complex at Mt. Halla National Park, Cheju Island, Korea. *Entomological News* 113 (2): 103 - 112.
- Schaffner, U., Boeve, J.L., Gfeller, H. and Schlunegger, U.P. (1994).** Sequestration of Veratrum alkaloids by specialist *Rhadinoceraea nodicornis* Konow (Hymenoptera: Tenthredinidae) and its ecoethological implications. *Journal of Chemical Ecology* 20 (12): 3233 - 3250.
- Schalk, J.M., Shepard, B.M. and Stoner, K.A. (1993).** Response of caterpillar pests and the parasite *Diadegma insulare* to collard cultivars and a pyrethrin insecticide. *HortScience* 28 (4): 308 - 310.
- Scharer, P. (1994).** Analysis of the factors influencing the abundance of the cereal leaf beetle (*Oulema* sp., Coleoptera: Chrysomelidae). *Agrarokologie* 12: 136 pp., 9 pp.
- Schaupp, W.C Jr., Volney, W.J.A. and Waters, W.E. (1991).** Parasitoids of endemic and epidemic populations of *Choristoneura occidentalis* Freeman and *Choristoneura retiniana* (Walsingham) (Lepidoptera: Tortricidae) in southern Oregon. *Canadian Entomologist* 123 (5): 1095 - 1102.
- Schaupp, W.C. Jr. and Kulman, H.M. (1992).** Attack behavior and host utilization of *Coccygomimus disparis* (Hymenoptera: Ichneumonidae) in the laboratory. *Environmental Entomology* 21 (2): 401 - 408.
- Schembri, S. (1992).** A preliminary list of the Ichneumonidae of the Maltese islands. *Bollettino della Societa Entomologica Italiana* 124 (1): 49 - 54.
- Schmidt, J.M., Carde, R.T. and Vet, L.E.M. (1993).** Host recognition by *Pimpla instigator* F. (Hymenoptera: Ichneumonidae): preferences and learned responses. *Journal of Insect Behavior* 6 (1): 1 - 11.
- Schmidt, O., Andersson, K., Will, A. and Schuchmann-Fedderson, I. (1990).** Viruslike particle proteins from a hymenopteran endoparasitoid are related to a protein component of the immune system in the lepidopteran host. *Archives of Insect Biochemistry and Physiology* 13 (1-2): 107 - 115.

- Schmidt, O., Schuchmann-Feddersen, I. and Harris, J.R. (1989).** Role of virus-like particles in parasitoid-host interaction of insects. *Subcellular biochemistry*. Volume 15. Virally infected cells 91 - 119.
- Schmidt, O. and Theopold, U. (1990).** Molecular mimicry in a parasitoid-host relationship. *Proceedings and abstracts, Vth International Colloquium on Invertebrate Pathology and Microbial Control, Adelaide, Australia, 20-24 August 1990*. 206.
- Schmidt, O. and Theopold, U. (1994).** Virus or not: host manipulation in insect parasitic interactions. *International Journal on Endocytobiosis and Cell Research* 10 (1-2): 47 - 52.
- Schmied, A. and Fuhrer, E. (1996).** The importance of ichneumonid wasps in areas where damage is caused by *Pristiphora abietina* Christ (Hymenoptera: Tenthredinidae). *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 103 (6): 644 - 657.
- Schnee, H. (1986).** Contribution to the knowledge of the biology of some species of *Erigorgus* (Hym.: Ichneumonidae). *Entomologische Nachrichten und Berichte* 30 (6): 280 - 281.
- Schnee, H. (1989).** Revision der von Gravenhorst beschriebenen und redeskribierten Anomaloninae mit Beschreibung zweier neuerer Arten (Hymenoptera, Ichneumonidae). *Deutsche Entomologische Zeitschrift* 36: 241-266.
- Schnee, H. (1993).** Interpretation of the Anomaloninae (Hymenoptera: Ichneumonidae) described by Boie, Hartig and Ratzeburg. *Beiträge zur Entomologie* 43 (1): 39 - 51.
- Schneider, A. and Madel, G. (1992).** Fecundity and vitality of adult parasitoids following exposure to neem (*Azadirachta indica*) treated surfaces. *Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie* 8 (1-3): 273 - 278.
- Schneider, M.I. and Vinuela, E. (1999).** Evaluation of tebufenozide on immature stages of *Hyposoter didymator* (Thunberg) a parasitoid of noctuid larvae. *Proceedings, 51st international symposium on crop protection, Gent, Belgium, 4 May 1999. Part I. Mededelingen Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen, Universiteit Gent* 64 (3a): 287 - 295.
- Schneider, M.V., Driessen, G., Beukeboom, L.W., Boll, R., Van Eunen, K., Selzner, A., Talsma, J. and Lapchin, L. (2003).** Gene flow between arrhenotokous and thelytokous populations of *Venturia canescens* (Hymenoptera). *Heredity* 90 (3): 260 - 267.
- Scholler, M. (1999).** Biological control of the warehouse moth *Ephestia elutella* in bulk grain. *Agrarökologie* 35: 143 pp., 14 pp.
- Schonrogge, K. (1991).** Biology of the oak sawflies *Caliroa cinxia* Klug and *Caliroa annulipes* Klug (Hym.: Tenthredinidae) and their larval parasitoids. *Journal of Applied Entomology* 111 (4): 365 - 379.
- Schonrogge, K. and Altenhofer, E. (1992).** On the biology and larval parasitoids of the leaf-mining sawflies *Profenusa thomsoni* (Konow) and *P. pygmaea* (Konow) (Hym.: Tenthredinidae). *Entomologist's Monthly Magazine* 128: 1536 - 1539, 99 - 108.
- Schroeder, L.M. (1999).** Prolonged development time of the bark beetle predator *Thanasimus formicarius* (Col.: Cleridae) in relation to its prey species *Tomicus piniperda*



- (L.) and *Ips typographus* (L.) (Col.: Scolytidae). Agricultural and Forest Entomology 1 (2): 127 - 135.
- Schultz, P.B. and Coffelt, M.A. (1989).** Evaluation of chrysanthemum resistance to oviposition and larval feeding of the European corn borer (Lepidoptera: Pyralidae). Journal of Economic Entomology 82 (5): 1423 - 1425.
- Schumacher, R.K., Austin, A.D. and Floyd, R.B. (2000).** Parasitoids of the autumn gum moth, *Mnesampela privata* (Guenee) (Lepidoptera: Geometridae) in south-eastern Australia, with description of two new larval parasitoids. Transactions of the Royal Society of South Australia Incorporated 124 (1): 1 - 15.
- Schurian, K.G., Fiedler, K. and Maschwitz, U. (1993).** Parasitoids exploit secretions of myrmecophilous lycaenid butterfly caterpillars (Lycaenidae). Journal of the Lepidopterists' Society 47 (2): 150 - 154.
- Schwarz, M. (1995).** Revision der westpaläarktischen Arten der Gattungen *Gelis* Thunberg mit apteren Weibchen und *Thaumatogelis* Schmiedeknecht (Hymenoptera, Ichneumonidae). Teil 1. Linzer Biologische Beiträge 27: 5-105.
- Schwarz, M. and Boriani, M. (1994).** Redescription of *Gelis longulus* (Hymenoptera: Ichneumonidae), a parasitoid of *Ocnerostoma piniariellum* (Lepidoptera: Yponomeutidae). European Journal of Entomology 91 (3): 331 - 334.
- Schwarz, M. and Shaw, M. R. (1998).** Western Palaearctic Cryptinae (Hymenoptera: Ichneumonidae) in the National Museums of Scotland, with nomenclatural changes, taxonomic notes, rearing records and special reference to the British check list. Part 1. Tribe Cryptini. Entomologist's Gazette 49: 101-127.
- Schwarz, M. and Shaw, M.R. (1999).** Western Palaearctic Cryptinae (Hymenoptera: Ichneumonidae) in the National Museums of Scotland, with nomenclatural changes, taxonomic notes, rearing records and special reference to the British check list. Part 2. Genus *Gelis* Thunberg (Phygadeuontini: Gelina). Entomologist's Gazette 50 (2): 117 - 142.
- Schwarz, M. and Shaw, M.R. (2000).** Western Palaearctic Cryptinae (Hymenoptera: Ichneumonidae) in the National Museums of Scotland, with nomenclatural changes, taxonomic notes, rearing records and special reference to the British check list. Part 3. Tribe Phygadeuontini, subtribes Chiroticina, Acrolytina, Hemitelina and Gelina (excluding *Gelis*), with descriptions of new species. Entomologist's Gazette 51 (3): 147 - 186.
- Sciaky, R. (1992).** A revision of the west-Palaearctic *Selenophorina* (Coleoptera: Carabidae: Harpalinae) (XXXVI contribution to the knowledge of Coleoptera: Carabidae). Bolletino di Zoologia Agraria e di Bachicoltura 24 (1): 37 - 65.
- Seaman, A.J., Nyrop, J.P. and Dennehy, T.J. (1990).** Egg and larval parasitism of the grape berry moth (Lepidoptera: Tortricidae) in three grape habitats in New York. Environmental Entomology 19 (3): 764 - 770.
- Sedivy, J. (1996).** *Xorides sinoxyli* sp.nov. from Israel (Hymenoptera: Ichneumonidae). Entomological Problems 27 (2):167 - 169.

- Seker, D.A. and Yanikoglu, A. (1999).** The changes in glycogen levels of *Pimpla turionellae* L. (Hymenoptera: Ichneumonidae) in the conditions of starvation, feeding, parasitism and aging. Turkish Journal of Zoology 23: suppl, 289 - 296.
- Sekhar, P.R. and Rai, P.S. (1989).** Studies on the biology and occurrence of natural enemies of safflower caterpillar, *Prospalta conducta* Walker (Lepidoptera: Noctuidae) in Karnataka. Journal of Research APAU 17 (1): 19 - 22.
- Sekita, N., Watanabe, T. and Yamada, M. (1996).** Population ecology of *Osmia cornifrons* (Hymenoptera: Megachilidae) in natural habitats. Bulletin of the Aomori Apple Experiment Station 29: 17 - 36.
- Selfa, J. (1994).** Contribution to the knowledge of the Ichneumoninae (Hymenoptera: Ichneumonidae) from the Iberian Peninsula. Zoologische Mededelingen 68: 15-25, 197 - 202.
- Selfa, J., Bordera, S. and Jimenez, R. (1988).** New contributions to the knowledge of the subfamily Ichneumoninae in Spain. II. Tribe Ichneumonini (Hymenoptera: Ichneumonidae). Eos Madrid 64 (1): 243 - 255.
- Selfa, J., Bordera, S. and Jimenez, R. (1990).** Two new Spanish species of the genus *Ichneumon* Linnaeus (Hymenoptera: Ichneumonidae). Nouvelle Revue d'Entomologie 7 (3): 267 - 270.
- Selfa, J. and Bordera, S. (1993).** Study of Peninsular Ichneumoninae of the National Museum of Natural Sciences. Phaeogenini (Hymenoptera: Ichneumonidae). Boletín de la Asociación Española de Entomología 17 (2): 37 - 47.
- Selfa, J. and Diller, E. (1994).** Illustrated key to the western Palearctic genera of Phaeogenini (Hymenoptera, Ichneumonidae, Ichneumoninae). Entomofauna 15: 237-251.
- Selfa, J. and Bordera, S. (1996).** The species of Ichneumoninae of the Universitat de Valencia II: *Ichneumoninae Stenopneusticae* (Hymenoptera: Ichneumonidae). Nouvelle Revue d'Entomologie 13 (3): 235 - 248.
- Selfa, J. and Anento, J.L. (1999).** New data on the Portuguese Ichneumoninae (Hymenoptera: Ichneumonidae). Nouvelle Revue d'Entomologie 16 (1): 65 - 66.
- Selvasundaram, R. and Muraleedharan, N. (1989).** Observations on the larval parasitoids of *Homona coffearia* (Nietner) (Tortricidae: Lepidoptera) in a tea garden at the Annamalais. Journal of Plantation Crops. 16: Supplement, 185 - 188.
- Semevskii, F.N. (1993).** Large-scale migrations of the green oak tortrix *Tortrix viridana* and of its specialized parasitoid *Dirophanes invisor*. Zoologicheskii Zhurnal 72 (4): 74 - 79.
- Sengonca, C. and Leisse, N. (1988).** Occurrence and importance of pupal parasitoids of the grape moth *Eupoecilia ambiguella* Hb. in the Ahr Valley. Journal of Applied Entomology 106 (2): 173 - 176.
- Sengonca, C. and Peters, G. (1991).** Occurrence of indigenous parasitoids of the important cabbage pests in the Cologne-Bonn production region. Zeitschrift für Angewandte Zoologie 78 (1): 91 - 99.

- Serini, G.B. (1990).** Parasitoids of *Parectopa robiniella* Clemens and *Phyllonorycter robiniellus* (Clemens) (Lepidoptera: Gracillariidae). *Bollettino di Zoologia Agraria e di Bachicoltura* 22 (2): 139 - 149.
- Seufert, P. and Fiedler, K. (1999).** Myrmecophily and parasitoid infestation of South-East Asian lycaenid butterfly larvae. *Ecotropica* 5 (1): 59 - 64.
- Shah, R.H., Patel, J.R. and Patel, N.C. (2001).** Occurrence of natural enemies of aonla hairy caterpillar, *Selepa celtis* Moore infesting aonla orchard. *Indian Journal of Entomology* 63 (2): 193 - 195.
- Shamuratova, N.G. (2002).** Natural enemies of lucerne pests in the Aral region. *Zashchita I Karantin Rastenii* 10: 38 - 39.
- Shanklin, D.R., Johnson, D.W. and Townsend, L.H. (1998).** Survey of parasitoids of the European corn borer (Lepidoptera: Pyralidae) in Southwestern Kentucky. *Journal of Entomological Science* 33 (3): 256 - 260.
- Shao, J.W., Meng, F.R., Zhou, H.Y., Sun, Z.Y., Li, G.L., Jia, R.F., Wang, X.C., Wang, S.J. and Sun, J.H. (1994).** A study on the bionomics of *Laspeyresia zebeana*. *Journal of Northeast Forestry University* 22 (5):100 - 104.
- Sharifi, S. (1968).** Monography of a parasitic hymenopter, *Xorides corcyrensis* (Hymenoptera: Ichneumonidae). *Ibid.*, 195-201 pp.
- Sharkey, M.J. and Wahl, D.B. (1992).** Cladistics of the Ichneumonoidea (Hymenoptera). *Journal of Hymenoptera Research* 1 (1):15 - 24.
- Sharma, D.D. and Agrawal, M.L. (1988).** Studies on the biology and immature stages of litchi fruit borer *Conopomorpha cramerella* (Snellen) (Lepidoptera: Gracillariidae). *Journal of Research Rajendra Agricultural University* 6 (1-2): 84 - 87.
- Sharma, D.R. and Batra, R.C. (1989).** Parasitoids of citrus leaf roller, *Psorosticha zizyphi* Stain. and influence of abiotic factors on the parasitization. *Journal of Insect Science* 2 (2): 162 - 164.
- Sharma, H.C., Sullivan, D.J. and Bhatnagar, V.S. (2002).** Population dynamics and natural mortality factors of the Oriental armyworm, *Mythimna separata* (Lepidoptera: Noctuidae), in South-Central India. *Crop Protection* 21 (9): 721 - 732.
- Sharma, K.C. and Bhalla, O.P. (1992).** Studies on the life of *Metasyrphus corollae* (Fab.) a predator of the cabbage aphid (*Brevicoryne brassicae* L.) on cauliflower seed crop. *Entomon* 17 (1 & 2): 49 - 53.
- Sharma, S.K., Devi, N. and Raj, D. (1999).** Bionomics and parasitization of diamondback moth, *Plutella xylostella* L. (Lep.: Plutellidae). *Journal of Entomological Research* 23 (4): 309 - 314.
- Sharov, A.A., Wagner, M.R. and Raffa, K.F. (1993).** Biology and population dynamics of the common pine sawfly, *Diprion pini* L., in Russia. Sawfly life history adaptations to woody plants 409 - 429.

- Shashidhar, V., Basalingappa, S. and Lingappa, S. (1998).** Parasitoids of the greater wax moth, *Galleria mellonella* (L.) in Dharwad, Karnataka, India. *Insect Environment* 3 (4): 100.
- Shaw, M. R. (1976).** Noctuid (Lep.) larvae consuming ichneumonoid (Hym.) larvae. *Entomologist's Monthly Magazine* [1975] 111: 188.
- Shaw, M. R. and Askew, R. R. (1976).** Ichneumonoidea (Hymenoptera) parasitic upon leaf-mining insects of the orders Lepidoptera, Hymenoptera and Coleoptera. *Ecological Entomology* 1: 127-133.
- Shaw, M. R. (1977).** Taxonomic and biological notes on the European species of *Benjaminia* Viereck (Hym., Ichneumonidae). *Nouvelle Revue d'Entomologie* 7: 85-89.
- Shaw, M. R. (1977).** On the distribution of some satyrid (Lep.) larvae at a coastal site in relation to their ichneumonid (Hym.) parasite. *Entomologist's Gazette* 28: 133-134.
- Shaw, M. R. (1978).** The status of *Trogus lapidator* (F.) (Hymenoptera: Ichneumonidae) in Britain, a parasite of *Papilio machaon* L. *Entomologist's Gazette* 29: 287-288.
- Shaw, M. R. (1984).** Foodplants and parasites (Hymenoptera: Ichneumonidae) of *Choreutis pariana* (Clerck) (Lepidoptera: Choreutidae) in Edinburgh. *Entomologist's Gazette* 35: 41-44.
- Shaw, M. R. (1986).** *Coleocentrus excitator* (Poda) (Hymenoptera: Ichneumonidae) new to Britain. *Entomologist's Gazette* 37: 221-224.
- Shaw, M. R. (1989).** A host record for *Stilbops limneriaeformis* Schmiedeknecht (Hymenoptera: Ichneumonidae, Stilbopinae) in Scotland. *Entomologist's Gazette* 40: 5-6.
- Shaw, M. R. (1991).** *Phrudus badensis* Hilpert (Hym., Ichneumonidae) new to Britain. *Entomologist's Monthly Magazine* 127: 157-158.
- Shaw, M.R. (1993).** Species of *Mesochorus* (Hymenoptera: Ichneumonidae) reared as hyperparasitoids of Lepidoptera via koinobiont ectoparasitoid Tryphoninae (Ichneumonidae). *Entomologist's Gazette* 44 (3): 181 - 182.
- Shaw, M.R. (1999).** Gregarious development in endoparasitic koinobiont Ichneumonidae (Hymenoptera). *Entomologist's Gazette* 50 (1): 55 - 56.
- Shaw, M.R. (2001).** Interactions between adults of some species of *Netelia* Gray (Hymenoptera: Ichneumonidae: Tryphoninae) and their caterpillar hosts (Lepidoptera). *Journal of Hymenoptera Research* 10 (1): 101 - 111.
- Shaw, M. R. (2004).** Notes on the biology of *Lycorina triangulifera* Holmgren (Hymenoptera: Ichneumonidae: Lycorininae). *Journal of Hymenoptera Research* 13: 302 – 308.
- Shaw, M.R. & Bennett, F.D. (2001).** Host prepupal mummification by *Colpognathus Wesmael* (Hymenoptera: Ichneumonidae, Ichneumoninae). *Entomologist's Gazette* 52: 201-203.
- Shaw, M.R. and Bailey, M. (1991).** Parasitoids (Hymenoptera: Braconidae: Ichneumonidae: Pteromalidae) and notes on the biology of the fern-boring sawfly *Heptamelus ochroleucus*

- (Stephens) (Hymenoptera: Tenthredinidae) in the English Lake District. *Entomologist* 110 (3): 103 - 109.
- Shaw, M. R. and Aeschlimann, J.-P. (1994).** Host ranges of parasitoids (Hymenoptera: Braconidae and Ichneumonidae) reared from *Epermenia chaerophyllella* (Goeze) (Lepidoptera: Epermeniidae) in Britain, with description of a new species of *Triclistus* (Ichneumonidae). *Journal of Natural History* 28: 619-629.
- Shaw, M. R. and Horstmann, K. (1997).** An analysis of host range in the *Diadegma nanus* group of parasitoids in Western Europe, with a key to species (Hymenoptera: Ichneumonidae: Campopleginae). *Journal of Hymenoptera Research* 6: 273-296.
- Shaw, M.R. and Wahl, D.B. (1989).** The biology, egg and larvae of *Acaenitus dubitator* (Panzer) (Hymenoptera: Ichneumonidae: Acaenitinae). *Systematic Entomology* 14 (1): 117 - 125.
- Shaw, M.R. and Kasparyan, D.R. (2002).** Some British records of *Olethrodotis modestus* (Gravenhorst) (Hymenoptera: Ichneumonidae: Ctenopelmatinae). *Entomologist's Record* 114: 137-139.
- Shaw, M.R. and Kasparyan, D.R. (2003).** Some genera of British and European Mesoleiini (Hym., Ichneumonidae: Ctenopelmatinae) in the National Museums of Scotland, including a new species of *Mesoleius* and a further twenty species new to Britain. *Entomologist's Monthly Magazine* 139: 17-28.
- Shaw, M.R., Kasparyan, D.R. and Fitton, M.G. (2003).** Revision of the British checklist of Ctenopelmatini (Hymenoptera: Ichneumonidae, Ctenopelmatinae). *Entomologist's Gazette* 54: 137-141.
- She, G. and He, R.R. (1988).** Field survey on parasites of rice stem borer in Chengdu, Sichuan province. *Chinese Journal of Biological Control* 4 (1): 6 - 9.
- Sheehan, W. and Hawkins, B.A. (1991).** Attack strategy as an indicator of host range in metopiine and pimpline Ichneumonidae (Hymenoptera). *Ecological Entomology* 16 (1): 129 - 131.
- Shelby, K.S., Adeyeye, O.A., Okot-Kotber, B.M. and Webb, B.A. (2000).** Parasitism-linked block of host plasma melanization. *Journal of Invertebrate Pathology* 75 (3): 218 - 225.
- Shelby, K.S., Cui, L. and Webb, B.A. (1998).** Polydnavirus-mediated inhibition of lysozyme gene expression and the antibacterial response. *Insect Molecular Biology* 7 (3): 265 - 272.
- Shelby, K.S. and Webb, B.A. (1994).** Polydnavirus infection inhibits synthesis of an insect plasma protein, arylphorin. *Journal of General Virology* 75 (9): 2285 - 2292.
- Shelby, K.S. and Webb, B.A. (1997).** Polydnavirus infection inhibits translation of specific growth-associated host proteins. *Insect Biochemistry and Molecular Biology* 27 (3): 263 - 270.
- Shelby, K.S. and Webb, B.A. (1999).** Polydnavirus-mediated suppression of insect immunity. *Journal of Insect Physiology* 45 (5): 507 - 514.

- Shelton, A.M., Wilsey, W.T., Hoebeke, E.R. and Schmaedick, M.A. (2002).** Parasitoids of cabbage Lepidoptera in Central New York. *Journal of Entomological Science* 37 (3): 270 - 271.
- Shelton, A.M., Xu, J.X. and Cheng, X.N. (2001).** Comparison of *Diadegma insulare* (Hymenoptera: Ichneumonidae) and *Microplitis plutellae* (Hymenoptera: Braconidae) as biological control agents of *Plutella xylostella* (Lepidoptera: Plutellidae): field parasitism. *Journal of Economic Entomology* 94 (1): 14 - 20.
- Sheng, J.K. and Lu, H.L. (1988).** Notes on the survey of the parasites of *Oulema oryzae* (Kuwayama) (Coleoptera: Chrysomelidae) in Jiangxi province. *Natural Enemies of Insects* 10 (1): 61.
- Sheng, M.L. (1990).** New records of Hymenoptera parasitizing wood-boring insects from China. *Entomotaxonomia* 12 (1): 56.
- Sheng, M.L. (2000).** The genus *Lissonota* (Hymenoptera: Ichneumonidae) from North China. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique, Entomologie* 70: 189 - 197.
- Sheng, M.L., Wu, L.Z., Wu, J.X. and He, Z.M. (1999).** A new species of the genus *Diaparsis* (Hymenoptera: Ichneumonidae) parasitizing *Lema decempunctata* with a new record from China. *Scientia Silvae Sinicae* 35 (1):66 - 68.
- Sheng, M.L., Zhang, Q.H. and Chen, G.F. (1998).** A new species of genus *Aptesia* parasitic on *Diprion jingyuanensis* (Hymenoptera: Ichneumonidae: Diprionidae). *Acta Entomologica Sinica* 41 (3): 316 - 318.
- Sheng, M.L. and Pei, H.C. (2002).** Two new species of genus *Cosmoconus* Foerster from Henan (Hymenoptera: Ichneumonidae). *Acta Entomologica Sinica* 45: Supplement, 96 - 98.
- Sheng, M.L. and Sun, S.P. (2002).** Chinese *Kristotomus* Mason with description of a new species (Hymenoptera: Ichneumonidae). *Acta Entomologica Sinica*. 45: Supplement, 93 - 95.
- Sheng, M.L. and Wei, D.Y. (1999).** A new species of genus *Himerta* (Hymenoptera: Ichneumonidae) from Hebei Province of China. *Journal of Northeast Forestry University* 27 (3): 81 - 82.
- Sheng, M.L. and Xu, G.T. (1997).** Three new species and two new host records of genus *Eriborus* Foerster (Hymenoptera: Ichneumonidae) from China. *Entomologia Sinica* 4 (2): 139 - 143.
- Shepard, B.M. and Barrion, A.T. (1998).** Parasitoids of insects associated with soybean and vegetable crops in Indonesia. *Journal of Agricultural Entomology* 15 (3): 239 - 272.
- Shetgar, S.S., Bilapate, G.G., Patil, V.V. and Londhe, G.M. (1990).** A note on the natural enemies of Bihar hairy caterpillar, *Spilosoma obliqua* (Walker). *Indian Journal of Entomology* 52 (1): 158.
- Shibata, E. (1994).** Population studies of *Callidiellum rufipenne* (Coleoptera: Cerambycidae) on Japanese cedar logs. *Annals of the Entomological Society of America* 87 (6): 836 - 841.

- Shojai, M. (1986).** Resultats de l'étude faunistiques des hymenopteres parasites (Trebrants) en Iran et l'importance de leur utilisation dans la lutte biologique. Proceeding I. Plant Prot. Cong. Iran; Univ. Tehran, 25-35 pp.
- Shojai, M. (1989).** Entomology (Ethology, social life and natural enemies) (biological control). Vol. III. Second ed. Univ. Tehran Publ., 406 pp (p. 341).
- Shojai, M., Esmaili, M., Ostovan, H., Khodaman, A., Daniali, M., Hosseini, M., Assadi, Y., Sadighfar, M., Korosh-Najad, A., Nasrollahi, A., Labbafi, Y., Azma, M., Ghavam, F. and Honarbakhsh, S. 2000.** Integrated pest management of codling moth and other important pests of pomoidea fruit trees. J. of Agricultural Sciences, Islamic Azad University 6(2): 15 – 46.
- Shojai, M., Ostovan, H., Hosseini, M., Sadighfar, M., Khodaman, A., Labbafi, Y., Nasrollahi, A., Ghavam, F. and Honarbakhsh, S. 2002.** Biocenotic potentials of apple orchards IPM in organic crop production program. J. of Agricultural Sciences, Islamic Azad University 8(1): 1 – 28.
- Short, M.W., Schmidt, S. and Lukacs, Z. (2002).** Parasitisation rates of some parasitoids (Hymenoptera: Ichneumonidae) of the autumn gum moth (Lepidoptera: Geometridae). Australian Entomologist 29 (2): 69 - 72.
- Shrivastava, S.K. and Yadav, K.P. (1991).** Distribution of *Heliothis armigera* (Hub.) and its biocontrol agents in Chhattisgarh, Madhya Pradesh. Agricultural Science Digest Karnal 11 (2): 107 - 109.
- Shu, S.Q. and Jones, R.L. (1993).** Evidence for a multicomponent sex pheromone in *Eriborus terebrans* (Gravenhorst) (Hym.: Ichneumonidae), a larval parasitoid of the European corn borer. Journal of Chemical Ecology 19 (11): 2563 - 2576.
- Shukla, B.C., Kaushik, U.K. and Gupta, R. (1999).** Activity of *Campoletis chlorideae* on *Helicoverpa armigera* in chickpea crop as influenced by intercropping with coriander under rice based cropping system. Journal of Applied Zoological Researches 10 (1): 32 - 35.
- Sidlyarevich, V., Lameko, V., Dabrowski, Z.T. and Narkiewicz-Jodko, J. (2000).** Some ecological peculiarities of the diamond back moth (*Plutella maculipennis* Curt.) population development. Proceedings of the international symposium on Lepidoptera pests in vegetables, Skierniewice, Poland, 2-3 December, 1998. Vegetable Crops Research Bulletin 52: 31 - 38.
- Sidor, C. (1989).** Contribution to the investigation of viral infection of larvae of the European sawfly (*Neodiprion sertifer*) in different instars with reference to their insect parasites. Zastita Bilja 40 (1): 57 - 66.
- Sieglaff, D.H., Mitchell, E.R. and Hu, G.Y. (1998).** Evaluation of rearing methods for *Diadegma insulare* (Hymenoptera: Ichneumonidae), an endoparasitoid of the diamondback moth (Lepidoptera: Plutellidae). Florida Entomologist 81 (4): 578 - 582.

- Silva, F.M.A., Fowler, H.G. and Lemos, R.N.S. (1997).** Parasitism of fall armyworm, *Spodoptera frugiperda* (Smith), in the Triangulo Mineiro region, MG. *Anais da Sociedade Entomologica do Brasil* 26 (2): 235 - 241.
- Silva De, M.T.B. (1993).** Control of the soyabean caterpillar (*Anticarsia gemmatalis* Hubner, 1818 - Lepidoptera: Noctuidae). IV. Natural biological control. *Ciencia Rural* 23 (2): 127 - 132.
- Silveira Da, J.C.F., Sasaki, E.T., Forner, M.A., Honda, M.S. and Calafiori, M.H. (1987).** Occurrence of parasitoids of *Spodoptera frugiperda* (J.E. Smith, 1797) on maize in Espirito Santo do Pinhal, SP. *Ecosystema*, publ. 1988, 12: 41 - 44.
- Silvie, P. (1993).** The parasitoids of *Syllepte derogata* (Fabricius, 1775) (Lepidoptera: Crambidae) in Togo. *Journal of African Zoology* 107 (4): 363 - 372.
- Simandl, J. (1992).** Parasitoids (Ichneumonidae) of an endemic population of pine sawflies (Hym.: Diprionidae). *Journal of Applied Entomology* 114 (5): 480 - 483.
- Simandl, J. (2000).** Bionomy and distribution of *Saperda perforata* (Coleoptera: Cerambycidae) in South Bohemia. *Sbornik Jihoceskeho Muzea v Ceskych Budejovicich, Prirodni Vedy* 40 (1-2): 55 - 57.
- Simbolotti, G. and Van Achterberg, C. (1992).** Revision of the West Palaearctic species of the genus *Bassus Fabricius* (Hymenoptera: Braconidae). *Zoologische Verhandelingen* 281: 80 pp.
- Simchuk, P.A. (1988).** Parasitoids of the potato moth *Phthorimaea operculella* Zeller (Lepidoptera: Gelechiidae) in the Crimea. *Entomologicheskoe Obozrenie* 67 (3): 510 - 513.
- Sime, K. (2002).** Chemical defence of *Battus philenor* larvae against attack by the parasitoid *Trogus pennator*. *Ecological Entomology* 27 (3): 337 - 345.
- Sime, K.R. (2000).** Two new records of pimpline ichneumonids attacking *Battus philenor* (Linnaeus) (Lepidoptera: Papilionidae). *Journal of Hymenoptera Research* 9 (1): 210 - 212.
- Sime, K.R. (2002).** Experimental studies of the host-finding behavior of *Trogus pennator*, a parasitoid of swallowtail butterflies. *Journal of Chemical Ecology* 28 (7): 1377 - 1392.
- Sime, K.R. and Brower, A.V.Z. (1998).** Explaining the latitudinal gradient anomaly in ichneumonid species richness: evidence from butterflies. *Journal of Animal Ecology* 67 (3): 387 - 399.
- Sime, K.R. and Wahl, D.B. (1998).** Taxonomy, mature larva, and observations on the biology of *Gnamptopelta obsidianator* (Brulle) (Hymenoptera: Ichneumonidae: Ichneumoninae). *Journal of Hymenoptera Research* 7 (2): 157 - 164.
- Sime, K.R. and Wahl, D.B. (2002).** The cladistics and biology of the *Callajoppa* genus-group (Hymenoptera: Ichneumonidae: Ichneumoninae). *Zoological Journal of the Linnean Society* 134 (1): 1 - 56.
- Singh, A.P. (1998).** Bionomics of cone worm, *Dioryctria abietella* Denis and Schiff. on West Himalayan spruce, *Picea smithiana*. *Annals of Forestry* 6 (1): 84 - 88.



- Singh, S.B., Yadav, S.S., Singh, S. and Dixit, A. (2002).** Naturally occurring key mortality factors of top borer (*Scirpophaga excerptalis* Wlk.) in sugarcane ratoon in U.P. Proceedings of the 64<sup>th</sup> Annual Convention of the Sugar Technologists' Association of India, Cochin Kerala, India, 17<sup>th</sup> - 19<sup>th</sup> August 2002. 69 - 84.
- Singh, S.P. (1991).** Natural enemies of *Papilio* spp. Indian Journal of Horticulture 48 (3): 237 - 242.
- Singh, S.P. and Singh, Y. (1991).** Parasitoids of *Heliothis armigera* (Hb.) on pigeonpea in Delhi. Indian Journal of Entomology 53 (1): 168.
- Singh, Y.P. and Kumar, V. (1991).** Record of new larval parasitoids of *Lymantria beatrix* Stoll. (Lepidoptera: Lymantriidae). Journal of the Bombay Natural History Society 88 (3): 43.
- Singh, Y.P. and Singh, P.P. (2002).** Natural parasites and extent of parasitism to shoot and fruit borer (*Leucinodes orbonalis*) of brinjal (*Solanum melongena*) at medium high altitude hills of Meghalaya. Indian Journal of Entomology 64 (2): 222 - 226.
- Sirot, E. (1996).** The pay-off from superparasitism in the solitary parasitoid *Venturia canescens*. Ecological Entomology 21 (3): 305 - 307.
- Sivaprakasam, N. (1997).** Parasitoids of fruit-borer (*Helicoverpa armigera*) of tomato (*Lycopersicon esculentum*) in Tamil Nadu. Indian Journal of Agricultural Sciences 67 (4): 177.
- Sivaprakasam, N., Balasubramanian, G., Venkatesan, S., Jeyaraman, V. and Narayanan, A. (1986).** Effect of host plants on parasitism by parasitoids of *Heliothis armigera* Hubner. Madras Agricultural Journal 73 (8): 468 - 470.
- Skatulla, U. and Gmach, T. (1989).** Notes on the determination of parasitism levels in pupae of the pine beauty moth in relation to pest forecasting. Anzeiger fur Schadlingskunde, Pflanzenschutz, Umweltschutz 62 (4): 65 - 67.
- Skillen, E.L., Pickering, J. and Sharkey, M.J. (2000).** Species richness of the campopleginae and ichneumoninae (Hymenoptera: Ichneumonidae) along a latitudinal gradient in Eastern North American old-growth forests. Environmental Entomology 29 (3): 460 - 466.
- Skinner, M., Parker, B.L., Wallner, W.E., Odell, T.M., Howard, D. and Aleong, J. (1993).** Parasitoids in low-level populations of *Lymantria dispar* (Lep.: Lymantriidae) in different forest physiographic zones. Entomophaga 38 (1): 15 - 29.
- Skovgard, H. and Jespersen, J.B. (1999).** Activity and relative abundance of hymenopterous parasitoids that attack puparia of *Musca domestica* and *Stomoxys calcitrans* (Diptera: Muscidae) on confined pig and cattle farms in Denmark. Bulletin of Entomological Research 89 (3): 263 - 269.
- Skovgard, H. and Jespersen, J.B. (2000).** Seasonal and spatial activity of hymenopterous pupal parasitoids (Pteromalidae and Ichneumonidae) of the house fly (Diptera: Muscidae) on Danish pig and cattle farms. Environmental Entomology 29 (3): 630 - 637.

- Slovak, M. (1987).** Cannibalism in larvae of the cabbage moth in relation to parasitism by the ichneumonid *Exetastes cinctipes* Retz. *Biologia Bratislava* 42 (10): 955 - 958.
- Slovak, M. (1987).** Depth of entering the soil by larvae of the cabbage moth in relation to parasitism by the ichneumonid *Exetastes cinctipes* Retz. *Biologia Bratislava* 42 (6): 597 - 601.
- Slovak, M. (1987).** Influence of temperature and host's nutritional conditions on development of the immature stages of *Exetastes cinctipes* Retz. (Hym.: Ichneumonidae). *Biologia Bratislava* 42 (6): 587 - 596.
- Slovak, M. (1988).** Breaking diapause of the ichneumonid *Exetastes cinctipes* by low temperature. *Biologia, Czechoslovakia* 43 (6): 549 - 554.
- Smith, G.K.M. (1996).** Insect parasites control three introduced sawflies. *Frontline, Technical Note Canadian Forest Service Sault Ste. Marie* 24: 4 pp.
- Smith, J.W. Jr., Rodriguez Del Bosque, L.A. and Agnew, C.W. (1990).** Biology of *Mallochia pyralidis* (Hymenoptera: Ichneumonidae), an ectoparasite of *Eoreuma loftini* (Lepidoptera: Pyralidae) from Mexico. *Annals of the Entomological Society of America* 83 (5): 961 - 966.
- Soczynski, G. and Gantner, M. (1998).** Parasitoids emerged from pupal cocoons of *Oulema gallaeciana* (Heyden) (Col.: Chrysomelidae) in the Lublin region. *Progress in Plant Protection*. 38 (2): 450 - 452.
- Sohati, P.H., Musonda, E.M. and Mukanga, M. (2001).** Distribution of cereal stemborers in Zambia and release of *Cotesia flavipes* Cameron, an exotic natural enemy of *Chilo partellus* (Swinhoe). *Insect Science and its Application* 21 (4): 311 - 316.
- Solbreck, C. (2000).** Ecology and biology of *Euphranta connexa* (Fabr.) (Diptera: Tephritidae) - a seed predator on *Vincetoxicum hirundinaria* Med. (Asclepiadaceae). *Entomologisk Tidskrift* 121 (1-2): 23 - 30.
- Soldevila, A., Webb, B.A. and Cui, L.W. (1997).** Expression and hemocyte-targeting of a *Campoletis sonorensis* polydnavirus cysteine-rich gene in *Heliothis virescens* larvae. *Archives of Insect Biochemistry and Physiology* 36 (4): 251 - 271.
- Soldevila, A.I., Heuston, S. and Webb, B.A. (1997).** Purification and analysis of a polydnavirus gene product expressed using a poly-histidine baculovirus vector. *Insect Biochemistry and Molecular Biology* 27 (3): 201 - 211.
- Soldevila, A.I., Webb, B.A. and Cui, L.W. (2000).** Relationships between polydnavirus gene expression and host range of the parasitoid wasp *Campoletis sonorensis*. *Journal of Insect Physiology* 46 (10): 1397 - 1407.
- Sole, M., Cano, J. and Guarro, J. (2002).** Molecular phylogeny of *Amauroascus*, *Auxarthron*, and morphologically similar onygenalean fungi. *Mycological Research* 106 (4): 388 - 396.
- Solsoloy, A.D., Domingo, E.O., Bilgera, B.U., Solsoloy, T.S., Bugawan, H.S. and Barluado, Z.D. (1994).** Occurrence, mortality factors and within-plant distribution of

- bollworm, *Helicoverpa armigera* (Hubn.) on cotton. Philippine Journal of Science 123 (1): 9 - 20.
- Song, M.J., Zhou, N.L., Zai, Y.D. and Zhang, X. (1994).** A preliminary study on *Euzoper pyriella* Yong. Plant Protection 20 (3): 13 - 15.
- Songa, J.M., Overholt, W.A., Okello, R.O. and Mueke, J.M. (2002).** Control of lepidopteran stemborers in maize by indigenous parasitoids in semi-arid areas of Eastern Kenya. Biological Agriculture and Horticulture 20 (1):77 - 90.
- Sontgen, J.M. and Sengonca, C. (1988).** Observations on the incidence of parasitoids of the apple clearwing moth, *Synanthedon myopaeformis* (Borkh.), in Nordrhein. Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie 6 (1-3): 262 - 266.
- Sood, A.K. and Bhalla, O.P. (1996).** Ecological studies on the cabbage white butterfly in the mid-hills of Himachal Pradesh. Journal of Insect Science 9 (2): 122 - 125.
- Sourakov, A. and Itchell, E.R. (2002).** Sex allocation in progeny of *Diadegma insulare* (Hymenoptera: Ichneumonidae). Journal of Entomological Science 37 (2): 210 - 211.
- Spitzer, K. and Van San, N. (1993).** Isolated populations of the winter moth, *Operophtera brumata* (Lepidoptera: Geometridae), their heavy metal content and parasitism. European Journal of Entomology 90 (3):311 - 321.
- Srinivas, P.R. (1989).** Extent of parasitism of gram-pod borer (*Heliothis armigera*) by ichneumonid larval parasites. Indian Journal of Agricultural Sciences 59 (6): 377 - 378.
- Srinivas, P.R. and Jayaraj, S. (1989).** Record of natural enemies of *Heliothis armigera* from Coimbatore district, Tamil Nadu. Journal of Biological Control 3 (1): 71 - 72.
- Srinivasan, K., Moorthy, P.N.K. and Devi, K.S. (1995).** Parasitism and progeny sex ratio of *Diadegma semiclausum* Hellen - a larval parasitoid of diamondback moth. Pest Management in Horticultural Ecosystems 1 (1): 9 - 13.
- Stary, B., Bezdecka, P., Capek, M., Stary, P., Zeleny, J. and Sedivy, J. (1988).** Atlas of insects beneficial to forest trees. Volume 2. Elsevier Science Publishers; Amsterdam; Netherlands, 100 pp.
- Stasiak, K., Demattei, M.V., Federici, B.A. and Bigot, Y. (2000).** Phylogenetic position of the *Diadromus pulchellus* ascovirus DNA polymerase among viruses with large double-stranded DNA genomes. Journal of General Virology 81 (12): 3059 - 3072.
- Stejskal, V., Lukas, J., Zd'arkova, E., Wakefield, M., Lukas, J. and Hubert, J. (2002).** Spatial arrangement of *Venturia canescens* and *Ephestia kuehniella* in the extremely infested pasta-producing factory: a case history. COST Action 842, 1999-2004. Biological control of pest insects and mites with special reference to entomophthorales. Proceedings of the Second Meeting of Working Group 4: Bio control of arthropod pests in the stored products, Prague, Czech Republic, 30-31st May 2002. 116 - 117.
- Stoll, G. (1996).** PLITS: Wirksamkeit von pflanzlichen Substanzen in der integrierten Bekämpfung von *Plutella xylostella* (L.) auf den Philippinen. Faculty of Agricultural Sciences, Hohenheim University; Germany, 188 pp.

- Stoltz, B. and Whitfield, J.B. (1992).** Viruses and virus-like entities in the parasitic Hymenoptera. *Journal of Hymenoptera Research* 1 (1): 125 - 139.
- Stoltz, D. and Makkay, A. (2000).** Co-replication of a reovirus and a polydnavirus in the ichneumonid parasitoid *Hyposoter exiguae*. *Virology New York* 278 (1): 266 - 275.
- Stoltz, D.B. (1990).** Evidence for chromosomal transmission of polydnavirus DNA. *Journal of General Virology* 71 (5): 1051 - 1056.
- Stouthamer, R., Luck, R.F. and Werren, J.H. (1992).** Genetics of sex determination and the improvement of biological control using parasitoids. *Environmental Entomology* 21 (3): 427 - 435.
- Stouthamer, R. and Bigler, F. (1991).** Genetic problems caused by sex determination in the rearing of parasitoids. Fifth Workshop of the IOBC Global Working Group on Quality Control of Mass Reared Arthropods 38 - 46.
- Strand, M.R. and Pech, L.L. (1995).** Immunological basis for compatibility in parasitoid-host relationships. *Annual Review of Entomology* 40: 31 - 56.
- Streibert, H.P. and Kaeding, D. (1994).** Use of Pegasus R (diafenthiuron) against diamondback moth (*Plutella xylostella*) and other lepidopterous pests in cruciferous crops. Proceedings Brighton Crop Protection Conference, Pests and Diseases, Brighton, UK, 21-24 November 1994, vol 2. 743 - 748.
- Streito, J.C. and Nibouche, S. (1997).** First observations on the parasitoids associated with lepidopterous pests of cotton in Burkina Faso. *Entomophaga* 42 (4): 543 - 557.
- Stuber, R. and Dickler, E. (1988).** Studies on the biology and behaviour of the apple clearwing *Synanthedon myopaeformis* (Bork.) (Lepidoptera: Sesiidae) as a basis for its control using the confusion method. *Mitteilungen aus der Biologischen Bundesanstalt für Land und Forstwirtschaft Berlin Dahlem* 241: 144 pp., 14 pp.
- Su, S.Y. and Zhou, B. (1988).** Studies on bionomics and control of *Phloeosinus sinensis*. *Scientia Silvae Sinicae* 24 (2): 239 - 242.
- Suasa-Ard, W. and Allsopp, P.G. (2000).** *Chilo tumidicostalis* (Hampson) (Lepidoptera: Pyralidae) and its natural enemies in Thailand. Sugarcane pest management in the New Millennium. 4<sup>th</sup> Sugarcane Entomology Workshop International Society of Sugar Cane Technologists, Khon Kaen, Thailand, 7 - 10 February 2000. 10 - 16.
- Subbiah, K. (1988).** New records of insect parasitoids of leaf-roller and flushworm of tea. *Current Science* 57 (12): 686.
- Sudheendrakumar, V.V. (1993).** Notes on hymenopteran parasites of *Eutectona machaeralis* recorded from Nilambur, Kerala. *Indian Forester* 119 (6): 510 - 511.
- Suganthi, M., Rao, G.V.R. and Kumar, S.T. (2002).** Relative efficacy of IPM strategies of chickpea on soil inhabiting and aerial natural enemies. *Legume Research* 25 (1): 66 - 68.
- Sugiura, S. and Osawa, N. (2001).** Parasitoid community associated with the leafroller *Eudemis gyrotis* (Lepidoptera: Tortricidae) on a bayberry, *Myrica rubra* (Myricaceae). *Entomological Science* 4 (1): 25 - 34.

- Sugiura, S. and Osawa, N. (2002).** Temporal response of parasitoids to the density of the leafroller *Eudemis gyrotis* (Lepidoptera: Tortricidae) on bayberry *Myrica rubra* (Myricaceae). *Environmental Entomology* 31 (6): 988 - 994.
- Suh, K.I., Lee, J.W. and Choi, W.Y. (1997).** A systematic study of the Mesochorinae (Hymenoptera: Ichneumonidae) from the eastern Palearctic region I. - a review of the genus *Mesochorus*. *Entomological Research Bulletin* 23: 1 - 28.
- Sulanc, M., Emre, I. and Yazgan, S. (1992).** Qualitative and quantitative effects of inorganic salts on development and synthesized protein amount of the male larvae of *Pimpla turionellae* L. (Hymenoptera: Ichneumonidae). *Doga, Turk Zooloji Dergisi* 16 (1): 92 - 100.
- Sulanc, M. and Emre, I. (2000).** Effects of B group vitamins and choline chloride on the development and protein synthesis in the male larvae of *Pimpla turionellae* L. (Hym.: Ichneumonidae). *Journal of Applied Entomology* 124 (3-4): 151 - 153.
- Sulistyowati, E. and Junianto, Y.D. (1995).** Inventory of natural enemies of cocoa pod borer (CPB), *Conopomorpha cramerella* (Snell.) in Maluku. *Pelita Perkebunan* 11 (2): 76 - 89.
- Sun, C.N. (1992).** Detoxifying enzymes of *Apanteles plutellae* and *Diadegma semiclausum*, parasitoids of diamondback moth larvae. *Resistant Pest Management* 4 (1): 21 - 22.
- Supatashvili, A.Sh. (1990).** Biological agents controlling *Diprion pini*. *Zashchita Rastenii Moskva* 6: 23 - 24.
- Surekha, K. (1995).** *Goryphus gibbosus* (Hymenoptera: Ichneumonidae) a new pupal parasite of *Opisina arenosella* (Xyloryctidae: Lepidoptera). *Journal of Ecobiology* 7 (3): 235 - 236.
- Sushil, S.N., Mishra, Y.D., Bhattacharya, A., Jaiswal, A.K. and Sharma, K.K. (1997).** Safety of endosulfan and dichlorvos to four parasitoids of lac insect predators. *Pest Management in Horticultural Ecosystems* 3 (1): 39 - 41.
- Szatmari, S., Balazs, K., Meszaros, Z., Jenser, G. and Polesny, F. (1998).** Lepidoptera living on raspberry in North-Hungary. Integrated plant protection in orchards, soft fruits. Workshop held in Vienna, Austria, 7-10 October, 1997. *Bulletin OILB SROP* 21 (10): 35 - 38.

## T

- Takahashi, H., Nishio, E. and Hayashi, H. (1993).** Pollination biology of the saprophytic species *Petrosavia sakurarii* (Makino) van Steenis in central Japan. *Journal of Plant Research* 106: 213 - 217.
- Talekar, N.S. (1988).** Biological control of diamondback moth in farmers' fields. AVRDC Publication, Asian Vegetable Research and Development Center 87 - 273: 50 - 54.
- Talekar, N.S. (1996).** Biological control of diamondback moth in Taiwan - a review. *Plant Protection Bulletin Taipei* 38 (3): 167 - 189.

- Talekar, N.S., Ooi, P.A.C., Lim, G.S., Ho, T.H., Manalo, P.L. and Waage, J.K. (1992).** Integrated management of diamondback moth: a collaborative approach in Southeast Asia. Integrated pest management in the Asia Pacific region 37 - 49.
- Talekar, N.S. and Yang, J.C. (1991).** Characteristic of parasitism of diamondback moth by two larval parasites. Entomophaga 36 (1): 95 - 104.
- Talekar, N.S. and Yang, J.C. (1993).** Influence of crucifer cropping system on the parasitism of *Plutella xylostella* (Lep.: Yponomeutidae) by *Cotesia plutellae* (Hym.: Braconidae) and *Diadegma semiclausum* (Hym.: Ichneumonidae). Entomophaga 38 (4): 541 - 550.
- Tanaka, K. (1992).** Size-dependent survivorship in the web-building spider *Agelena limbata*. Oecologia 90 (4): 597 - 602.
- Tanaka, T. and Vinson, S.B. (1991).** Interaction of venoms with the calyx fluids of three parasitoids, *Cardiochiles nigriceps*, *Microplitis croceipes* (Hymenoptera: Braconidae), and *Campoletis sonorensis* (Hymenoptera: Ichneumonidae) in effecting a delay in the pupation of *Heliothis virescens* (Lepidoptera: Noctuidae). Annals of the Entomological Society of America 84 (1): 87 - 92.
- Tarasco, E. (1995).** Larval morphology and biology of *Coelichneumon rudis* (Boyer de Fonscolombe) (Hymenoptera: Ichneumonidae), endoparasitoid of *Thaumetopoea pityocampa* (Denis et Schiffermuller) pupae. Entomologica 29: 5 - 51.
- Tarasco, E. (1996).** A gynandromorphic specimen of *Coelichneumon rudis* (Boyer de Fonscolombe, 1847) (Hymenoptera: Ichneumonidae). Entomologica 30: 101 - 104.
- Tarwacki, G. (1998).** Pine tortricids (Tortricidae) and their parasitoids in the Forest Experimental Department (LZD) at Rogow. Sylwan 142 (8): 89 - 95.
- Tarwacki, G. (1999).** Importance of the complex-centre method of forest protection in increasing environment resistance. Sylwan 143 (12): 73 - 79.
- Teder, T. (1998).** Limited variability of genitalia in the genus *Pimpla* (Hymenoptera: Ichneumonidae): inter- or intraspecific causes? Netherlands Journal of Zoology 48 (4): 335 - 347.
- Teder, T., Tammaru, T. and Pedmanson, R. (1999).** Patterns of host use in solitary parasitoids (Hymenoptera: Ichneumonidae): field evidence from a homogeneous habitat. Ecography 22 (1): 79 - 86.
- Teder, T., Tanhuanpaa, M., Ruohomaki, K., Kaitaniemi, P. and Henriksson, J. (2000).** Temporal and spatial variation of larval parasitism in non-outbreaking populations of a folivorous moth. Oecologia 123 (4): 516 - 524.
- Teggelli, G.R., Biradar, A.P., Vastrad, S.M. and Rajashaker, D.W. (1998).** Impact of pesticides on an endoparasitoid, *Campoletis chlorideae* Uchida. Advances in Agricultural Research in India, publ. 2000, 10: 25 - 29.
- Teggelli, R.G., Biradar, A.P., Balikai, R.A. and Rao, K.J. (1997).** Host specificity and host age on the development of *Campoletis chlorideae* Uchida. Advances in Agricultural Research in India 8: 93 - 97.

- Tejada, L.O. and Cansino, E.R. (1986).** Genera of Ichneumonidae (Hymenoptera) of the northeast of Mexico. Informe de Investigacion, Division de Ciencias Agropecuarias y Maritimas, Instituto Tecnologico y de Estudios Superiores de Monterrey, Mexico 19: 82.
- Telekar, N.S. and Yang, J.C. (1990).** Use of parasitoids for controlling the diamondback moth, *Plutella xylostella* (Lep.: Yponomeutidae).FFTC-NARC International Seminar on 'The use of parasitoids and predators to control agricultural pests', Tukuba Science City, Ibaraki ken 305, Japan, 2-7 October 1989. 21 pp.
- Tempest, A.M., Bergmann, E.C., De Faria, A.M., Penteado-Dias, A.M., Imenes, S.D.L. and Guerra, T.M. (1998).** About the Ichneumonidae in a rubber-tree crop (*Hevea brasiliensis*). Arquivos do Instituto Biologico Sao Paulo 65 (2): 63 - 68.
- Tenhumberg, B. and Poehling, H.M. (1989).** Population dynamics of syrphids (Diptera: Syrphidae) in winter wheat. Mitteilngen der Deutschen Gesellschaft fur Allgemeine und Angewandte Entomologie 7 (1-3): 318 - 324.
- Tereshkin, A. (1992).** A new tribe, a new genus and a new species of the Ichneumoninae Stenopneusticae from Europe and Siberia (Hymenoptera, Ichneumonidae). Entomofauna 13: 193-201.
- Tereshkin, A. (1994).** A new genus and species of the Ichneumoninae Stenopneusticae from the Far East (Hymenoptera, Ichneumonidae). Entomofauna 15: 117-123.
- Thakur, J.N., Singh, J.P., Verma, O.P. and Diwakar, M.C. (1995).** Bioecological studies on gram pod borers *Heliothis* species under Jammu conditions. Journal of Advanced Zoology 16 (2): 118 - 122.
- Thakur, N.S.A. and Deka, T.C. (1994).** *Hyposter ebeninus* Gravenhorst (Hymenoptera: Ichneumonidae): a larval endoparasite of cabbage butterfly, *Pieris brassicae* L. from Meghalaya, India. Indian Journal of Hill Farming 7 (1): 81 - 85.
- Thaler, J.S. (1999).** Jasmonate-inducible plant defences cause increased parasitism of herbivores. Nature London 399: 6737, 686 - 688.
- Thayer, M.K., Newton, A.F. Jr. and Pape, T. (1992).** Case 793. Metopiinae Foester, 1868 (Insecta, Hymenoptera), Metopiini Raffray, 1904 (Insecta: Coleoptera), and Metopiini Townsend, 1908 (Insecta: Diptera): proposed removal of homonymy. Bulletin of Zoological Nomenclature 49 (3): 200 - 204.
- Theilmann, D.A. and Summers, M.D. (1988).** Identification and comparison of *Campoletis sonorensis* virus transcripts expressed from four genomic segments in the insect hosts *Campoletis sonorensis* and *Heliothis virescens*. Virology New York 167 (2): 329 - 341.
- Theopold, U., Kinuthia, W., Schmidt, O., Li, D.M., Austin, A.D. and Downton, M. (2000).** Protection by immune disguise: a new lesson from a parasitoid wasp. Fourth International Hymenoptera Conference, held in Canberra, Australia, in January 1999. Hymenoptera: evolution, biodiversity and biological control 23 - 28.
- Theopold, U., Krause, E. and Schmidt, O. (1994).** Cloning of a VLP-protein coding gene from a parasitoid wasp *Venturia canescens*. Archives of Insect Biochemistry and Physiology 26 (2-3): 137 - 145.

- Thibout, E. (1988).** The specificity of *Diadromus pulchellus* (Hymenoptera: Ichneumonidae) towards its host *Acrolepiopsis assectella*, the leek moth. *Entomophaga* 33 (4): 439 - 452.
- Thibout, E., Guillot, J.F., Ferary, S., Limouzin, P. and Auger, J. (1995).** Origin and identification of bacteria which produce kairomones in the frass of *Acrolepiopsis assectella* (Lep.: Hyponomeutoidea). *Experientia* 51 (11): 1073 - 1075.
- Thibout, E., Guillot, J.F. and Auger, J. (1993).** Microorganisms are involved in the production of volatile kairomones affecting the host seeking behaviour of *Diadromus pulchellus*, a parasitoid of *Acrolepiopsis assectella*. *Physiological Entomology* 18 (2): 176 - 182.
- Thibout, E., Lecomte, C. and Auger, J. (1988).** *Diadromus pulchellus*: search for a host and specificity. *Colloques de l'INRA* 48: 7 - 14.
- Thiery, D., Xuereb, A., Villemant, C., Sentenac, G., Delbac, L., Kuntzman, P. and Lozzia, C. (2001).** The parasites of grape tortricids: noticed on several species present in 3 French vine regions. Proceedings of the IOBC-WPRS Working Group "Integrated Control in Viticulture" at Ponte de Lima, Portugal, 2-7 March, 2001. *Bulletin OILB SROP* 24 (7): 135 - 141.
- Thies, C. and Tschardt, T. (1999).** Landscape structure and biological control in agroecosystems. *Science Washington* 285: 893 - 895.
- Thoeny, W.T. and Nordin, G.L. (1991).** Parasitoids reared from the locust twig borer, *Ecdytolopha insiticiana* Zell. (Lepidoptera: Tortricidae) in southeastern Kentucky. *Journal of the Kansas Entomological Society* 64 (1): 77 - 79.
- Thomas, J.A., Knapp, J.J., Akino, T., Gerty, S., Wakamura, S., Simcox, D.J., Wardlaw, J.C. and Elmes, G.W. (2002).** Parasitoid secretions provoke ant warfare: subterfuge used by a rare wasp may be the key to an alternative type of pest control. *Nature London* 417: 688, 505 - 506.
- Thomas, J.A. and Elmes, G.W. (1993).** Specialized searching and the hostile use of allomones by a parasitoid whose host, the butterfly *Maculinea rebeli*, inhabits ant nests. *Animal Behaviour* 45 (3): 593 - 602.
- Tilakaratna, D. (1991).** Parasites of the teak defoliator, *Hyblaea puera*. Sri Lanka Forester, publ. 1995? 20 (1-2): 23 - 25.
- Tillman, P.G. and Powell, J.E. (1989).** Comparison of acceptance of larval instars of the tobacco budworm (Lepidoptera: Noctuidae) by *Microplitis croceipes*, *Microplitis demolitor*, *Cotesia kazak* (Hymenoptera: Braconidae), and *Hyposoter didymator* (Hymenoptera: Ichneumonidae). *Journal of Agricultural Entomology* 6 (4): 201 - 209.
- Tillman, P.G. and Powell, J.E. (1991).** Developmental time in relation to temperature for *Microplitis croceipes*, *M. demolitor*, *Cotesia kazak* (Hymenoptera: Braconidae), and *Hyposoter didymator* (Hymenoptera: Ichneumonidae), endoparasitoids of the tobacco budworm (Lepidoptera: Noctuidae). *Environmental Entomology* 20 (1): 61 - 64.
- Tillman, P.G. and Powell, J.E. (1992).** Intraspecific host discrimination and larval competition in *Microplitis croceipes*, *Microplitis demolitor*, *Cotesia kazak* (Hym.:)



- Braconidae) and *Hyposoter didymator* (Hym.: Ichneumonidae), parasitoids of *Heliothis virescens* (Lep.: Noctuidae). *Entomophaga* 37 (2): 229 - 237.
- Tingle, F.C., Mitchell, ER. and McLaughlin, JR. (1994).** Lepidopterous pests of cotton and their parasitoids in a double-cropping environment. *Florida Entomologist* 77 (3): 334 - 341.
- Tipping, P.W. (1993).** Field studies with *Cassida rubiginosa* (Coleoptera: Chrysomelidae) in Canada thistle. *Environmental Entomology* 22 (6): 1402 - 1407.
- Tkaczuk, C. (1996).** *Pachynematus pumilio*, a little known pest of black currants. *Ochrona Roslin* 40 (7): 7 - 8.
- Tkaczuk, C., Mietkiewski, R. and Filipowicz, A. (1995).** Natural enemies of the blackcurrant sawfly (*Pachynematus pumilio* Knw.). *Materialy ogólnopolskiej konferencji naukowej Nauka Praktyce Ogrodniczej z okazji XXV-lecia Wydziału Ogrodniczego Akademii Rolniczej w Lublinie* 395 - 398.
- Tobias, V.I. (1988).** Family Paxylomatidae (Hymenoptera) of the fauna of the USSR. (in Russian) *Trudy Vsesoyuznogo éntomologicheskogo obshchestva* 70: 131-143.
- Tobias, V.I. (1993).** Dependence of wing venation in the Hymenoptera on the ecological environment of their habitats. *Entomologicheskoe Obozrenie* 72 (3): 497 - 506.
- Toft, R.J., Malham, J.P. and Beggs, J.R. (1999).** Mortality and emergence pattern of overwintering cocoons of the wasp parasitoid *Sphecophaga vesparum vesparum* (Hymenoptera: Ichneumonidae) in New Zealand. *Environmental Entomology* 28 (1): 9 - 13.
- Togashi, I. and Ishikawa, T. (1994).** Parasites reared from cocoons of *Monema flavescens* Walker and *Latoia sinica* (Moore) (Lepidoptera: Limacodidae) in Ishikawa prefecture. *Transactions of the Shikoku Entomological Society* 20 (3-4): 321 - 325.
- Tolkanitz, V.I. (1974).** Revision of the *Netelia* (Hymenoptera, Ichneumonidae) and a review of the species of the fauna of the USSR. (in Russian) *Zoologicheskii Zhurnal* 53: 376-393.
- Tolkanits, V.I. (1999).** New and little-known Palearctic species of the genus *Exochus* (Hymenoptera: Ichneumonidae: Metopiinae). *Zoologicheskii Zhurnal* 78 (2): 191 - 201.
- Tolkanits, V.I. (2001).** New species of the genus *Exochus* (Hymenoptera: Ichneumonidae, Metopiinae) from Russia and Mongolia. *Zoologicheskii Zhurnal* 80 (11): 1404 - 1408.
- Tolkanits, V.I. and Seregina, L.Ya. (1988).** Ichneumon flies (Hymenoptera: Ichneumonidae) - natural enemies of the oak leaf roller in the south of the Ukraine. *Vestnik Zoologii* 1: 22 - 29.
- Tong, X.W. (1987).** New host records of Chalcidoidea. *Natural Enemies of Insects* 9 (1): 54.
- Tooker, J.F. and Hanks, L.M. (2000).** Flowering plant hosts of adult hymenopteran parasitoids of central Illinois. *Annals of the Entomological Society of America* 93 (3): 580 - 588.
- Topping, C.J., Sunderland, K.D. and Van Baarlen, P. (1996).** Host location by *Gelis festinans*, an eggsac parasitoid of the linyphiid spider *Erigone atra*. *Entomologia Experimentalis et Applicata* 81 (2): 155 - 163.

- Torres, J.A. (1992).** Lepidoptera outbreaks in response to successional changes after the passage of Hurricane Hugo in Puerto Rico. *Journal of Tropical Ecology* 8 (3): 285 - 298.
- Torres-Vila, L.M., Rodriguez-Molina, M.C., Palo, E., Del Estal, P. and Lacasa, A. (2000).** The larval parasitoid complex of *Helicoverpa armigera* on tomato in the Vegas del Guadiana (Extremadura). *Boletín de Sanidad Vegetal, Plagas* 26 (3): 323 - 333.
- Townes, H. (1961).** A key to the genera of Ichneumonidae recorded from the Indo-Australian area. *In: Townes, H., M. Townes, H. and Gupta, V.K. (1961).* A catalogue and reclassification of the Indo-Australian Ichneumonidae. *Memoirs of the American Entomological Institute* 1: 1-522.
- Townes, H. (1961).** Generic descriptions. *In: Townes, H., Townes, M. and Gupta, V.K. 1961.* A catalogue and reclassification of the Indo-Australian Ichneumonidae. *Memoirs of the American Entomological Institute* 1: 1-522.
- Townes, H. (1965).** Descriptions of new genera. *In: Townes, H., S. Momoi, & M. Townes. 1965.* A catalogue and reclassification of the Eastern Palearctic Ichneumonidae. *Memoirs of the American Entomological Institute* 5: 1-661.
- Townes, H. (1966).** Descriptions of new genera. *In: Townes, H. & M. Townes. 1966.* A catalogue and reclassification of the Neotropic Ichneumonidae. *Memoirs of the American Entomological Institute* 8: 1-367.
- Townes, H. (1969).** The genera of Ichneumonidae, part 1. *Memoirs of the American Entomological Institute* 11: 1-300.
- Townes, H. (1970).** The genera of Ichneumonidae, part 2. *Memoirs of the American Entomological Institute* 12: 1-537.
- Townes, H. (1970).** The genera of Ichneumonidae, part 3. *Memoirs of the American Entomological Institute* 13: 1-307.
- Townes, H. (1971).** The genera of Ichneumonidae, part 4. *Memoirs of the American Entomological Institute* 17: 1-372.
- Townes, H. (1972).** Ichneumonidae as biological control agents. *Proceedings Tall Timbers conference on ecological animal control by habitat management* 3:235-248.
- Townes, H. (1973).** Descriptions of new genera. *In: Townes, H. & M. Townes. 1973.* A catalogue and reclassification of the Ethiopian Ichneumonidae. *Memoirs of the American Entomological Institute* 19: 1-416.
- Townes, H. (1973).** *Gessia*, new name. *In: Townes, H. & M. Townes. 1973.* A catalogue and reclassification of the Ethiopian Ichneumonidae. *Memoirs of the American Entomological Institute* 19: 1-416.
- Townes, H. (1973).** Two ichneumonids (Hymenoptera) from the early Cretaceous. *Proceedings of the Entomological Society of Washington* 75: 216-219.
- Townes, H. (1978).** *Brachychroa*, new genus. *In: Townes, H. & M. Townes. 1978.* Ichneumon-flies of America North of Mexico: 7. Subfamily Banchinae, tribes Lissonotini and Banchini. *Memoirs of the American Entomological Institute* 26: 1-614.

- Townes, H. (1983).** Revisions of twenty genera of Gelini (Hymenoptera). *Memoirs of the American Entomological Institute* 35: 1-281.
- Townes, H., Gupta, V. and Townes, M. (1992).** The ichneumon-flies of America North of Mexico: Part 11. Subfamily Tryphoninae, Tribes Oedemopsini, Tryphonini, and Idiogrammatini. *Memoirs of the American Entomological Institute* 51: 1-292.
- Townes, H., Momoi, S. and Townes, M. (1965).** A catalogue and reclassification of the Eastern Palearctic Ichneumonidae. *Memoirs of the American Entomological Institute* 5: 1-661.
- Townes, H. and Townes, M. (1966).** A catalogue and reclassification of the Neotropic Ichneumonidae. *Memoirs of the American Entomological Institute* 8: 1-367.
- Townes, H. and Townes, M. (1973).** A catalogue and reclassification of the Ethiopian Ichneumonidae. *Memoirs of the American Entomological Institute* 19: 1-416.
- Townes, H. and Townes, M. (1978).** Ichneumon-flies of America North of Mexico: 7. Subfamily Banchinae, tribes Lissonotini and Banchini. *Memoirs of the American Entomological Institute* 26: 1-614.
- Townes, H., Townes, M. and Gupta, V.K. (1961).** A catalogue and reclassification of the Indo-Australian Ichneumonidae. *Memoirs of the American Entomological Institute* 1: 1-522.
- Tran, T.V. and Takasu, K. (2000).** Host age selection by the host-feeding pupal parasitoid *Diadromus subtilicornis* (Gravenhorst) (Hymenoptera: Ichneumonidae). *Applied Entomology and Zoology* 35 (4): 549 - 556.
- Tran, T.V. and Takasu, K. (2000).** Life history of the pupal parasitoid *Diadromus subtilicornis* (Gravenhorst) (Hymenoptera: Ichneumonidae) as influenced by temperature, photoperiod, and availability of food and hosts. *Entomological Science* 3 (2): 255 - 264.
- Triggiani, O. (1986).** Observations on the biology and behaviour of *Dioryctria pineae* (Stgr.) (Lepidoptera: Phycitidae) and on its parasitoid *Elachertus geniculatus* (Ratz.) (Hymenoptera: Chalcidoidea). *Entomologica* 21: 141 - 153.
- Triggiani, O. and Tarasco, E. (1998).** Biological and morphological aspects of *Curculio (Balaninus) propinquus* (Desbrochers) (Coleoptera: Curculionidae). *Entomologica* 32: 179 - 193.
- Tripathi, C.P.M., Kumar, N. and Kumar, A. (2000).** Sex ratio of *Campoletis chloridae* Uchida in response to *Helicoverpa armigera* (Hubner) density. *Insect Science and its Application* 20 (1): 73 - 76.
- Trudeau, D. and Gordon, D.M. (1989).** Factors determining the functional response of the parasitoid *Venturia canescens*. *Entomologia Experimentalis et Applicata* 50 (1): 3 - 6.
- Tsankov, G. and Georgiev, G. (1991).** Records on parasitoids of smaller poplar borer, *Saperda populnea* (Coleoptera: Cerambycidae) along the Danube in Bulgaria. *Entomophaga* 36 (4): 493 - 498.
- Tsankov, G. and Georgiev, G. (1992).** New species parasitizing the willow moth *Hyponomeuta rorellus* in Bulgaria. *Nauka za Gorata* 29 (4): 68 - 73.

- Tsankov, G. and Gupta, V.K. (1988).** The ichneumonoid parasites of the European pine shoot moth caterpillars in Bulgaria (Hymenoptera: Ichneumonoidea). *Advances in parasitic Hymenoptera research: Proceedings of the II Conference on the Taxonomy and Biology of Parasitic Hymenoptera.* recd. 1993, 391 - 394.
- Tsankov, G. and Stalev, Z. (1993).** Parasitoids of larvae and pupae of *Rhyacionia buoliana* in the Strandzha region: species composition, distribution and effect. *Nauka za Gorata* 30 (1): 57 - 63.
- Tseng, H.K. (1988).** Utilization of portable bottle for the rearing methods of enemy of tea pest-parasitoids. *Taiwan Tea Research Bulletin* 7: 27 - 30.
- Tudor, I. and Marcu, O. (1997).** Research into ichneumonid parasites in pupae of *Tortrix viridana*. *Buletinul Universitatii din Brasov. Seria B, Economie Forestiera* 19: 55 - 59.
- Turkar, K.S., Gupta, R., Banerjee, S.K. and Wanjari, R.R. (2000).** Influence of intercropping chickpea with coriander on parasitisation of *Heliothis armigera* (Hubner) by *Campoletis chlorideae* Uchida. *Journal of Entomological Research* 24 (3): 279 - 281.
- Turkmen, S. (1987).** A new pest of legumes (*Amicta oberthuri* Hey.) (Lepidoptera: Psychidae) in southeast Anatolia. *Turkiye I. Entomoloji Kongresi Bildirileri*, 13-16 Ekim, Ege Universitesi, Bornovo, Izmir 239 - 248.
- Turnock, W.J. (1988).** Density, parasitism, and disease incidence of larvae of the bertha armyworm, *Mamestra configurata* Walker (Lepidoptera: Noctuidae), in Manitoba, 1973-1986. *Canadian Entomologist* 120 (5): 401 - 413.
- Tybirk, K. (1989).** Flowering, pollination and seed production of *Acacia nilotica*. *Nordic Journal of Botany* 9 (4): 375 - 381.

## U

- Uchoa-Fernandes, M.A., De Campos, W.G. and De Campos, W.G. (1993).** Parasitoids of larvae and pupae of the tomato worm *Scrobipalpus absoluta* Meyrick, 1917 (Lepidoptera: Gelechiidae). *Revista Brasileira de Entomologia* 37 (3): 399 - 402.
- Uckan, F. (1999).** The morphology of the venom apparatus and histology of venom gland of *Pimpla turionellae* (L.) (Hym.: Ichneumonidae) females. *Turkish Journal of Zoology* 23 (4): 461 - 466.
- Udayagiri, S. (1988).** Life history and new records of natural enemies of *Trilochoa varians* (Walker) (Lepidoptera: Bombycidae). *Annals of Entomology* 6 (1): 1 - 6.
- Uematsu, H. (1988).** Reproductive capacity of *Diadromus subtilicornis* (Hymenoptera: Ichneumonidae), a pupal parasitoid of the diamondback moth, *Plutella xylostella* (Lepidoptera: Plutellidae). *Proceedings of the Association for Plant Protection of Kyushu* 34: 144 - 146.
- Uematsu, H., Yamashita, T. and Sakanoshita, A. (1987).** Parasitoids of the diamondback moth, *Plutella xylostella* L. and their seasonal incidence in Miyazaki district, in the southern part of Japan. *Proceedings of the Association for Plant Protection of Kyushu* 33: 136 - 138.

- Uematsu, H. and Yamashita, T. (1999).** Seasonal occurrence of diamondback moth, *Plutella xylostella* (L.) and parasitoids in some commercial cruciferous fields in Miyazaki, Southern Japan. *Japanese Journal of Applied Entomology and Zoology* 43 (3): 113 - 121.
- Ueno, T. (1995).** Abdominal tip movements during oviposition by two parasitoids (Hymenoptera: Ichneumonidae) as an index of predicting the sex of depositing eggs. *Applied Entomology and Zoology* 30 (4): 588 - 590.
- Ueno, T. (1996).** Effect of superparasitism on offspring survival of sex ratio in the parasitoid wasp, *Itopectis naranyae* (Hymenoptera: Ichneumonidae). *Proceedings of the Kanto TosanPlant Protection Society* 43: 175 - 178.
- Ueno, T. (1996).** Estimation of host-size dependent sex ratio in the polyphagous parasitoid *Itopectis naranyae* Ashmead (Hymenoptera: Ichneumonidae) in the field. *Applied Entomology and Zoology* 31 (4): 613 - 615.
- Ueno, T. (1996).** Physical elimination of competitive eggs and larvae in hosts by early instar larvae of *Itopectis naranyae* (Hymenoptera: Ichneumonidae). *Proceedings of the Kanto Tosan Plant Protection Society* 43: 179 - 181.
- Ueno, T. (1998).** Adaptiveness of sex ratio control by the pupal parasitoid *Itopectis naranyae* (Hymenoptera: Ichneumonidae) in response to host size. *Evolutionary Ecology* 12 (6): 643 - 654.
- Ueno, T. (1998).** Selective host-feeding on parasitized hosts by the parasitoid *Itopectis naranyae* (Hymenoptera: Ichneumonidae) and its implication for biological control. *Bulletin of Entomological Research* 88 (4): 461 - 466.
- Ueno, T. (1998).** Sex allocation by a parasitoid wasp (Hymenoptera: Ichneumonidae) to different host species: a question for the mechanism of host size estimation. *Journal of Insect Behavior* 11 (6): 811 - 821.
- Ueno, T. (1999).** Adult size and reproduction in the ectoparasitoid *Agrothereutes lanceolatus* Walker (Hym.: Ichneumonidae). *Journal of Applied Entomology* 123 (6): 357 - 361.
- Ueno, T. (1999).** Behavioural evidence for recognition of parasitized hosts by superparasitizing *Pimpla nipponica* (Hymenoptera: Ichneumonidae). *Canadian Journal of Zoology* 77 (3): 413 - 417.
- Ueno, T. (1999).** Host suitability and sex ratio in two strains of the parasitoid *Itopectis naranyae* (Hymenoptera: Ichneumonidae). *Annual Report of the Kanto Tosan Plant Protection Society* 46: 81 - 83.
- Ueno, T. (1999).** Host suitability and sex-ratio differences in wild-caught and laboratory-reared parasitoid *Pimpla parnarae* (Hymenoptera: Ichneumonidae). *Annals of the Entomological Society of America* 92 (4): 609 - 614.
- Ueno, T. (1999).** Host-feeding and acceptance by a parasitic wasp (Hymenoptera: Ichneumonidae) as influenced by egg load and experience in a patch. *Evolutionary Ecology* 13 (1): 33 - 44.

- Ueno, T. (1999).** Multiparasitism and host feeding by solitary parasitoid wasps (Hymenoptera: Ichneumonidae) based on the pay-off from parasitized hosts. *Annals of the Entomological Society of America* 92 (4): 601 - 608.
- Ueno, T. (1999).** Reproduction and host-feeding in the solitary parasitoid wasp *Pimpla nipponica* (Hymenoptera: Ichneumonidae). *Invertebrate Reproduction and Development* 35 (3): 231 - 237.
- Ueno, T. (2002).** Biology of the ectoparasitoid wasp *Agrothereutes lanceolatus* (Hymenoptera: Ichneumonidae): host acceptance and larval development on a laboratory host. *Journal of the Faculty of Agriculture, Kyushu University* 47 (1): 37 - 43.
- Ueno, T., Godfray, H.C.J. and Shimada, M. (1999).** Host-size-dependent sex ratio in a parasitoid wasp. Special issue. Parasitoids: a model system to answer questions in behavioral, evolutionary and population ecology. *Researches on Population Ecology* 41 (1): 47 - 57.
- Ueno, T. and Hirai, K. (1999).** Larval and pupal parasitoids of oriental corn borer *Ostrinia furnacalis* (Lepidoptera: Pyralidae). *Japanese Journal of Applied Entomology and Zoology* 43 (3): 135 - 137.
- Ueno, T. and Tanaka, T. (1994).** Can a female parasitoid recognize a previously rejected host? *Animal Behaviour* 47 (4): 988 - 990.
- Ueno, T. and Tanaka, T. (1994).** Comparative biology of six polyphagous solitary pupal endoparasitoids (Hymenoptera: Ichneumonidae): differential host suitability and sex allocation. *Annals of the Entomological Society of America* 87 (5): 592 - 598.
- Ueno, T. and Tanaka, T. (1996).** Self-host discrimination by a parasitic wasp [*Pimpla nipponica*]: the role of short-term memory. *Animal Behaviour* 52 (5): 875 - 883.
- Ueno, T. and Tanaka, T. (1997).** Comparison between primary and secondary sex ratios in parasitoid wasps using a method for observing chromosomes. *Entomologia Experimentalis et Applicata* 82 (1): 105 - 108.
- Ugur, A. (1996).** The effect of superparasitism on development of *Pimpla turionellae* (L.) (Hymenoptera: Ichneumonidae). *Turkiye Entomoloji Dergisi* 20 (1): 51 - 58.
- Ugur, A., Kansu, I.A. and Kedici, R. (1987).** Investigations of the development of *Pimpla turionellae* (L.) (Hymenoptera: Ichneumonidae) in the pupae of *Agrotis segetum* (Denis and Schiff.) (Lepidoptera: Noctuidae) reared on various food plants. *Turkiye I. Entomoloji Kongresi Bildirileri*, 13-16 Ekim 1987, Ege Universitesi, Bornova, Izmir 481 - 490.
- Ugur, A. and Kansu, I.A. (1990).** The determination of an optimal parasite:host ratio and duration of parasite contact with the host in the rearing of *Pimpla turionellae* (L.) (Hymenoptera: Ichneumonidae). *Proceedings of the Second Turkish National Congress of Biological Control* 67 - 72.
- Ugur, A. and Kansu, I.A. (1994).** Some effects of hosts on rearing *Pimpla turionellae* (L.) (Hymenoptera: Ichneumonidae). *Turkiye III. Biyolojik Mucadele Kongresi Bildirileeri*, 25-28 Ocak 1994, Ege Universitesi Ziraat Fakultesi, Bitki Koruma Bolumu, Izmir 215 - 220.

- Ulber, B., Paul, V.H., Foller, I., Evans, N. and Williams, I. (2000).** Bibliography of parasitoid species and levels of parasitism of rape stem weevil *Ceutorhynchus napi* Gyll. and cabbage stem weevil *Ceutorhynchus pallidactylus* (Mrsh.). Working group "Integrated control in oilseed crops". Proceedings of the meeting at Prague (Czech Republic) May 31 June 2, 1999. Bulletin OILB SROP 23 (6): 131 - 134.
- Ulber, B. and Von Klingenberg, A. (1994).** Investigations on the occurrence of Tersilochinae (Hym.: Ichneumonidae) as parasitoids of oilseed rape pests in the Gottingen region in 1990 and 1991, and on their emergence following various tillage techniques. Untersuchungen zum Auftreten der Tersilochinae (Hym.: Ichneumonidae) als Larvalparasitoide einiger Rapsschadlinge im Raum Gottingen 1990 und 1991 und zu deren Schlupfabundanz nach unterschiedlicher Bodenbearbeitung. Journal of Applied Entomology 117 (3): 287 - 299.
- Ulpah, S. and Kok, L.T. (1996).** Interrelationship of *Bacillus thuringiensis* Berliner to the diamondback moth (Lepidoptera: Noctuidae) and its primary parasitoid, *Diadegma insulare* (Hymenoptera: Ichneumonidae). Journal of Entomological Science 31 (4): 371 - 377.
- Ulpah, S. and Kok, L.T. (1996).** Interrelationship of *Bacillus thuringiensis* Berliner to the diamondback moth (Lepidoptera: Noctuidae) and its primary parasitoid, *Diadegma insulare* (Hymenoptera: Ichneumonidae). Journal of Entomological Science 31 (4): 371 - 374.
- Ulrich, W. (1999).** The number of species of Hymenoptera in Europe and assessment of the total number of Hymenoptera in the world. Polskie Pismo Entomologiczne 68 (2): 151 - 164.
- Ulrichs, C., Mewis, I., Schnitzler, W.H. and Burleigh, J.R. (2001).** Effectivity of synthetic insecticides against *Maruca vitrata* F. and the parasitoid *Bassus asper* CHOU & SHARKEY in the Philippines. Vortrage der Entomologentagung in Dusseldorf vom 26. bis. 31. Marz 2001. Mitteilungen der Deutschen Gesellschaft fur allgemeine und angewandte Entomologie 13 (1-6): 279 - 282.
- Ulrichs, C., Mewis, I., Schnitzler, W.H. and Burleigh, J.R. (2001).** Parasitoids of the bean podborer, *Maruca vitrata* F. (Lepidoptera: Pyraustinae), a pest of *Vigna sesquipedalis* in the Philippine lowlands. Vortrage der Entomologentagung in Dusseldorf vom 26. bis. 31. Marz 2001. Mitteilungen der Deutschen Gesellschaft fur allgemeine und angewandte Entomologie 13 (1-6): 283 - 288.
- Upadhyay, R.N., Dubey, O.P. and Vaishampayan, S.M. (1999).** Studies on the common predatory and parasitic species of insects collected on light trap. JNKVV Research Journal, publ. 2000. 33 (1-2): 50 - 57.
- Urano, T. and Ito, K. (1993).** Life histories of the parasitoid wasps and their percent parasitism on inoculated larvae of the cryptomeria bark borer, *Semanotus japonicus* Lacordairei (Coleoptera: Cerambycidae). Journal of the Japanese Forestry Society 75 (5): 409 - 415.

- Urban, J. (1997).** Causes and progression of a pale tussock moth outbreak in the White Carpathian mountains and the Hostynska highlands. *Zpravy Lesnickeho Vyzkumu* 42 (4): 11 - 15.
- Urban, J. (2000).** Linden and its important leaf-eating pest. *Zpravy Lesnickeho Vyzkumu* 45 (2): 13 - 19.
- Urban, J. (2001).** Contribution to the knowledge of the development and harmfulness of *Caliroa annulipes* Klug (Hymenoptera: Tenthredinidae). *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis* 49 (3): 7 - 27.
- Urban, J. and Sedivy, J. (1997).** Factors regulating the mass outbreak of the pale tussock moth (*Calliteara pudibunda* L.) (Lepidoptera, Lymantriidae). *Lesnictvi Forestry* 43 (2): 67 - 78.
- Usha-Chauhan., Bhalla, O.P. and Sharma, K.C. (1997).** Biology and seasonality of the diamondback moth, *Plutella xylostella* (L.) (Lepidoptera: Yponomeutidae) and its parasitoids on cabbage and cauliflower. *Pest Management in Horticultural Ecosystems* 3 (1): 7 - 12.
- Uzun, S. (1987).** Parasites of the cabbage butterfly (*Pieris brassicae* (L.)) (Lepidoptera: Pieridae) harmful on cabbage and cauliflower in the Izmir district. *Turkiye Entomoloji Dergisi* 11 (4): 237 - 245.

## V

- Valicente, F.H. (1989).** Survey of the natural enemies of *Spodoptera frugiperda* (J.E. Smith, 1797) (Lepidoptera: Noctuidae) in different regions of Minas Gerais. *Anais da Sociedade Entomologica do Brasil* 18 (1): 119 - 130.
- Valicente, F.H. and Barreto, M.R. (1999).** Survey of natural enemies of the fall armyworm, *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae), in the Cascavel region, PR, Brazil. *Anais da Sociedade Entomologica do Brasil* 28 (2): 333 - 337.
- Van Achterberg, C. (1990).** Illustrated key to the subfamilies of the Holarctic Braconidae (Hymenoptera: Ichneumonoidea). *Zoologische Mededelingen* 64 (1-6): 1 - 20.
- Van Achterberg, C. (1999).** The West Palearctic species of the subfamily Paxylommatinae (Hymenoptera: Ichneumonidae), with special reference to the genus *Hybrizon* Fallen. *Zoologische Mededelingen* 73: 11 - 26.
- Van Den Berg, H., Nyambo, B.T. and Waage, J.K. (1990).** Parasitism of *Helicoverpa armigera* (Lepidoptera: Noctuidae) in Tanzania: analysis of parasitoid-crop associations. *Environmental Entomology* 19 (4): 1141 - 1145.
- Van Nouhuys, S. and Tay, W.T. (2001).** Causes and consequences of small population size for a specialist parasitoid wasp. *Oecologia* 128 (1): 126 - 133.
- Van Rossem, G. (1981).** A revision of Western Palearctic oxytorine genera (Hymenoptera, Ichneumonidae). *Spixiana (Suppl.)* 4: 79-135.
- Van Rossem, G. (1982).** A revision of Western Palearctic oxytorine genera. Part II. Genus *Eusterinx* (Hymenoptera, Ichneumonidae). *Spixiana* 5: 149-170.



- Van Rossem, G. (1987).** A revision of Western Palearctic oxytorine genera. Part VI. (Hymenoptera, Ichneumonidae). Tijdschrift voor Entomologie 130: 49-108.
- Van Rossem, G. (1988).** A revision of Palearctic oxytorine genera. Part VII. (Hymenoptera, Ichneumonidae). Tijdschrift voor Entomologie 131: 103-112.
- Van Rossem, G. (1989).** The genus *Cryptus Fabricius*, 1804 of the Canary Islands (Hymenoptera, Ichneumonidae). Tijdschrift voor Entomologie 132 (2): 251 - 258.
- Van Rossem, G. (1990).** Key to the genera of the palearctic Oxytorinae, with the description of three new genera (Hymenoptera: Ichneumonidae). Zoologische Mededelingen 63: 309-323.
- Vargas-Camplis, J. (1995).** Seasonal fluctuation of larval parasites of bollworm complex before boll weevil eradication program in northern Tamaulipas Mexico. Proceedings Beltwide Cotton Conferences, San Antonio, TX, USA, January 4-7, 1995: Vol 2: 790 - 791.
- Vargas-Osuna, E., Munoz-Ledesma, J., Aldebis, H.K. and Santiago-Alvarez, C. (1994).** Pathogens and parasitoids for the control of *Thaumetopoea pityocampa* (D. y Schiff.) (Lep.: Notodontidae). Boletín de Sanidad Vegetal, Plagas 20 (2): 511 - 515.
- Varkonyi, G. (1998).** *Metopius harpyiae* (Hymenoptera: Ichneumonidae: Metopiinae) new to Finland. Entomologica Fennica 9 (1): 52.
- Varkonyi, G., Hanski, I., Rost, M. and Itamies, J. (2002).** Host-parasitoid dynamics in periodic boreal moths. Oikos 98 (3): 421 - 430.
- Vas, J., Marko, V., Abraham, L. and Meszaros, Z. (2001).** Study of Neuropteroidea (Raphidioptera: Neuroptera) communities by using Malaise traps in an untreated orchard and its environment. Acta Phytopathologica et Entomologica Hungarica 36 (1-2): 115 - 122.
- Velez-Angel, R. (1988).** Recognition, identification and biology of species of Gelechiidae (Lepidoptera) on solanaceous plants from the Antioquia department. I. *Scrobipalpula absoluta* (Meyrick). Revista Colombiana de Entomología 14 (2): 25 - 32.
- Venkatesan, T., Rao, N.S. and Joshi, S. (1999).** Influence of mating age, parental sex-ratio, and host age on the female progeny emergence of *Campoletis chlorideae* Uchida (Hymenoptera: Ichneumonidae). Shashpa 6 (2): 145 - 148.
- Venkatesan, T., Singh, S.P., Joshi, S. and Rao, N.S. (1995).** Breeding of *Campoletis chlorideae* Uchida (Hymenoptera : Ichneumonidae) on *Corcyra cephalonica* Stainton. Journal of Biological Control 9 (2): 128 - 129.
- Vera, M.L., Valverde, L., Popich, S.B. and Ajmat De Toledo, Z.D. (1995).** Preliminary evaluation of natural enemies of *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) in Tucuman, Argentina. Acta Entomologica Chilena 19: 135 - 141.
- Verhoog, M.D., Brakefield, P.M. and Van Boven, A. (1996).** Melanic moths and the ability to encapsulate parasitoid eggs and larvae. Proceedings of the 7th meeting of Experimental and Applied Entomologists in the Netherlands on 15 December 1995. Proceedings of the

- Section Experimental and Applied Entomology of the Netherlands Entomological Society 7: 127 - 133.
- Verkerk, R.H.J. and Wright, D.J. (1993).** Biological activity of neem seed kernel extracts and synthetic azadirachtin against larvae of *Plutella xylostella*. *Pesticide Science* 37 (1): 83 - 91.
- Verkerk, R.H.J. and Wright, D.J. (1997).** Field-based studies with the diamondback moth tritrophic system in Cameron Highlands of Malaysia: implications for pest management. *International Journal of Pest Management* 43 (1): 27 - 33.
- Verma, A.K. and Makhmoor, H.D. (1989).** Development of life table for *Metasyrphus confrater* (Wiedemann) (Diptera: Syrphidae), a predator of the cabbage aphid (Homoptera: Aphididae) in cauliflower crop ecosystem. *Entomon* 14 (3-4): 227 - 232.
- Verma, R.R. and Joshi, R. (1988).** Studies on the natural enemies of insect pests of temperate region. *Progressive Horticulture* 20 (3-4): 335 - 336.
- Verma, T.D., Verma, K.S. and Karan, R. (1996).** Natural enemies of *Dioryctria abietella* Denis & Schiff., a pest of conifers in the north-west Himalaya. *Pest Management and Economic Zoology* 4 (1-2): 115 - 117.
- Verma, T.S. and Lal, OP. (1985).** A new record of *Itamoplex* sp. (Hymenoptera: Ichneumonidae) parasitizing eggplant shoot and fruit borer in Kulu valley, Himachal Pradesh. *Bulletin of Entomology* 26 (2): 219 - 222.
- Viafara-Millan, H.F., Roa, F.G. and Diaz, A.E. (1999).** Natural parasitism of *Neoleucinodes elegantalis* (Lepidoptera: Pyralidae) in producers zones of Solanaceae from Cauca and Valle del Cauca, Colombia. *Revista Colombiana de Entomologia* 25 (3-4): 151 - 159.
- Vikberg, V. (1999).** *Lathroplex anthreni* sp. n. (Hymenoptera: Ichneumonidae), a parasitoid on *Anthrenus museorum* (Linnaeus) in southern Finland. *Entomologica Fennica* 10 (4): 239 - 244.
- Viktorov, G.A. (1970).** *Aubertiana* Viktorov, a new name for *Aubertia* Viktorov (Hymenoptera, Ichneumonidae). (in Russian) *Zoologicheskii Zhurnal* 49: 308-309.
- Vilkamaa, P. and Komonen, A. (2001).** Redescription and biology of *Trichosia* (Baeosciara) sinuata Menzel & Mohrig (Diptera: Sciaridae). *Entomologica Fennica* 12 (1): 46 - 49.
- Vincelette, F. (1994).** The utilisation of parasitoid wasps in biological control. *Faberies* 19 (1): 12 - 26.
- Vinson, S.B. (1988).** Biochemical interrelations between plants, herbivores and parasitoids. *Colloques del'INRA* 48: 23 - 25.
- Vinson, S.B. (1990).** The interaction of two parasitoid polydnviruses with the endocrine system of *Heliothis virescens*. *Proceedings and abstracts, Vth International Colloquium on Invertebrate Pathology and Microbial Control, Adelaide, Australia, 20-24 August 1990.* 195 - 199.
- Vinson, S.B., Hegazi, E.M., Pennacchio, F., Bradleigh-Vinson, S. and Lanzrein, B. (1998).** A possible mechanism for the physiological suppression of conspecific eggs and

- larvae following superparasitism by solitary endoparasitoids. Parasitoid-host interactions at the physiological and molecular level. *Journal of Insect Physiology* 44 (9): 703 - 712.
- Vinson, S.B., Williams, H.J. and Lu, J. (1994).** Identification of different compounds from different plants responsible for the orientation of *Campoletis sonorensis* to potential host sites. *Norwegian Journal of Agricultural Sciences SUPP16*: 207 - 210.
- Vinson, S.B. and Hu, J.S. (2000).** Interaction between the larval endoparasitoid *Campoletis sonorensis* (Hymenoptera: Ichneumonidae) and its host the tobacco budworm (Lepidoptera: Noctuidae). *Annals of the Entomological Society of America* 93 (2): 220 - 224.
- Vinson, S.B. and Williams, H.J. (1991).** Host selection behaviour of *Campoletis sonorensis*: a model study. *Biological Control* 1 (2): 107 - 117.
- Vinuela, E., Medina, M.P., Schneider, M., Gonzalez, M., Budia, F., Adan, A., Del Estal, P., Vogt, H., Vinuela, E. and Jacas, J. (2001).** Comparison of side-effects of spinosad, tebufenozide and azadirachtin on the predators *Chrysoperla carnea* and *Podisus maculiventris* and the parasitoids *Opius concolor* and *Hyposoter didymator* under laboratory conditions. IOBC-WPRS Working Group 'Pesticides and Beneficial Organisms'. Proceedings of the meeting at Castello de la Plana, Spain, 18-20 October, 2000. *Bulletin OILB SROP* 24 (4): 25 - 34.
- Virla, E.G., Colomo, M.V., Berta, C. and Valverde, L. (1999).** The complex of parasitoids of fall armyworm of maize, *Spodoptera frugiperda*, in Argentina (Insecta: Lepidoptera). *Neotropica* 45: 113 - 114, 3 - 12.
- Visalakshy, P.N.G. and Jayanth, K.P. (1990).** Studies on the biology and seasonal abundance of *Hypena laceratalis* Walker (Lepidoptera: Noctuidae) on *Lantana camara* L. in India. *Entomon* 15 (3-4): 231 - 234.
- Visitpanich, J. (1994).** The parasitoid wasps of the coffee stem borer, *Xylotrechus quadripes* Chevrolat (Coleoptera: Cerambycidae) in northern Thailand. *Japanese Journal of Entomology* 62 (3): 597 - 606.
- Vogt, H. (1997).** The importance of using selective insecticides against key pests in apple orchards to preserve the parasitoid fauna of leafminers. *Entomological research in organic agriculture. Selected papers from the European Workshop, Austrian Federal Ministry of Science and Research, Vienna, Austria, 14-16 March, 1995. Biological Agriculture and Horticulture* 15 (1-4): 241 - 248.
- Volkoff, A.N., Beliveau, C., Rocher, J., Hilgarth, R., Levasseur, A., Duonor-Cerutti, M., Cusson, M. and Webb, B.A. (2002).** Evidence for a conserved polydnavirus gene family: Ichnovirus homologs of the CsIV repeat element genes. *Virology* 300 (2): 316 - 331.
- Volkoff, A.N., Cerutti, P., Rocher, J., Ohresser, M.C.P., Devauchelle, G. and Duonor-Cerutti, M. (1999).** Related RNAs in lepidopteran cells after in vitro infection with *Hyposoter didymator* virus define a new polydnavirus gene family. *Virology New York* 263 (2): 349 - 364.

- Volkoff, A.N., Ravallec, M., Bossy, J.P., Cerutti, P., Rocher, J., Cerutti, M. and Devauchelle, G. (1995).** The replication of *Hyposoter didymator* polydnavirus: cytopathology of the calyx cells in the parasitoid. *Biology of the Cell* 83 (1): 1 - 13.
- Volkoff, A.N., Rocher, J., Cerutti, P., Ohresser, M.C.P., d'-Aubenton-Carafa, Y., Devauchelle, G. and Duonor-Cerutti, M. (2001).** Persistent expression of a newly characterized *Hyposoter didymator* polydnavirus gene in long-term infected lepidopteran cell lines. *Journal of General Virology* 82 (4): 963 - 969.
- Vrablova, M., Toth, P. and Cagan, L. (2002).** *Diadegma angitiaeforma* (Hymenoptera, Ichneumonidae) - first records from Slovakia and a new host for the species. *Biologia Bratislava* 57 (5): 630.

## W

- Wacker, K., Bartsch, P. and Clemen, G. (2001).** The development of the tooth pattern and dentigerous bones in *Polypterus senegalus* (Cladistia: Actinopterygii). *Annals of Anatomy* 183 (1): 37 - 52.
- Wackers, F.L., Bjornsen, A. and Dorn, S. (1996).** A comparison of flowering herbs with respect to their nectar accessibility for the parasitoid *Pimpla turionellae*. *Proceedings of the 7th meeting of Experimental and Applied Entomologists in the Netherlands on 15 December 1995. Proceedings of the Section Experimental and Applied Entomology of the Netherlands Entomological Society* 7: 177 - 182.
- Wackers, F.L., Mitter, E. and Dorn, S. (1998).** Vibrational sounding by the pupal parasitoid *Pimpla* (*Coccygomimus*) *turionellae*: an additional solution to the reliability-detectability problem. *Biological Control* 11 (2): 141 - 146.
- Waern, P. and Ekbohm, B. (1995).** The diamondback moth (*Plutella xylostella*) as an insect pest in spring oilseed rape. *Vaxtskyddsnotiser* 59 (4): 111 - 115.
- Wagner, D.L., Tobi, D.R., Parker, B.L., Wallner, W.E. and Leonard, J.G. (1989).** Immature stages and natural enemies of *Korscheltellus gracilis* (Lepidoptera: Hepialidae). *Annals of the Entomological Society of America* 82 (6): 717 - 724.
- Wahl, D.B. (1987).** A revision of *Venturia* north of Central America (Hymenoptera: Ichneumonidae). *University of Kansas Science Bulletin* 53: 275-356.
- Wahl, D.B. (1988).** A review of the mature larvae of the Banchini and their phylogenetic significance, with comments on the Stilbopinae (Hymenoptera: Ichneumonidae). *In* V.K. Gupta (ed.), *Advances in Parasitic Hymenoptera Research*. New York: E.J. Brill. 546 pp.
- Wahl, D. (1990).** A review of the mature larvae of Diplazontinae, with notes on larvae of Acaenitinae and Orthocentrinae and proposal of two new subfamilies (Insecta: Hymenoptera, Ichneumonidae). *Journal of Natural History* 24 (1): 27 - 52.
- Wahl, D.B. (1991).** A new species of *Dusona* from New Zealand, and the application of *Dusona* vs. *Delopia* (Hymenoptera: Ichneumonidae: Campopleginae). *Proceedings of the Entomological Society of Washington* 93 (4): 946 - 950.

- Wahl, D.B. (1991).** The status of Rhimphoctona, with special reference to the higher categories within Campopleginae and the relationships of the subfamily (Hymenoptera: Ichneumonidae). *Transactions of the American Entomological Society* 117 (3-4): 193 - 213.
- Wahl, D.B. (1993).** Cladistics of the genera of Labeninae (Hymenoptera: Ichneumonidae). *Entomologia Generalis* 18: 91-105.
- Wahl, D.B. (1993).** Cladistics of the genera of Mesochorinae (Hymenoptera: Ichneumonidae). *Systematic Entomology* 18 (4): 371 - 387.
- Wahl, D.B. (1996).** Further notes on the family-group names of the Ichneumoninae (Hymenoptera: Ichneumonidae). *Journal of Hymenoptera Research* 5: 273.
- Wahl, D.B. (1996).** Xenothyriini, a new tribe of Labeninae, and a key to the tribes and genera of Labeninae (Hymenoptera: Ichneumonidae). *Proceedings of the Entomological Society of Washington* 98: 158-161.
- Wahl, D.B. (1997).** The cladistics of the genera and subgenera of Xoridinae. *In*, Gauld, I.D., *The Ichneumonidae of Costa Rica, 2. Memoirs of the American Entomological Institute* 57: 454-460.
- Wahl, D. B. and M. J. Sharkey. (1993).** Superfamily Ichneumonoidea. pp. 358-362. *In*: H. Goulet and J.T. Huber (eds.), *Hymenoptera of the world, an identification guide to families*. Agriculture Canada Research Branch Monograph No. 1894E, 668pp.
- Wahl, D.B., Mason, W.R.M. and Townes, H. (1995).** The family-group names of the Ichneumoninae (Hymenoptera: Ichneumonidae). *Journal of Hymenoptera Research* 4: 285 - 293.
- Wahl, D.B. and Gauld, I.D. (1998).** The cladistics and higher classification of the Pimpliformes (Hymenoptera: Ichneumonidae). *Systematic Entomology* 23: 101-134.
- Wahl, D.B. and Gauld, I.D. (1998).** The cladistics and higher classification of the Pimpliformes (Hymenoptera: Ichneumonidae). *Systematic Entomology* 23 (3): 265 - 298.
- Wahl, D.B. and Gupta, V.K. (1988).** A review of the mature larvae of the Banchini and their phylogenetic significance, with comments on the Stilbopinae (Hymenoptera: Ichneumonidae). *Advances in parasitic Hymenoptera research: Proceedings of the II. Conference on the Taxonomy and Biology of Parasitic Hymenoptera. recd. 1993*, 147 - 161.
- Wakisaka, S., Tsukuda, R. and Nakasuji, F. (1991).** Life tables of the diamondback moth, *Plutella xylostella* (L.) (Lepidoptera: Yponomeutidae) and effects of rainfall, temperature and host plants on survival and reproduction. *Japanese Journal of Applied Entomology and Zoology* 35 (2): 115 - 122.
- Waladde, S.M., Leutle, M.F. and Villet, M.H. (2001).** Parasitism of *Plutella xylostella* (Lepidoptera, Plutellidae): field and laboratory observations. *South African Journal of Plant and Soil* 18 (1): 32 - 37.

- Walter, S. (1991).** Ichneumonidae (Hymenoptera) in pollution-damaged spruce forests of the Harz Mountains. *Zoologische Jahrbucher, Abteilung für Systematik, Ökologie und Geographie der Tiere* 118 (3-4): 449 - 464.
- Wan, S.F. (1989).** A new genus and two new species of Acaenitinae from China (Hymenoptera: Ichneumonidae). *Acta Entomologica Sinica* 32 (3): 357 - 360.
- Wan, Z.J. and Wang, T.K. (1989).** A study on *Arge nipponensis* Rohwer. *Insect Knowledge* 26 (2): 85 - 88.
- Wang, S.F. (1987).** A new genus and species of the Tryphoninae from China (Hymenoptera: Ichneumonidae). (in Chinese) *Sinozoologica* 5: 153-154.
- Wang, W.X. (1987).** Investigation of parasites of *Buzura suppressaria* Guenee. *Chinese Journal of Biological Control* 3 (1): 46.
- Wang, S.F. (1989).** A new genus and two new species of Acaenitinae from China (Hymenoptera: Ichneumonidae). (in Chinese) *Acta Entomologica Sinica* 32: 357-360.
- Wang, S.F. (1989).** Notes on Chinese *Phalgea* Cameron (Hymenoptera: Ichneumonidae, Acaenitinae). *Acta Entomologica Sinica* 32 (2): 228 - 229.
- Wang, C.Z., Yang, Q.H. and Zhou, M.Z. (1997).** Effects of gossypol on growth of the cotton bollworm and development of its parasitoid *Campoletis chloridae*. *Entomologia Sinica* 4 (2): 182 - 188.
- Wang, G., Wei, H.X., Zhao, M.G., Shi, J.L., Pan, H.Y. and Wan, Y.Y. (1992).** A report on parasite natural enemies of Xingan larch casebearer (*Coleophora dahurica* Flkv). Collection of achievements on the technique cooperation project of P. R. China and F. R. Germany 21 - 24.
- Wang, Ge., Wang, G., Sun, J.W., Ma, S.C. and Sheng, M.L. (1996).** A new species of *Endasys* Forester (Hymenoptera: Ichneumonidae) parasitizing *Pachynematus itoi* Okutani (Hymenoptera: Tenthredinidae). *Entomotaxonomia* 18 (3): 230 - 232.
- Wang, H.J., Wu, Z.Y., Lu, C.F., Li, G.P. and Xu, T.S. (2000).** A study on bionomics of bamboo puss moth, *Norraca retrofusca* (Notodontidae: Lepidoptera) in China. *Forest Research Beijing* 13 (6): 583 - 588.
- Wang, Q., Shi, G.L. and O'-Callaghan, M. (1999).** Parasitic natural enemies of lemon tree borer. Proceedings of the Fifty Second New Zealand Plant Protection Conference, Auckland Airport Centra, Auckland, New Zealand, 10 – 12 August, 1999. 60 - 64.
- Wang, R.Y., Zhang, N.Q., Wang, B. and Li, Z.H. (1996).** A brief report on the genus *Rhyssa* Gravenhorst in China (Hymenoptera: Ichneumonidae: Pimplinae). *Journal of Shandong Agricultural University* 27 (3): 249 - 257.
- Wang, S.F. and Zhang, L.X. (1989).** A new species of the genus *Habronyx* [*H. majorocellus*] from Liaoning (Hymenoptera: Ichneumonidae: Anomalinae). *Acta Entomologica Sinica* 32 (4): 480 - 481.
- Wang, W.X. and Liu, W.Q. (1993).** Field surveys on the pupal parasitism of *Dendrolimus punctatus* (Lep.: Lasiocampidae) in Guangxi and Hunan. *Chinese Journal of Biological Control* 9 (3): 143 - 144.

- Wang, X.G., Liu, S.S. and Chen, X.O. (1997).** Host age preference and suitability of *Diadromus collaris*, a major pupal parasite of *Plutella xylostella*. Chinese Journal of Biological Control 13 (3): 101 - 105.
- Wang, X.G. and Liu, S.S. (1998).** Bionomics of *Diadromus collaris* (Hymenoptera: Ichneumonidae), a major pupal parasitoid of *Plutella xylostella*. Acta Entomologica Sinica 41 (4): 389 - 395.
- Wang, S.-F. and Yao J. (1994).** A new genus and species of the Hemigasterinae from China (Hymenoptera: Ichneumonidae). (in Chinese) Sinozoologica 11: 175-178.
- Wang, S. and Yan. J. (1998).** A new genus and species of Pionini from China (Hymenoptera: Ichneumonidae, Ctenopelmatinae). Scientia Silvae Sinicae 34: 42-44.
- Wang, Y.K., Yin, C.L., Bai, Z.Q., Zhang, Y.R., Li, J.J. and Yang, Q.S. (2000).** A preliminary study on the use and protection of the natural enemies of insect pests in the seed stands of *Picea crassifolia*. Forest Research Beijing 13 (4): 416 - 422.
- Wardle, A.R. (1990).** Learning of host microhabitat colour by *Exeristes roborator* (F.) (Hymenoptera: Ichneumonidae). Animal Behaviour 39 (5): 914 - 923.
- Wardle, A.R. and Borden, J.H. (1989).** Learning of an olfactory stimulus associated with host microhabitat by *Exeristes roborator*. Entomologia Experimentalis et Applicata 52 (3): 271 - 279.
- Wardle, A.R. and Borden, J.H. (1990).** Learning of host microhabitat form by *Exeristes roborator* (F.) (Hymenoptera: Ichneumonidae). Journal of Insect Behavior 3 (2): 251 - 263.
- Wardle, A.R. and Borden, J.H. (1990).** Maturation feeding without learning in adult *Exeristes roborator* (Hymenoptera: Ichneumonidae). Journal of Economic Entomology 83 (1): 126 - 130.
- Wardle, A.R. and Borden, J.H. (1991).** Effect of prior experience on the response of *Exeristes roborator* (Hymenoptera: Ichneumonidae) to a natural host and microhabitat in a seminatural environment. Environmental Entomology 20 (3): 889 - 898.
- Washburn, J.O., Kirkpatrick, B.A. and Volkman, L.E. (1996).** Insect protection against viruses. Nature London. 383: 6603, 767.
- Watanabe, M. and Hirai, Y. (1998).** A solitary endoparasitic wasp, *Bathyplectes* sp. (Hymenoptera: Ichneumonidae), of the clover leaf weevil, *Hypera punctata* and its parasitism in Japan. Proceedings of the Kanto Tosan Plant Protection Society 45: 235 - 237.
- Watt, A.D., Stork, N.E. and Hunter, M.D. (1997).** Forests and insects. Chapman & Hall Ltd; London; UK, xv + 406 pp.
- Watt, A.D. (1989).** The growth and survival of *Panolis flammea* larvae in the absence of predators on Scots pine and lodgepole pine. Ecological Entomology 14 (2): 225 - 234.
- Wayman, L. D. (1994).** Spatial distribution and sex ratios of parasitic Hymenoptera (Ichneumonidae: Campopleginae and Ichneumoninae; Braconidae: *Aphidius ervi*) in a disturbed Georgia piedmont landscape. M. S. thesis, Univ. of Georgia, Athens, 123pp.
- Weaver, J.E., Zivkovich, C.A., Baniecki, J.F., Yohn, G.W., Smith, B.D. and**

- Workman, D.J. (1994).** Biological agents associated with the alfalfa weevil - a survey of Jefferson, Grant, and Hardy Counties, West Virginia, for 1993. Circular Agricultural and Forestry Experiment Station, West Virginia University 160: 18 pp.
- Weaver, J.E. and Shaffer, J.B. (1988).** Biocontrol of the alfalfa weevil in West Virginia - a status report. Bulletin, Agricultural and Forestry Experiment Station, West Virginia University. 700: 13 pp.
- Webb, B.A., Cui, L.W., Pennacchio, F., Bradleigh-Vinson, S. and Lanzrein, B. (1998).** Relationships between polydnavirus genomes and viral gene expression. Parasitoid-host interactions at the physiological and molecular level. *Journal of Insect Physiology* 44 (9): 785 - 793.
- Webb, B.A. and Cui, L.W. (1996).** Isolation and characterization of a member of the cysteine-rich gene family from *Campoletis sonorensis* polydnavirus. *Journal of General Virology* 77 (4): 797 - 809.
- Webb, B.A. and Luckhart, S. (1994).** Evidence for an early immunosuppressive role for related *Campoletis sonorensis* venom and ovarian proteins in *Heliothis virescens*. *Archives of Insect Biochemistry and Physiology* 26 (2-3): 147 - 163.
- Webb, B.A. and Luckhart, S. (1996).** Factors mediating short- and long-term immune suppression in a parasitized insect. *Journal of Insect Physiology* 42 (1): 33 - 40.
- Webb, B.A. and Summers, M.D. (1990).** Venom and viral expression products of the endoparasitic wasp *Campoletis sonorensis* share epitopes and related sequences. *Proceedings of the National Academy of Sciences of the United States of America*. 87: 13, 4961 - 4965.
- Webb, B.A. and Summers, M.D. (1992).** Stimulation of polydnavirus replication by 20-hydroxyecdysone. *Experientia* 48 (10): 1018 - 1022.
- Weeks, P.J.D., Gauld, I.D., Gaston, K.J. and O'Neill, M.A. (1997).** Automating the identification of insects: a new solution to an old problem. *Bulletin of Entomological Research* 87 (2): 203 - 211.
- Weinstein, P. and Austin, A.D. (1991).** The host relationships of trigonalid wasps (Hymenoptera: Trigonalidae), with a review of their biology and catalogue to world species. *Journal of Natural History* 25 (2): 399 - 433.
- Weiss, E. and Stettmer, C. (1991).** Weeds in agricultural ecosystems attract flower-visiting beneficial insects. *Agrarokologie* 1: 104 pp., 7 pp.
- Wells, M.L., Hagan, D.V. and McPherson, R.M. (2001).** Survey of parasitoids associated with *Rhyacionia frustrana* (Comstock) in Bulloch County, Georgia. *Journal of Entomological Science* 36 (1): 101 - 104.
- West, R.J. (1991).** Notes on the biology and control of the black army cutworm, *Actebia fennica* (Lepidoptera: Noctuidae), in black spruce plantations. *Proceedings of the Entomological Society of Ontario* 122: 53 - 63.
- West, R.J., Dixon, P.L., Quednau, F.W., Lim, K.P. and Hiscock, K. (1994).** Establishment of *Olesicampe geniculatae* Quednau and Lim (Hymenoptera: Ichneumonidae) to control



- the mountain ash sawfly, *Pristiphora geniculata* (Hartig) (Hymenoptera: Tenthredinidae), in Newfoundland. Canadian Entomologist 126 (1): 7 - 11.
- West, R.J., Kenis, M., Butt, G.W. and Bennett, S.M. (1999).** Parasitoid complex of *Zeiraphera canadensis* (Lepidoptera: Tortricidae) and evaluation of *Tycherus osculator* (Hymenoptera: Ichneumonidae) as a biological control agent. Canadian Entomologist 131 (4): 465 - 474.
- Whiteman, N.K. and Landwer, B.H.P. (2000).** Parasitoids reared from *Polistes* (Hymenoptera: Vespidae: Polistinae) nests in Missouri, with a state record of *Elasmus polistis* Burks (Hymenoptera: Elasmidae). Journal of the Kansas Entomological Society 73 (3): 186 - 188.
- Whitfiel, J.B., Suh, K.I. and Lee, J.W. (1999).** Phylogeny of the subfamily Mesochorinae (Hymenoptera: Ichneumonidae) based on partial mitochondrial DNA sequences. Korean Journal of Entomology 29 (1): 31 - 35.
- Whitfield, J.B., Johnson, N.F. and Hamerski, M.R. (1989).** Identity and phylogenetic significance of the metapostnotum in nonaculeate Hymenoptera. Annals of the Entomological Society of America 82 (6): 663 - 673.
- Whitman, D.W. and Eller, F.J. (1990).** Parasitic wasps orient to green leaf volatiles. Chemoecology 1 (2): 69 - 75.
- Wieber, A.M., Cook, S.P., Webb, R.E., Tatman, K.M. and Reardon, R.C. (1995).** Niche partitioning by four *Gelis* spp. (Hymenoptera: Ichneumonidae) hyperparasitoids of the primary gypsy moth parasitoid *Cotesia melanoscela* (Hymenoptera: Braconidae). Annals of the Entomological Society of America 88 (4): 427 - 433.
- Wieber, A.M., Webb, R.E., Reardon, R.C., Tatman, K.M. and Thorpe, K.W. (2002).** Temporal patterns of hyperparasitoid emergence from *Cotesia melanoscelus* (Hymenoptera: Braconidae) cocoons. Entomological News 113 (4): 253 - 266.
- Wiech, K., Jankowska, B., Finch, S., Hartfield, C. and Brunel, E. (1999).** Preliminary observations on *Diadegma fenestralis* a parasitoid of the diamond-back moth, *Plutella maculipennis*. Integrated control in field vegetable crops. Proceedings of the meeting at Chania, Crete, 6-8 October, 1997. Bulletin OILB SROP 22 (5): 145 - 149.
- Wiech, K. and Kalmuk, J. (2002).** The role of parasitoids in decreasing the number of diamond back moth (*Plutella xylostella*) in horticultural crops. The BCPC Conference: Pests and diseases, Volumes 1 and 2. Proceedings of an international conference held at the Brighton Hilton Metropole Hotel, Brighton, UK, 18-21 November 2002. 329 - 334.
- Wiech, K. and Pniak, M. (2002).** Parasitoids of leaf roller caterpillars occurring on common privet hedgerows. Progress in Plant Protection 42 (2): 464 - 466.
- Williams, D.W., Fuester, R.W., Metterhouse, W.W., Balaam, R.J., Bullock, R.H., Chianese, R.J. and Reardon, R.C. (1992).** Incidence and ecological relationships of parasitism in larval populations of *Lymantria dispar* (Lepidoptera: Lymantriidae). Biological Control 2 (1): 35 - 43.

- Williams, R. and Randolph, S. (2002).** Insect and invertebrate group the rose bedeguar project: rearing & recognising inhabitants. *Cecidology* 17 (1): 17 - 24.
- Wilson, J.A.Jr. and DuRant, J.A. (1991).** Parasites of the European corn borer (Lepidoptera: Pyralidae) in South Carolina. *Journal of Agricultural Entomology* 8 (2): 109 - 116.
- Winiarska, W. (1995).** *Apanteles melanoscelis* Ratz. - an effective parasitoid of gypsy moth caterpillars. *Ochrona Roslin* 39 (2): 6.
- Wnuk, A. and Wojciechowicz, E. (1993).** Syrphid predators (Diptera: Syrphidae) occurring in colonies of the cabbage aphid, *Brevicoryne brassicae* (L.), on cabbage and cauliflower. *Polskie Pismo Entomologiczne* 62 (1-4): 215 - 229.
- Wolf, R. and Wolf, D. (1988).** Activation by calcium ionophore injected into unfertilized ovarian eggs explanted from *Pimpla turionellae* (Hymenoptera). *Zoologische Jahrbucher, Abteilung fur Allgemeine Zoologie und Physiologie der Tiere* 92 (4): 501 - 512.
- Wright, L.M. (1995).** Seasonality of the Ichneumonoidea and alate Formicidae in a tropical moist forest at Barro Colorado Island, Panama. Senior honors thesis, Univ. of Georgia, Athens, 85pp.
- Wright, D.J., Waage, J.K. and Gu, D.J. (1995).** Effects of sublethal doses of insecticides on the mating behaviour of the male parasitoid, *Diadegma eucerophagum*. *Journal of South China Agricultural University* 16 (2): 55 - 59.
- Wright, D.J., Waage, J.K. and Gu, D.J. (1995).** Influence of sublethal effects of insecticides on the population dynamics of *Plutella xylostella* and its parasitoid *Diadegma eucerophaga* in the field. *Entomologia Sinica* 2 (1): 77 - 82.
- Wyler, T. and Lanzrein, B. (2003).** Ovary development and polydnavirus morphogenesis in the parasitic wasp *Chelonus inanitus*. II. Ultrastructural analysis of calyx cell development, virion formation and release. *Journal of General Virology* 84 (5): 1151 - 1163.
- Wytwer, J. and Sawoniewicz, J. (1997).** Diversity indices of double classification in the study of Ichneumonidae (Hymenoptera: Ichneumonidae) communities. *Fragmenta Faunistica* 41: 10 - 16, 151 - 165.

## X

- Xiao, T.G., Song, H.Y. and Chen, C.M. (1992).** A survey on the parasitic insects of the larvae of the Chinese tea oecophorid (*Casmara patrona* Meyrick). *Journal of Hunan Agricultural College* 18 (2): 288 - 292.
- Xie, G.F., Li, X., Bi, H.M., Han, G.Y. and Xie, M. (1993).** A study on the bionomics of two species of Zeiraphera Treit. on larch. *Entomological Knowledge* 30 (5): 280 - 282.
- Xu, D.M. and Stoltz, D. (1991).** Evidence for a chromosomal location of polydnavirus DNA in the ichneumonid parasitoid *Hyposoter fugitivus*. *Journal of Virology* 65 (12): 6693 - 6704.

- Xu, J.X., Shelton, A.M. and Cheng, X.N. (2001).** Variation in susceptibility of *Diadegma insulare* (Hymenoptera: Ichneumonidae) to permethrin. *Journal of Economic Entomology* 94 (2): 541 - 546.
- Xu, J.X., Wu, J.C. and Cheng, X.N. (2001).** A method of rearing of *Diadegma insulare* in greenhouse. *Entomological Knowledge* 38 (4): 311 - 313.

## Y

- Yabas, C. and Zeren, O. (1987).** Studies on the biology of the cabbage webworm (*Hellula undalis* F.) (Lepidoptera: Pyralidae). *Turkiye I. Entomoloji Kongresi Bildirileri*, 13-16 Ekim 1987, Ege Universitesi, Nornova, Izmir 229 - 238.
- Yadav, K.P. and Shrivastava, S.K. (1990).** *Campoletis chlorideae* (Uchida) (Hymenoptera: Ichneumonidae) a new host of *Brachymeria secundaria* (Ruschka) (Hymenoptera: Chalcididae). *Entomon* 15 (3-4): 287.
- Yadava, C.P. and Lal, S.S. (1988).** Relationship between certain abiotic and biotic factors and the occurrence of gram pod borer, *Heliothis armigera* (Hbn.) on chickpea. *Entomon* 13 (3-4): 269 - 273.
- Yakovlev, Y.B. (1993).** Mushrooms and insects associated with them in Petrozavodsk city parks. *Aquilo Ser. Botanica* 31: 131 - 136.
- Yamamoto, Y., Otori, M., Ohbayashi, T., Iwabuchi, K. and Mitsuhashi, J. (1997).** In vitro rearing of the larval endo-parasitoid, *Venturia canescens* (Gravenhorst) (Hymenoptera: Ichneumonidae) II. Larval development. *Applied Entomology and Zoology* 32 (1): 256 - 258.
- Yamane, S. (1990).** A revision of the Japanese Eumenidae (Hymenoptera: Vespoidea). *Insecta Matsumurana* 43: 189 pp., 9 pp.
- Yang, G.H. (1998).** The preliminary study on *Lycorina spilonotae* Chao. *Journal of Henan Agricultural Sciences* 9: 20 - 21.
- Yang, J.C., Chu, Y.I. and Talekar, N.S. (1993).** Biological studies of *Diadegma semiclausum* (Hym.: Ichneumonidae), a parasite of diamondback moth. *Entomophaga* 38 (4): 579 - 586.
- Yang, J.C., Chu, Y.I. and Talekar, N.S. (1994).** Studies on the characteristics of parasitism of *Plutella xylostella* (Lep.: Plutellidae) by a larval parasite *Diadegma semiclausum* (Hym.: Ichneumonidae). *Entomophaga* 39 (3-4): 397 - 406.
- Yang, M.X. (1993).** Three species of ichneumons parasitizing the eggs of spiders. *Plant Protection* 19 (5): 46 - 48.
- Yasarakinci, N. and Kornosor, S. (1990).** The effectiveness of the natural enemies and parasitoids of *Heliothis virescens* (Hufn.) (Lep.: Noctuidae) a pest of chickpea and lentil in southeastern Anatolia. *Proceedings of the Second Turkish National Congress of Biological Control* 83 - 89.
- Yastrebov, I.O. (1986).** *Hyposoter* - a parasitoid of the cabbage white butterfly. *Zashchita Rastenii* 8: 47 - 48.

- Yastrebov, I.O. (1991).** A new method of determining the harmful effect of the cabbage moth. *Vestnik Sel'skokhozyaistvennoi Nauki Moskva* 7: 141 - 144.
- Yastrebov, I.O. (1991).** Nectar plants near cabbage plantings. *Zashchita Rastenii* 1: 17.
- Yastrebov, I.O. (1992).** Effective parasite of *Barathra brassicae*. *Zashchita Rastenii Moskva* 11: 27.
- Yasuhara, A., Momoi, S. and Kondo, H. (1998).** Photoperiodic control of diapause in five species of *Coccygomimus* (Hymenoptera: Ichneumonidae) from Hokkaido, Japan. *Japanese Journal of Applied Entomology and Zoology* 42 (3): 123 - 128.
- Yasuhara, A. and Momoi, S. (1997).** Photoperiodic response and life cycle of *Coccygomimus luctuosus* Smith (Hymenoptera: Ichneumonidae). *Japanese Journal of Applied Entomology and Zoology* 41 (3): 133 - 139.
- Yeargan, K.V. and Braman, S.K. (1989).** Comparative behavioral studies of indigenous hemipteran predators and hymenopteran parasites of the green cloverworm (Lepidoptera: Noctuidae). *Journal of the Kansas Entomological Society* 62 (2): 156 - 163.
- Yeargan, K.V. and Braman, S.K. (1989).** Life history of the hyperparasitoid *Mesochorus discitergus* (Hymenoptera: Ichneumonidae) and tactics used to overcome the defensive behavior of the green cloverworm (Lepidoptera: Noctuidae). *Annals of the Entomological Society of America* 82 (3): 393 - 398.
- Yin, L.H., Zhang, C., Qin, J.D. and Wang, C.Z. (2003).** Polydnavirus of *Campoletis chloridae*: characterization and temporal effect on host *Helicoverpa armigera* cellular immune response. *Archives of Insect Biochemistry and Physiology* 52 (2): 104 - 113.
- Yitaferu, K. and Gebre-Amlak, A. (1994).** Phenology and survival of the maize stalk borer, *Busseola fusca* in eastern Ethiopia. *Insect Science and its Application* 15 (2): 177 - 184.
- Yonce, C.E., Tedders, W.L. and Wood, B.W. (1996).** Cold tolerance of hickory shuckworm (Lepidoptera: Tortricidae) larvae and associated parasites. *Journal of Entomological Science* 31 (1): 13 - 19.
- You, L.S., Lei, R.H., Jiang, J.X., Bo, L.Y. and Xiao, Z.S. (2002).** Bionomic of *Campoletis chloridae* (Hym.: Ichneumonidae) as a parasitoid of the cotton bollworm *Helicoverpa armigera* (Lep.: Noctuidae). *Entomologia Sinica* 9 (3): 29 - 37.
- You, L.S., Li, J.W., Xiong, S.L. and Chen, Z.F. (1999).** Investigation on the parasitoid wasps of Siberian cocklebur stem borer [*Ostrinia orientalis*] and study on the ichneumon wasp *Trathala flavoorbitalis*. *Wuyi Science Journal* 15: 55 - 63.
- Yu, D.S., Kokko, E.G., Barron, J.R., Schaalje, G.B. and Gowen, B.E. (1992).** Identification of ichneumonid wasps using image analysis of wings. *Systematic Entomology* 17 (4): 389 - 395.
- Yu, D.S. and Horstmann, K. (1997).** A Catalogue of World Ichneumonidae (Hymenoptera). The American Entomological Institute. Gainesville, Florida.
- Yuan, G.H., Sun, T.S., Zheng, X.J. and Zheng, X.Y. (2000).** Study on the biological characters of *Lycorina spilonatae* Chao and its utilization. *China Fruits* 1: 29 - 31.

## Z

- Zakhals, C.J. (1989).** The Acaenitinae of the Netherlands and the host of *Acaenitus dubitator* (Hymenoptera: Ichneumonidae). *Entomologische Berichten* 49 (2): 21 - 25.
- Zaman, M. and Karimullah. (1987).** Lepidoptera of jute cultivars in Peshawar. *Pakistan Journal of Agricultural Research* 8 (3): 290 - 297.
- Zandigiaco, P. (1993).** *Choristoneura lafauryana* (Rag.) on soyabean in Friuli. *Informatore Fitopatologico* 43 (6): 46 - 50.
- Zanuncio, J.C., Teixeira, C.A.D. and Sossai, M.F. (1999).** Natural enemies of *Nomophila* sp. (Lepidoptera: Pyralidae), a cutworm of *Eucalyptus grandis* (Myrtaceae) seedlings in Vicoso, Minas Gerais, Brazil. *Anais da Sociedade Entomologica do Brasil* 28 (2): 357 - 358.
- Zapryanov, A.A. (1992).** *Anthonomus pyri* Kalt (Coleoptera: Curculionidae) as a pest problem in pear orchards and the role of natural enemies for reducing its population. *Acta Phytopathologica et Entomologica Hungarica* 27 (1-4): 687 - 690.
- Zhao, J.Z., Ayers, G.S., Grafius, E.J. and Stehr, F.W. (1992).** Effects of neighboring nectar-producing plants on populations of pest Lepidoptera and their parasitoids in broccoli plantings. *Great Lakes Entomologist* 25 (4): 253 - 258.
- Zhou, G.F., Overholt, W.A. and Kimani-Njogu, S.W. (2003).** Species richness and parasitism in an assemblage of parasitoids attacking maize stem borers in coastal Kenya. *Ecological Entomology* 28 (1): 109 - 118.
- Zhou, J., Yin, L.H. and Wang, C.Z. (2002).** Polydnavirus gene products and their physiological effects on host insects. *Entomological Knowledge* 39 (3): 161 - 165.
- Zhu, H.Q. and Zhao, G. (1988).** Studies on the biology of *Casinaria* sp. (Hym.: Ichneumonidae), a parasite of *Semiothisa cineraria* Bremer et Grey. *Natural Enemies of Insects* 10 (3): 148 - 150.
- Zijp, J.P., Blommers, L., Polesny, F., Muller, W. and Olszak, R.W. (1996).** The extraordinary life history of *Scambus pomorum* (Hymenoptera: Ichneumonidae): parasitoid and predator. International conference on integrated fruit production, at Cedzyna, Poland, 28 August - 2 September 1995. *Bulletin OILB SROP* 19 (4): 212 - 213.
- Zijp, J.P. and Blommers, L.H.M. (2002).** Apple sawfly *Hoplocampa testudinea* (Hym.: Tenthredinidae) and its parasitoid *Lathrolestes ensator* in Dutch apple orchards (Hym.: Ichneumonidae: Ctenopelmatinae). *Journal of Applied Entomology* 126 (6): 265 - 274.
- Zijp, J.P. and Blommers, L.H.M. (2002).** Impact of the parasitoid *Lathrolestes ensator* (Hym.: Ichneumonidae: Ctenopelmatinae) as antagonist of apple sawfly *Hoplocampa testudinea* (Hym.: Tenthredinidae). *Journal of Applied Entomology* 126 (7-8): 366 - 377.
- Zijp, J.P. and Blommers, L.H.M. (2002).** Survival mode between the yearly reproduction periods, and reproductive biology of *Scambus pomorum* (Hymenoptera: Ichneumonidae: Pimplinae), a parasitoid of the apple blossom weevil *Anthonomus pomorum* (Coleoptera: Curculionidae). *Entomologia Generalis* 26 (1): 29 - 46.

- Zizp, J.P. and Blommers, L.H.M. (1992).** *Syrrhizus delusorius* and *Scambus pomorum*, two parasitoids of the apple blossom weevil. Proceedings of the Section Experimental and Applied Entomology of the Netherlands Entomological Society 3: 46 - 50.
- Zlatanova, A.A. (1987).** The codling moth. Zashchita Rastenii Moskva 11: 54.
- Zolubas, P., Gedminas, A. and Shields, K. (2001).** Gypsy moth parasitoids in the declining outbreak in Lithuania. Journal of Applied Entomology 125 (5): 229 - 234.
- Zwakhals, C.J. (1997).** *Trogus lapidator* found in the Netherlands (Hymenoptera: Ichneumonidae: Ichneumoninae). Entomologische Berichten 57 (2): 28.